

Name \_\_\_\_\_

ten-digit Student ID number \_\_\_\_\_

Lecture Time \_\_\_\_\_

Recitation Instructor \_\_\_\_\_

Section Number \_\_\_\_\_

## Instructions:

1. Fill in all the information requested above. On the scantron sheet fill in your name, student ID number, and the section number of your recitation with an extra 0 at the left. See list below. Blacken the correct circles.
2. On the bottom under Test/Quiz Number, write 02 and fill in the little circles.
3. This booklet contains 25 problems, each worth 8 points. The maximum score is 200 points.
4. For each problem mark your answer on the scantron sheet and also circle it in this booklet.
5. Work only on the pages of this booklet.
6. Books, notes, calculators are not to be used on this test.
7. At the end turn in your exam and scantron sheet to your recitation instructor.

TA	Lecture time	Rec. time	Sect. #	TA	Lecture time	Rec. time	Sect. #
Yun Ge	11:30	7:30	0001	Chris Bush	1:30	7:30	0011
		8:30	0002			8:30	0012
Huijie Wang	11:30	9:30	0003	Chi Weng Cheong	1:30	8:30	0031
		10:30	0004	Jing Feng Lau	1:30	9:30	0013
Wei-Nan Lin	11:30	11:30	0005			10:30	0014
		12:30	0006	Raakesh Pankanti	1:30	11:30	0015
Feng Chen	11:30	1:30	0007			12:30	0016
		2:30	0008	Phani Surapaneni	1:30	1:30	0017
Abhijeet Bhalerao	11:30	3:30	0009			2:30	0018
		4:30	0010	Himanshu Markandeya	1:30	3:30	0019
Kevin Mugo	11:30 or 1:30	1:30	0027			4:30	0020

1. The domain of the function  $y = \ln(2^{x/3})$  is
- A. All real numbers.
  - B.  $x > 0$
  - C.  $x > \ln\left(\frac{2}{3}\right)$
  - D.  $x > \ln\left(\frac{3}{2}\right)$
  - E.  $x > -\frac{1}{3}$
2. Express  $\tan(\sin^{-1}(x))$  as an algebraic function of  $x$ .
- A.  $\frac{1}{\sqrt{1-x^2}}$
  - B.  $\frac{1}{\sqrt{x^2-1}}$
  - C.  $\frac{x}{\sqrt{1-x^2}}$
  - D.  $\sqrt{1-x^2}$
  - E.  $\frac{\sqrt{1-x^2}}{x}$

3.  $\lim_{x \rightarrow -2} \frac{\frac{1}{2} + \frac{1}{x}}{2+x} =$

A.  $\infty$   
B.  $-\infty$   
C. 0  
D.  $-\frac{1}{4}$   
E.  $\frac{1}{2}$

4.  $\lim_{x \rightarrow -\infty} \frac{\sqrt{x^2 + 2}}{x} =$

A.  $\infty$   
B.  $-\infty$   
C. 0  
D. 1  
E.  $-1$

5. Find an equation of the tangent line to the curve,  $y = \frac{x}{x+1}$  at the point  $\left(1, \frac{1}{2}\right)$ .

A.  $y = \frac{1}{2}$

B.  $x + 2y = 2$

C.  $x - 2y = 0$

D.  $4y + x = 3$

E.  $4y - x = 1$

6. The radius of a sphere is increasing at a rate of 2 mm/sec. How fast is the volume increasing when the radius is 10 mm? ( $V = \frac{4}{3} \pi r^3$ )

A.  $640\pi \text{ mm}^3/\text{sec}$

B.  $1600\pi \text{ mm}^3/\text{sec}$

C.  $1000\pi \text{ mm}^3/\text{sec}$

D.  $800\pi \text{ mm}^3/\text{sec}$

E.  $1200\pi \text{ mm}^3/\text{sec}$

7. Use a linear approximation (or differentials) to estimate  $\sqrt[3]{994}$ .

- A. 9.95
- B. 9.8
- C. 10.02
- D. 9.99
- E. 9.98

8. A population of bacteria doubles every 2 days. How long will it take for the population to triple?

- A.  $2 \ln\left(\frac{3}{2}\right)$  days
- B.  $2 \frac{\ln 3}{\ln 2}$  days
- C. 3 days
- D. 2.5 days
- E.  $2 \ln 6$  days

9. If  $f(x) = \ln(\sin(x^2))$ , then  $f'(x) =$ 

A.  $\frac{\sin(x^2)}{2x \cos(x^2)}$

B.  $\frac{\cos(x^2)}{2x \sin(x^2)}$

C.  $\frac{2x \cos(x^2)}{\sin(x^2)}$

D.  $\frac{2x \sin(x^2)}{\cos(x^2)}$

E.  $\frac{2x}{\sin(x^2) \cos(x^2)}$

10. If  $f(x) = x^{\sin x}$ , then  $f'(x) =$ 

A.  $x^{\cos x}$

B.  $x^{\sin x} \cos x$

C.  $x^{\sin x} \left( \cos x \ln x + \frac{\sin x}{x} \right)$

D.  $x^{\sin x} (\cos x \ln x)$

E.  $x^{\sin x} \frac{\cos x}{x}$

11. Find an equation of the tangent line to the curve  $(x^2 + y^2)^2 = 4x^2y$  at the point  $(x, y) = (1, 1)$ .

- A.  $y = 1$
- B.  $y = x$
- C.  $y = 2x - 1$
- D.  $y = -x + 2$
- E.  $y = -2x + 3$

12. The absolute maximum value of the function  $f(x) = \frac{x}{x^2 + 1}$  on the interval  $[0, 2]$  is

- A.  $\frac{2}{5}$
- B. 0
- C. 1
- D.  $\frac{1}{2}$
- E.  $\frac{3}{4}$

13. Let  $f$  be a function whose derivative  $f'$  is given by  $f'(x) = (x - 2)^2 x^4(x + 2)^5$ . Then  $f$  has a

- A. local minimum at  $x = 0$
- B. local minimum at  $x = -2$
- C. local maximum at  $x = 0$
- D. local minimum at  $x = 2$
- E. local maximum at  $x = 2$

14.  $\lim_{x \rightarrow \frac{\pi}{2}} (\sec x - \tan x) =$

- A. 0
- B. 1
- C.  $\infty$
- D.  $-1$
- E.  $\frac{1}{2}$

15. A box with square base and open top must have a volume of  $4000 \text{ cm}^3$ . If the cost of the material used is  $\$1/\text{cm}^2$ , the smallest possible cost of the box is

- A. \$500
- B. \$600
- C. \$1000
- D. \$1200
- E. \$2000

16. If  $\int_{-2}^2 f(x)dx = 2$  and  $\int_0^2 f(x)dx = 3$ , then  $\int_{-2}^0 f(x)dx =$

- A. -1
- B. 1
- C. -5
- D. 5
- E. -3

17.  $\int_0^1 (x^2 - \sqrt{x} + 1)dx =$

A.  $-\frac{1}{6}$   
B.  $\frac{5}{6}$   
C.  $\frac{2}{3}$   
D.  $\frac{1}{3}$   
E. 1

18.  $\int_0^{\frac{\pi}{2}} \sin(2x)dx =$

A. 1  
B. -1  
C. 2  
D. -2  
E. 0

19.  $\int_0^1 x^2(x^3 + 1)^{17} dx =$

A.  $\frac{2^{18}}{18}$

B.  $\frac{2^{18}}{54}$

C.  $\frac{2^{18} - 1}{18}$

D.  $\frac{2^{18} - 1}{54}$

E.  $\frac{2^{18} - 1}{3}$

20. If  $g(x) = \int_0^{2x} e^{t^2} dt$ , then  $g'(x) =$

A.  $e^{2x^2}$

B.  $2e^{x^2}$

C.  $2e^{2x^2}$

D.  $2e^{4x^2}$

E.  $e^{4x^2}$

21. The function  $2x^3 - 9x^2 - 24x + 1$  is increasing on

- A.  $(-\infty, -1)$  only
- B.  $(-1, 4)$  only
- C.  $(4, \infty)$  only
- D.  $(-\infty, 1)$  and  $(4, \infty)$
- E.  $(-\infty, -1)$  and  $(4, \infty)$

22. The function  $x^4 - 6x^3 + 12x^2 + 1$  is concave down on


- A.  $(-\infty, 1)$  only
- B.  $(1, 2)$  only
- C.  $(2, \infty)$  only
- D.  $(-\infty, -1)$  and  $(2, \infty)$
- E.  $(-\infty, 1)$  and  $(2, \infty)$

23. The graph of the function  $f$  is given below, and  $\lim_{x \rightarrow 1^+} f(x) = a$ ,  $\lim_{x \rightarrow 1^-} f(x) = b$ .  
 Which of the following is true?

MA 16100

Fall 2008

1. The graph of  $y = f(x)$  is shown below.



A.  $a = 2, b = 1$

B.  $a = 1, b = 2$

C.  $a = \text{dne}, b = \text{dne}$

D.  $a = \text{dne}, b = 2$

E.  $a = 1, b = \text{dne}$

(dne means “does not exist”)

24. The graph of the function  $y = \sin(x)$  is shrunk horizontally by a factor of 2, then translated 1 unit to left. The resulting graph is that of

A.  $y = \sin\left(\frac{1}{2}x + 1\right)$

B.  $y = \sin\left(\frac{1}{2}x - 1\right)$

C.  $y = \sin\left(\frac{1}{2}x + 2\right)$

D.  $y = \sin(2x + 1)$

E.  $y = \sin(2x + 2)$

25. For what value of the constant  $c$  is the function  $f$  continuous on  $(-\infty, \infty)$ ?

$$f(x) = \begin{cases} 2cx^2 + 3x & \text{if } x < 2 \\ x^3 + cx^2 & \text{if } x \geq 2 \end{cases}$$

- A.  $\frac{1}{4}$
- B. 4
- C.  $\frac{1}{2}$
- D. 2
- E. No value of  $c$  makes  $f$  continuous on  $(-\infty, \infty)$