

1. What is the determinant of the following matrix?

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 1 \\ 3 & 4 & 1 & 2 \\ 4 & 1 & 2 & 3 \end{bmatrix}$$

- A. 0
B. 8
C. 55
D. 120
E. 160
2. If $\det \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix} = 4$, then $\det \begin{bmatrix} a_1 & a_2 & 4a_3 - 2a_2 \\ b_1 & b_2 & 4b_3 - 2b_2 \\ \frac{1}{2}c_1 & \frac{1}{2}c_2 & 2c_3 - c_2 \end{bmatrix} =$
- A. 8
B. 6
C. 4
D. 2
E. 1
3. Let A be an $n \times n$ matrix. Which of the following statements must be true?
- (i) If $A^T = -A$, then $\det A$ must be zero.
(ii) If $A^2 = A$, then A must be the identity matrix or the zero matrix.
(iii) If $\det A \neq 0$, then the homogeneous system $A\mathbf{x} = \mathbf{0}$ only has the trivial solution.
- A. None
B. (iii) only
C. (i) and (iii) only
D. (ii) and (iii) only
E. (i), (ii) and (iii)

4. Which of the following vectors in R_3 is a linear combination of

$$v_1 = [4 \ 2 \ -3], v_2 = [2 \ 1 \ -2], v_3 = [-2 \ -1 \ 4]?$$

- A. $[1 \ 0 \ 0]$
B. $[1 \ 1 \ 1]$
C. $[-2 \ 2 \ 3]$
D. $[6 \ 3 \ 7]$
E. None of the above.
5. Which of the following sets of 2×2 matrices are vector spaces? (Here \oplus and \odot are the usual addition and scalar multiplication of matrices.)
- (i) {all matrices $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ such that $a + b = 3c - d$.}
(ii) {all 2×2 matrices A such that $A\mathbf{x} = \mathbf{0}$ has a nontrivial solution.}
(iii) {all 2×2 matrices A such that $A\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ is consistent.}
- A. (i) and (ii)
B. (ii) only
C. (ii) and (iii)
D. (i) only
E. All of them are vector spaces.
6. Let P_3 be the set of all polynomials of degree 3 or less. Which of the following subsets are subspaces of P_3 ? (Here \oplus and \odot are the standard addition and scalar multiplication.)
- (i) {all polynomials $p(x)$ such that $p(1) \neq 0$ }
(ii) {all polynomials $p(x)$ such that $p(x) = p(-x)$.}
(iii) {all polynomials $p(x)$ with $p(0) = p(1)$.}
- A. (i) and (ii)
B. (ii) only
C. (ii) and (iii)
D. (i) and (iii)
E. All of them are vector spaces.

7. Let $A = \begin{bmatrix} 2 & 0 & 1 \\ 0 & 1 & t \\ 2 & 0 & 1 \\ 0 & 1 & t \end{bmatrix}$. Choose a value of t so that the null space of A has dimension 0.

- A. 0
- B. 1
- C. -1
- D. 2
- E. No such value of t exists.

8. Let A be a 3×5 matrix with rank 3. Which of the following statements is true?

- A. A consistent linear system $A\mathbf{x} = \mathbf{b}$ must have a unique solution.
- B. The null space of A has dimension 3.
- C. The columns of A form a basis for the column space of A .
- D. The system $A\mathbf{x} = \mathbf{b}$ is consistent for every \mathbf{b} in \mathbb{R}^3 .
- E. The rows of A form a linearly dependent set.

9. Choose a value of a so that the vector $\begin{bmatrix} -9 \\ 0 \\ 1 \end{bmatrix}$ is orthogonal to the vector $\begin{bmatrix} a \\ 1 \\ a^3 \end{bmatrix}$.

- A. 1
- B. 2
- C. 3
- D. 4
- E. None of the above

10. Let $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$. Then $\begin{bmatrix} 1 \\ -i \end{bmatrix}$

- A. is an eigenvector with eigenvalue 1.
- B. is an eigenvector with eigenvalue -1 .
- C. is an eigenvector with eigenvalue i .
- D. is an eigenvector with eigenvalue $-i$.
- E. is not an eigenvector.

11. Consider the following differential equation:

$$dx/dt = x + 5y$$

$$dy/dt = 5x + y$$

Which of the following is a solution:

- A. $x = 3e^{3t} + 2e^{-2t}$ and $y = 2e^{3t} - 2e^{-2t}$
- B. $x = 3e^{4t} + 2e^{-4t}$ and $y = 2e^{6t} - 3e^{-t}$.
- C. $x = 2e^{6t} + 2e^{-4t}$ and $y = 2e^{6t} - 2e^{-4t}$.
- D. $x = 3e^{3t} + 2e^{2t}$ and $y = 2e^{3t} - 2e^{2t}$.
- E. $x = 8e^t + 4e^{-3t}$ and $y = 9e^t - 4e^{-3t}$.

12. Suppose the vector $v = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$ is orthogonal to $\begin{bmatrix} 1 \\ 0 \\ -1 \\ 1 \end{bmatrix}$, $\begin{bmatrix} 1 \\ -1 \\ 0 \\ -1 \end{bmatrix}$ and

$\begin{bmatrix} 2 \\ -1 \\ -1 \\ 1 \end{bmatrix}$. Then which of the following statement is always correct:

- A. $a = 0, b = c = -d$
- B. $b = 0, a = b = -d$
- C. $c = a = 0, a = b = -c$
- D. $d = 0, a = b = c$
- E. $a = b = 0, c = 2d + 1$

13. Let $C[-\pi, \pi]$ be the real vector space of continuous functions defined on $[-\pi, \pi]$. Define an inner product on $C[-\pi, \pi]$ by

$$\langle f, g \rangle = \int_{-\pi}^{\pi} f(t)g(t)dt.$$

Then which of the following set of vectors are orthogonal?

- A. $1, t, t^2$
 - B. $\sin^2 t, 1, \cos t$
 - C. $1, e^t, e^{2t}$
 - D. $1, t, t^2 - \frac{\pi^2}{3}$,
 - E. None of the above
14. Let A be an $n \times n$ matrix, which of the following statements is FALSE?
- A. If A is a symmetric matrix, then A^T is also symmetric.
 - B. The product AA^T is always symmetric.
 - C. If A is skew symmetric, then A^3 is symmetric.
 - D. If A is symmetric, then $A + A^2$ is symmetric.
 - E. The sum $A + A^T$ is always symmetric.
15. A and B are $n \times n$ invertible matrices, which of the following statements is FALSE?
- A. $(A^2)^{-1} = (A^{-1})^2$
 - B. $(A^{-1})^T = (A^T)^{-1}$.
 - C. $(AB^{-1})^{-1} = BA^{-1}$.
 - D. $(A + B^{-1})^{-1} = A^{-1} + B$.
 - E. $(aA)^{-1} = \frac{1}{a}A^{-1}$ for any nonzero real number a .

16. Consider the linear system

$$\begin{cases} x + 2y + 3z = a \\ 2x - y + z = b \\ 3x + y + 4z = c \end{cases}$$

Under which condition will this system be consistent?

- A. $a = c - 2b$
- B. $a = c - b$.
- C. $b = a + c$.
- D. $b = a - 2c$.
- E. $c = 2a + b$.

17. Which of the following statements is FALSE?

- A. The rows of any invertible $n \times n$ matrix span \mathbb{R}^n .
- B. The rows of any orthogonal $n \times n$ matrix span \mathbb{R}^n .
- C. The rows of any elementary $n \times n$ matrix span \mathbb{R}^n .
- D. The rows of any symmetric $n \times n$ matrix span \mathbb{R}^n .
- E. The rows of any nonsingular $n \times n$ matrix span \mathbb{R}^n .

18. Let $A = \begin{bmatrix} 2 & 3 & 4 \\ 4 & 5 & 6 \\ 1 & 2 & 3 \\ 1 & 0 & 1 \end{bmatrix}$. Which of the following collections of vectors is linearly independent?

- A. The rows of the matrix A .
- B. The rows of the matrix A^T .
- C. The rows of the matrix $A \cdot A^T$.
- D. The first three rows of A .
- E. None of the above.

19. In this problem, A is a 5×8 matrix. Find the TRUE statement.
- A. If $\text{rref}(A)$ has five leading 1's then the columns of A are linearly independent.
 - B. If the number of leading 1's of $\text{rref}(A)$ equals three, then one can pick three columns of A that span the column space of A .
 - C. If columns 1,3 and 8 of $\text{rref}(A)$ have no leading 1 then A must have five linearly independent rows.
 - D. If columns 1, 3, 4, 5 and 8 of $\text{rref}(A)$ are exactly the ones that have no leading 1 then there are five linearly independent rows in A .
 - E. None of the above.

20. What are the eigenvalues of the matrix $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}$?

- A. 1, 2.
- B. 1, 2, 3.
- C. $\pm 1, 2$.
- D. $\pm 1, 2, 3$.
- E. None of the above.

21. Which of the following matrices is NOT diagonalizable?

- A. $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$.
- B. $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$.
- C. $\begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$.
- D. $\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$.
- E. $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$.

22. For an $n \times n$ real matrix A , which of the following statements are true?
- i) If $A^T = A$, then all eigenvalues of A are real.
 - ii) If A is an orthogonal matrix, then the columns of A form an orthonormal set.
 - iii) If all eigenvalues of A are equal to 1, then A is similar to the identity matrix I_n .

- A. i) only.
- B. ii) only.
- C. i) and ii) only .
- D. i), ii), iii).
- E. None of the above.

23. Assume A and B are $n \times n$ symmetric matrices. Which of the following statements are always true?

- (i) A^2 is symmetric.
- (ii) AB is symmetric.
- (iii) ABA is symmetric.

- A. none of these
- B. (i) only
- C. (ii) only
- D. (i) and (iii) only.
- E. (i), (ii) and (iii)

24. Let $A = \begin{bmatrix} 1 & 0 & 2 \\ -1 & 3 & 5 \\ 0 & 4 & -2 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 0 & 3 \end{bmatrix}$. Compute the $(2, 3)$ -entry of $2A[B + A^T]$.

- A. 4
- B. 8
- C. 26
- D. 52
- E. 104

25. Let W denote the vector space spanned by the vectors

$$\mathbf{u}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 2 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 2 \end{bmatrix},$$

and let $\mathbf{v} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ -1 \end{bmatrix}$. Find the distance from \mathbf{v} to W .

- A. 0
- B. 1
- C. 2
- D. $\sqrt{2}$
- E. 4