

Martingale transforms and their projection operators on manifolds

Rodrigo Bañuelos*, Fabrice Baudoin†
Department of Mathematics
Purdue University
West Lafayette, 47906

September 29, 2011

Abstract

We prove the boundedness on L^p , $1 < p < \infty$, of operators on manifolds which arise by taking conditional expectation of transformations of stochastic integrals. These operators include various classical operators such as second order Riesz transforms and operators of Laplace transform-type.

Contents

1	Introduction	1
2	Martingale inequalities	3
3	Proof of Theorem 1.1 and some refinements	7
3.1	An estimate of the constant in (1.3)	12
4	Applications	14
4.1	Multipliers of Laplace transform-type for Schrödinger operators on manifolds . .	14
4.2	Second order Riesz transforms on Lie groups of compact type	16

1 Introduction

The classical martingale inequalities of Burkholder and Gundy [8] play a fundamental role in many areas of probability and its applications. These inequalities have their roots in the celebrated 1966 martingale transforms inequality of Burkholder [6]. In 1984 [7], Burkholder obtained

*Supported in part by NSF Grant #0603701-DMS

†Supported in part by NSF Grant 0907326-DMS

the sharp constants in his 1966 martingale inequalities. In recent years, these sharp inequalities have had many applications to the study of basic singular integrals and Fourier multipliers on Euclidean space \mathbb{R}^d with the Lebesgue measure (the ordinary Laplacian case) and with Gaussian measure (the Ornstein–Uhlenbeck case). For an account of some of this literature we refer the reader to the overview article [2]. The purpose of this paper is to show that these martingale transform techniques apply to wide range of operators on manifolds, including multipliers arising from Schrödinger operators and Riesz transforms on Lie groups. For the Laplacian, and other self-adjoint diffusions without a zero order term, Burkholder’s sharp inequalities can be applied and our L^p bounds are exactly as those on \mathbb{R}^d . However, in the case of Schrödinger operators the sharp Burkholder inequalities do not apply in any direct way and in this case we obtain our results by proving a version of the Burkholder-Gundy inequalities with an exponential weight which does not produce best constants. Our results in this paper (see Remark 2.1 below) also correct a gap contained in [21], [22] and [23]. (See also [20] for details on the correct formulation of some of the results in those papers.)

To introduce our operators and state our results, let \mathbb{M} be a smooth manifold endowed with a smooth measure μ . Let X_1, \dots, X_d be locally Lipschitz vector fields defined on \mathbb{M} . We consider the Schrödinger operator,

$$L = -\frac{1}{2} \sum_{i=1}^d X_i^* X_i + V,$$

where X_i^* denotes the formal adjoint of X_i with respect to μ and where $V : \mathbb{M} \rightarrow \mathbb{R}$ is a non-positive smooth function, often referred to it in this paper as a *potential*. We assume that L is essentially self-adjoint with respect to μ on the space $\mathcal{C}_0^\infty(\mathbb{M})$ of smooth and compactly supported functions and that the solution of the Stratonovitch stochastic differential equation associated with L is non-explosive. We denote by $(P_t)_{t \geq 0}$ the heat semigroup with generator L . We can write $L = \frac{1}{2} \sum_{i=1}^d X_i^2 + X_0 + V$, for some locally Lipschitz vector field X_0 .

Let $A_{ij} : [0, +\infty) \times \mathbb{M} \rightarrow \mathbb{R}$, $1 \leq i, j \leq d$ be bounded and smooth real valued functions and let $A(t, x) = (A_{ij})$ be the $d \times d$ matrix with A_{ij} entries. Set

$$\|A\| = \| \|A(t, x)\| \|_{L^\infty([0, +\infty) \times \mathbb{M})},$$

where $|A(t, x)|$ is the usual quadratic norm of the $d \times d$ matrix $A(t, x)$. That is,

$$|A(t, x)| = \sup\{|A(t, x)\xi|; \xi \in \mathbb{R}^d, |\xi| = 1\}.$$

Our goal in this paper is to study the continuity on $L_\mu^p(\mathbb{M})$ for $1 < p < \infty$ of the operator

$$(1.1) \quad \mathcal{S}_A f = \sum_{i,j=1}^d \int_0^\infty P_t X_i^* A_{ij}(t) X_j P_t f dt$$

and to obtain precise information on the size of their norms. The following two remarks shed some light on the structure of these operators in two important special cases.

Remark 1.1. If $L = -\frac{1}{2} \sum_{i=1}^d X_i^* X_i + V$ is subelliptic, then the semigroup P_t admits a smooth symmetric kernel $p(t, x, y)$ (see [5], [17]) and in that case, it is easily seen that

$$\mathcal{S}_A f(x) = \int_{\mathbb{M}} K(x, z) f(z) d\mu(z),$$

with

$$K(x, z) = \sum_{i,j=1}^d \int_0^{+\infty} \int_{\mathbb{M}} A_{ij}(t, y) X_i^y p(t, y, x) X_j^y p(t, y, z) d\mu(y) dt.$$

Remark 1.2. Let (\mathbb{M}, g) be a complete Riemannian manifold. In this framework the Laplace-Beltrami operator Δ is essentially self-adjoint on $C_0^\infty(\mathbb{M})$. We may then consider the Schrödinger operator

$$L = -\frac{1}{2} \Delta + V$$

where, as above, V is a non-positive smooth potential. In this case, the operator \mathcal{S}_A can be written as

$$\mathcal{S}_A f = \int_0^\infty P_t \operatorname{div}(A(t) \nabla P_t f) dt.$$

For $1 < p < \infty$ let p^* denote the maximum of p and q , where $\frac{1}{p} + \frac{1}{q} = 1$. Thus $p^* = \max\{p, \frac{p}{p-1}\}$ and

$$(1.2) \quad p^* - 1 = \begin{cases} \frac{1}{p-1}, & 1 < p \leq 2, \\ p - 1, & 2 \leq p < \infty. \end{cases}$$

The following is the main result of this paper.

Theorem 1.1. For any $1 < p < \infty$ there is a constant C_p depending only on p such that for every $f \in L^p_\mu(\mathbb{M})$,

$$(1.3) \quad \|\mathcal{S}_A f\| \leq C_p \|A\| \|f\|_p.$$

If the potential $V \equiv 0$, then

$$(1.4) \quad \|\mathcal{S}_A f\|_p \leq (p^* - 1) \|A\| \|f\|_p,$$

and this bound is sharp.

As we shall see below, these operators include the multipliers of Laplace transform-type and second order Riesz transforms on Lie groups of compact type. The fact that the bound in (1.4) is best possible follows from the fact that the best constant in the L^p inequality for second order Riesz transforms $R_j R_k$ on \mathbb{R}^d is $p^* - 1$; see §4.2 below and [13] and [3].

The paper is organized as follows. In §2, we recall various versions of Burkholder's sharp inequalities and prove a version of the Burkholder-Gundy inequality needed for (1.3). The proof of Theorem 1.1 is given in §3 where we also give an explicit bound for the constant in (1.3). In §4, we give some concrete examples of our operators.

2 Martingale inequalities

In this section we recall the sharp martingale inequalities of Burkholder and prove a version of the Burkholder-Gundy inequalities used in the proof of (1.3) in Theorem 1.1.

Let $f = \{f_n, n \geq 0\}$ be a martingale with different sequence $df = \{df_k, k \geq 0\}$, where $df_k = f_k - f_{k-1}$ for $k \geq 1$ and $df_0 = f_0$. Given a predictable sequence of random variables $\{v_k, k \geq 0\}$ uniformly bounded for all k the martingale difference sequence $\{v_k df_k, k \geq 0\}$ generates a new martingale called the *martingale transform* of f and denoted by $v * f$. We set $\|f\|_p = \sup_{n \geq 0} \|f_n\|_p$ for $0 < p < \infty$. Burkholder's 1966 result [6] asserts that the operator $f \rightarrow v * f = g$ is bounded on L^p for all $1 < p < \infty$. In his 1984 seminal paper [7] Burkholder determined the norm of this operator by proving the following result.

Theorem 2.1. *Let $f = \{f_n, n \geq 0\}$ be a martingale and let $g = v * f$ be its martingale transform by the predictable sequence $v = \{v_k, k \geq 0\}$ with v_k taking values in $[-1, 1]$ for all k . Then*

$$(2.1) \quad \|g\|_p \leq (p^* - 1)\|f\|_p, \quad 1 < p < \infty,$$

and the constant $p^* - 1$ is best possible.

In [9], K.P. Choi used the techniques of Burkholder to identify the best constant in the martingale transforms where the predictable sequence v takes values in $[0, 1]$ instead of $[-1, 1]$. While Choi's constant is not as explicit as the $p^* - 1$ constant of Burkholder, one does have a lot of information on it.

Theorem 2.2. *Let $f = \{f_n, n \geq 0\}$ be a real-valued martingale and let $g = v * f$ be its martingale transform by a predictable sequence $v = \{v_k, k \geq 0\}$ with values in $[0, 1]$ for all k . Then*

$$(2.2) \quad \|g\|_p \leq c_p \|f\|_p, \quad 1 < p < \infty,$$

with the best constant c_p satisfying

$$c_p = \frac{p}{2} + \frac{1}{2} \log \left(\frac{1 + e^{-2}}{2} \right) + \frac{\alpha_2}{p} + \dots$$

where

$$\alpha_2 = \left[\log \left(\frac{1 + e^{-2}}{2} \right) \right]^2 + \frac{1}{2} \log \left(\frac{1 + e^{-2}}{2} \right) - 2 \left(\frac{e^{-2}}{1 + e^{-2}} \right)^2.$$

Motivated by Theorems 2.1 and 2.2 the following definition was introduced in [3].

Definition 2.1. *Let $-\infty < b < B < \infty$ and $1 < p < \infty$ be given and fixed. We define $C_{p,b,B}$ as the least positive number C such that for any real-valued martingale f and for any transform $g = v * f$ of f by a predictable sequence $v = \{v_k, k \geq 0\}$ with values in $[b, B]$, we have*

$$(2.3) \quad \|g\|_p \leq C \|f\|_p.$$

Thus, for example, $C_{p,-a,a} = a(p^* - 1)$ by Burkholder's Theorem 2.1 and $C_{p,0,a} = a c_p$ by Choi's Theorem 2.2. It is also the case that for any b, B as above,

$$(2.4) \quad \max \left\{ \left(\frac{B-b}{2} \right) (p^* - 1), \max\{|B|, |b|\} \right\} \leq C_{p,b,B} \leq \max\{B, |b|\} (p^* - 1)$$

For the applications to the above operators we will need versions of these inequalities for continuous-time martingales. Suppose that $(\Omega, \mathcal{F}, \mathbb{P})$ is a complete probability space, filtered by $(\mathcal{F}_t)_{t \geq 0}$, a nondecreasing and right-continuous family of sub- σ -fields of \mathcal{F} . Assume, as usual, that \mathcal{F}_0 contains all the events of probability 0. Let X, Y be adapted, real valued martingales which have right-continuous paths with left-limits (r.c.l.l.). Denote by $[X, X]$ the quadratic variation process of X : we refer the reader to Dellacherie and Meyer [10] for details. Following [4] and [27], we say that Y is differentially subordinate to X if the process $|Y_0| \leq |X_0|$ and $([X, X]_t - [Y, Y]_t)_{t \geq 0}$ is nondecreasing and nonnegative as a function of t . We have the following extension of the Burkholder inequalities proved in [4] for continuous-path martingales and in [27] in the general case. Set $\|X\|_p = \sup_{t \geq 0} \|X_t\|$, $0 < p < \infty$.

Theorem 2.3. *If Y is differentially subordinate to X , then*

$$(2.5) \quad \|Y\|_p \leq (p^* - 1) \|X\|_p, \quad 1 < p < \infty,$$

and the inequality is sharp.

The case of non-symmetric multipliers is covered by the following result proved in [3].

Theorem 2.4. *Suppose $-\infty < b < B < \infty$ and X_t, Y_t are two real valued martingales with which have right-continuous paths with left-limits with $|Y_0| \leq |X_0|$ and which satisfy*

$$(2.6) \quad \left[\frac{B-b}{2} X, \frac{B-b}{2} X \right]_t - \left[Y - \frac{b+B}{2} X, Y - \frac{b+B}{2} X \right]_t \geq 0$$

and nondecreasing for all $t \geq 0$ (differential subordination). Then

$$\|Y\|_p \leq C_{p,b,B} \|X\|_p, \quad 1 < p < \infty,$$

and the inequality is sharp.

Note that when $B = a > 0$ and $b = -a$, we have the case of Theorem 2.3. As we shall see, (2.5) will give the bound in (1.4) and (2.6) will give some extensions.

For the general case when $V \neq 0$ (for the inequality (1.3)) none of the above results apply and we shall need a variation of the Burkholder-Gundy inequalities. While this bound is not sharp, it applies to a wide class of processes and can even be used to study operators on manifolds acting on forms. We shall comment more on this a little later. We start by recalling the following domination inequality due to Lenglart [18] (See also Revuz-Yor [25], p.162-163).

Proposition 2.1. (Lenglart) *Let $(N_t)_{t \geq 0}$ be a positive adapted right-continuous process and $(A_t)_{t \geq 0}$ be an increasing process. Assume that for every bounded stopping time τ ,*

$$\mathbb{E}(N_\tau) \leq \mathbb{E}(A_\tau).$$

Then, for every $k \in (0, 1)$,

$$\mathbb{E} \left(\left(\sup_{0 \leq t \leq T} N_t \right)^k \right) \leq \frac{2-k}{1-k} \mathbb{E} \left(A_T^k \right).$$

We shall use this lemma to prove the following

Theorem 2.5. *Let $T > 0$ and $(M_t)_{0 \leq t \leq T}$ be a continuous local martingale. Consider the process*

$$Z_t = e^{\int_0^t V_s ds} \int_0^t e^{-\int_0^s V_u du} dM_s,$$

where $(V_t)_{0 \leq t \leq T}$ is a non positive adapted and continuous process. For every $0 < p < \infty$, there is a universal constant C_p , independent of T , $(M_t)_{0 \leq t \leq T}$ and $(V_t)_{0 \leq t \leq T}$ such that

$$\mathbb{E} \left(\left(\sup_{0 \leq t \leq T} |Z_t| \right)^p \right) \leq C_p \mathbb{E} \left(\langle M \rangle_T^{\frac{p}{2}} \right).$$

Proof. By stopping it is enough to prove the result for bounded M . Let $q \geq 2$. From Itô's formula we have

$$dZ_t = Z_t V_t dt + dM_t$$

and

$$\begin{aligned} d|Z_t|^q &= q|Z_t|^{q-1} \mathbf{sgn}(Z_t) dZ_t + \frac{1}{2} q(q-1) |Z_t|^{q-2} d \langle M \rangle_t \\ &= q|Z_t|^q V_t dt + q \mathbf{sgn}(Z_t) |Z_t|^{q-1} dM_t + \frac{1}{2} q(q-1) |Z_t|^{q-2} d \langle M \rangle_t. \end{aligned}$$

Since $V_t \leq 0$, as a consequence of the Doob's optional sampling theorem, we get that for every bounded stopping time τ ,

$$\mathbb{E} (|Z_\tau|^q) \leq \frac{1}{2} q(q-1) \mathbb{E} \left(\int_0^\tau |Z_t|^{q-2} d \langle M \rangle_t \right).$$

From the Lenglart's domination inequality, we deduce then that for every $k \in (0, 1)$,

$$\mathbb{E} \left(\left(\sup_{0 \leq t \leq T} |Z_t|^q \right)^k \right) \leq \frac{2-k}{1-k} \left(\frac{1}{2} q(q-1) \right)^k \mathbb{E} \left(\left(\int_0^T |Z_t|^{q-2} d \langle M \rangle_t \right)^k \right).$$

We now bound

$$\begin{aligned} \mathbb{E} \left(\left(\int_0^T |Z_t|^{q-2} d \langle M \rangle_t \right)^k \right) &\leq \mathbb{E} \left(\left(\sup_{0 \leq t \leq T} |Z_t| \right)^{k(q-2)} \left(\int_0^T d \langle M \rangle_t \right)^k \right) \\ &\leq \mathbb{E} \left(\left(\sup_{0 \leq t \leq T} |Z_t| \right)^{kq} \right)^{1-\frac{2}{q}} \mathbb{E} \left(\langle M \rangle_T^{\frac{kq}{2}} \right)^{\frac{2}{q}}. \end{aligned}$$

As a consequence, we obtain

$$\mathbb{E} \left(\left(\sup_{0 \leq t \leq T} |Z_t|^q \right)^k \right) \leq \frac{2-k}{1-k} \left(\frac{1}{2} q(q-1) \right)^k \mathbb{E} \left(\left(\sup_{0 \leq t \leq T} |Z_t| \right)^{kq} \right)^{1-\frac{2}{q}} \mathbb{E} \left(\langle M \rangle_T^{\frac{kq}{2}} \right)^{\frac{2}{q}}.$$

Letting $p = qk$ yields the claimed result. \square

We should note here that in the above proof the fixed time T can be replaced by a stopping time. More interesting and useful for applications is the fact that this result admits the following multidimensional generalization whose proof identical to the above proof and which we leave to the interested reader.

Theorem 2.6. *Let $T > 0$ and $(M_t)_{0 \leq t \leq T}$ be an \mathbb{R}^d valued continuous local martingale. Consider the solution of the matrix equation*

$$d\mathcal{M}_t = \mathcal{V}_t \mathcal{M}_t dt, \quad \mathcal{M}_0 = \mathbf{Id},$$

where $(\mathcal{V}_t)_{0 \leq t \leq T}$ is an adapted and continuous process taking values in the set of symmetric and non positive $d \times d$ matrices. Consider the process

$$Z_t = \mathcal{M}_t \int_0^t \mathcal{M}_s^{-1} dM_s.$$

For every $0 < p < \infty$, there is a universal constant C_p , that is independent from $d, T, (M_t)_{0 \leq t \leq T}$ and $(\mathcal{V}_t)_{0 \leq t \leq T}$ such that

$$(2.7) \quad \mathbb{E} \left(\left(\sup_{0 \leq t \leq T} \|Z_t\| \right)^p \right) \leq C_p \mathbb{E} \left(\|\langle M \rangle_T\|^{\frac{p}{2}} \right).$$

Remark 2.1. *Theorem 2.6 may be used to correct a gap contained in the papers [21], [22], [23] of X. D. Li. In these papers the author considers quantities like $\mathcal{M}_t \int_0^t \mathcal{M}_s^{-1} dM_s$ but writes them as $\int_0^t \mathcal{M}_t \mathcal{M}_s^{-1} dM_s$ to give L^p estimates using the classical Burkholder-Gundy inequality or even explicit expressions for the constants using the Burkholder bounds of Theorem 2.3. This, however, is not possible due to the non-adaptedness of the process $\mathcal{M}_t \mathcal{M}_s^{-1}$ which prevents us from bringing the \mathcal{M}_t inside the integral. Therefore, in those papers, several of the explicit constants given there involving $(p^* - 1)$ need to be replaced with less precise constants C_p depending only on p . The necessary changes and correct formulations for Li's results are contained in [20].*

3 Proof of Theorem 1.1 and some refinements

Consider the Schrödinger operator $L = \frac{1}{2} \sum_{i=1}^d X_i^2 + X_0 + V$. The diffusion $(Y_t)_{t \geq 0}$ with generator $\frac{1}{2} \sum_{i=1}^d X_i^2 + X_0$ can then be constructed via the Stratonovitch stochastic differential equation

$$dY_t = X_0(Y_t)dt + \sum_{i=1}^d X_i(Y_t) \circ dB_t^i,$$

which we assume non-explosive. In the sequel we shall denote by $(Y_t)_{t \geq 0}$ the solution of this equation started with an initial *distribution* μ . We will use E_x to denote the expectation associated with the process Y starting at x and E to denote the expectation of the process starting with μ . (See [2] for some literature related to the possible intricacies associated to the process starting with the possibly infinite measure μ .)

Let us recall that by the Feynman-Kac formula, the semigroup P_t acting on $f \in C_0^\infty(\mathbb{M})$ can be written as

$$P_t f(x) = E_x \left(e^{\int_0^t V(Y_s) ds} f(Y_t) \right).$$

Let us fix $T > 0$ in what follows. We first consider the relevant martingales associated to our operators. We have

Lemma 3.1. *Let $f \in C_0^\infty(\mathbb{M})$. The process $\left(e^{\int_0^t V(Y_s) ds} (P_{T-t} f)(Y_t) \right)_{0 \leq t \leq T}$ is a martingale and we have*

$$e^{\int_0^T V(Y_s) ds} f(Y_T) = (P_T f)(Y_0) + \sum_{i=1}^d \int_0^T e^{\int_0^t V(Y_s) ds} (X_i P_{T-t} f)(Y_t) dB_t^i.$$

The proof of the lemma is clear. For $0 \leq t \leq T$, set

$$Z_t^1 = e^{\int_0^t V(Y_s) ds}, \quad Z_t^2 = (P_{T-t} f)(Y_t)$$

and apply the Itô formula.

The next expression provides the probabilistic connection to our operators. For $f \in C_0^\infty(\mathbb{M})$, $0 < T < \infty$, set

$$\mathcal{S}_A^T f(x) = \mathbb{E} \left(e^{\int_0^T V(Y_s) ds} \sum_{i,j=1}^d \int_0^T e^{-\int_0^t V(Y_s) ds} A_{ij}(T-t, Y_t) (X_j P_{T-t} f)(Y_t) dB_t^i \mid Y_T = x \right).$$

If g is another function in $C_0^\infty(\mathbb{M})$, the Itô isometry and the lemma give

$$\begin{aligned} \int_{\mathbb{M}} (\mathcal{S}_A^T f) g d\mu &= \mathbb{E} \left(e^{\int_0^T V(Y_s) ds} \sum_{i,j=1}^d \int_0^T e^{-\int_0^t V(Y_s) ds} A_{ij}(T-t, Y_t) (X_j P_{T-t} f)(Y_t) dB_t^i g(Y_T) \right) \\ &= \sum_{i,j=1}^d \mathbb{E} \left(e^{\int_0^T V(Y_s) ds} g(X_T) \int_0^T e^{-\int_0^t V(Y_s) ds} A_{ij}(T-t, Y_t) (X_j P_{T-t} f)(Y_t) dB_t^i \right) \\ &= \sum_{i,j=1}^d \mathbb{E} \left(\int_0^T A_{ij}(T-t, Y_t) (X_i P_{T-t} g)(Y_t) (X_j P_{T-t} f)(Y_t) dt \right) \\ &= \sum_{i,j=1}^d \int_0^T \int_{\mathbb{M}} A_{ij}(t) (X_i P_t g)(X_j P_t f) d\mu dt. \end{aligned}$$

Therefore, we have

$$(3.1) \quad \mathcal{S}_A^T f = \sum_{i,j=1}^d \int_0^T (P_t X_i^* A_{ij}(t) X_j P_t f) dt.$$

Lemma 3.2. *For any $1 < p < \infty$, there is a constant C_p depending only on p such that for every $f \in L_\mu^p(\mathbb{M})$,*

$$(3.2) \quad \|\mathcal{S}_A^T f\| \leq C_p \|A\| \|f\|_p.$$

If the potential $V \equiv 0$, then

$$(3.3) \quad \|\mathcal{S}_A^T f\| \leq (p^* - 1) \|A\| \|f\|_p.$$

In particular these constants do not depend on T .

Proof. We first observe that if $V \equiv 0$ then the martingale

$$\sum_{i,j=1}^d \int_0^t A_{ij}(T-t, Y_t) (X_j P_{T-t} f)(Y_t) dB_t^i$$

is differentially subordinate to $\|A\| P_{T-t} f(Y_t)$. It follows from the contraction of the conditional expectation on L^p , $1 < p < \infty$, and Theorem 2.3, that

$$\|\mathcal{S}_A^T f\| \leq (p^* - 1) \|A\| \|f\|_p,$$

which is the estimate in (3.3).

We now deal with the case of $V \neq 0$. We first prove an L^p estimates for

$$\int_0^T \sum_{i=1}^d (X_i P_{T-t} f)^2(Y_t) dt$$

which is the quadratic variation of the martingale appearing in Lemma 3.1. Using Itô's formula with $0 \leq t \leq T$ for $(P_{T-t} f)(Y_t)^2$, we obtain

$$\begin{aligned} f(Y_T)^2 &= P_T f(Y_0)^2 + \int_0^T \left(\frac{1}{2} \sum_{i=1}^d X_i^2 + X_0 + \frac{\partial}{\partial t} \right) (P_{T-t} f)^2(Y_t) dt \\ &\quad + \sum_{i=1}^d \int_0^T X_i (P_{T-t} f)^2(Y_t) dB_t^i. \end{aligned}$$

Therefore

$$\mathbb{E} \left(\int_0^T \left(\frac{1}{2} \sum_{i=1}^d X_i^2 + X_0 + \frac{\partial}{\partial t} \right) (P_{T-t} f)^2(Y_t) dt \right) \leq \|f\|_2^2.$$

We now compute

$$\begin{aligned}
\left(\frac{1}{2}\sum_{i=1}^d X_i^2 + X_0 + \frac{\partial}{\partial t}\right) (P_{T-t}f)^2 &= 2(P_{T-t}f) \left(\frac{1}{2}\sum_{i=1}^d X_i^2 + X_0\right) P_{T-t}f + \sum_{i=1}^d (X_i P_{T-t}f)^2 \\
&\quad - 2(P_{T-t}f) \left(\frac{1}{2}\sum_{i=1}^d X_i^2 + X_0 + V\right) P_{T-t}f \\
&= \sum_{i=1}^d (X_i P_{T-t}f)^2 - 2V(P_{T-t}f)^2 \\
&\geq \sum_{i=1}^d (X_i P_{T-t}f)^2.
\end{aligned}$$

This implies that

$$\mathbb{E} \left(\int_0^T \sum_{i=1}^d (X_i P_{T-t}f)^2(Y_t) dt \right) \leq \mathbb{E} \left(\int_0^T \left(\frac{1}{2} \sum_{i=1}^d X_i^2 + X_0 + \frac{\partial}{\partial t} \right) (P_{T-t}f)^2(Y_t) dt \right) \leq \|f\|_2^2.$$

Combining this with Theorem 2.5 and again with the fact that the conditional expectation is a contraction on L^2 , we obtain

$$\|\mathcal{S}_A^T f\|_2 \leq \|A\| \|f\|_2,$$

which proves the result for $p = 2$.

Assume now that $2 < p < \infty$. The above computations show that $(P_{T-t}f)^2(Y_t)$ is a submartingale. From the Lenglart-Lépingle-Pratelli estimate and Doob's maximal inequality, we deduce that

$$\begin{aligned}
\mathbb{E} \left(\left(\int_0^T \left(\frac{1}{2} \sum_{i=1}^d X_i^2 + X_0 + \frac{\partial}{\partial t} \right) (P_{T-t}f)^2(Y_t) dt \right)^{\frac{p}{2}} \right) &\leq p^{p/2} \mathbb{E} \left(\sup_{0 \leq t \leq T} ((P_{T-t}f)^2(Y_t))^{p/2} \right) \\
&\leq p^{p/2} \left(\frac{p}{p-2} \right)^{p/2} \mathbb{E}(f(Y_T)^p) \\
&\leq p^{p/2} \left(\frac{p}{p-2} \right)^{p/2} \|f\|_p^p.
\end{aligned}$$

We conclude

$$\begin{aligned}
\mathbb{E} \left(\left(\int_0^T \sum_{i=1}^d (X_i P_{T-t}f)(Y_t)^2 dt \right)^{\frac{p}{2}} \right) &\leq \mathbb{E} \left(\left(\int_0^T \left(\frac{1}{2} \sum_{i=1}^d X_i^2 + X_0 + \frac{\partial}{\partial t} \right) (P_{T-t}f)^2(Y_t) dt \right)^{\frac{p}{2}} \right) \\
&\leq p^{p/2} \left(\frac{p}{p-2} \right)^{p/2} \|f\|_p^p
\end{aligned}$$

Combining this with Theorem 2.5, proves the result in the range $2 < p < \infty$. Finally, the adjoint of \mathcal{S}_A^T acting on $L_\mu^p(\mathbb{M})$ is $\mathcal{S}_{A^*}^T$ acting on $L_\mu^q(\mathbb{M})$, where $\frac{1}{p} + \frac{1}{q} = 1$. The result is then obtained by duality. \square

Proof of Theorem 1.1. It remains to show that we can let $T \rightarrow \infty$ in Lemma 3.2. To see this observe that by applying the lemma with the matrix

$$A_{ij}(t, x) = \frac{(X_i P_t f)(x)(X_j P_t g)(x)}{\left(\sum_{i=1}^d (X_i P_t f)^2(x)\right)^{1/2} \left(\sum_{i=1}^d (X_i P_t g)^2(x)\right)^{1/2}}$$

we obtain the uniform bound

$$(3.4) \quad \int_0^T \int_{\mathbb{M}} \left(\sum_{i=1}^d (X_i P_t f)^2(x)\right)^{1/2} \left(\sum_{i=1}^d (X_i P_t g)^2(x)\right)^{1/2} d\mu dt \leq a_p \|f\|_p \|g\|_q$$

with $\frac{1}{p} + \frac{1}{q} = 1$, where $a_p = C_p$ in the case of $V \neq 0$ and otherwise it is $p^* - 1$. Since

$$\int_{\mathbb{M}} (\mathcal{S}_A^T f) g d\mu = \sum_{i,j=1}^d \int_0^T \int_{\mathbb{M}} A_{ij}(t) (X_i P_t g)(X_j P_t f) d\mu dt,$$

we deduce that

$$\begin{aligned} \lim_{T \rightarrow \infty} \int_{\mathbb{M}} (\mathcal{S}_A^T f) g d\mu &= \sum_{i,j=1}^d \int_0^{+\infty} \int_{\mathbb{M}} A_{ij}(t) (X_i P_t g)(X_j P_t f) d\mu dt \\ &= \int_{\mathbb{M}} (\mathcal{S}_A f) g d\mu \end{aligned}$$

and moreover that

$$\left| \int_{\mathbb{M}} (\mathcal{S}_A f) g d\mu \right| \leq a_p \|A\| \|f\|_p \|g\|_q.$$

This shows that the same bounds in Lemma 3.2 hold with \mathcal{S}_A^T replaced by \mathcal{S}_A . This completes the proof of Theorem 1.1. \square

The following corollary of the above proof generalizes to manifolds the key estimate of Nazarov and Volberg [24] and Dragičević and Volberg [11], [12], for the Laplacian in \mathbb{R}^d ; see [2, Corollary 3.9.1].

Corollary 3.1. *For all $f, g \in C_0^\infty(\mathbb{M})$, $1 < p < \infty$,*

$$(3.5) \quad \int_0^\infty \int_{\mathbb{M}} \left(\sum_{i=1}^d (X_i P_t f)^2(x)\right)^{1/2} \left(\sum_{i=1}^d (X_i P_t g)^2(x)\right)^{1/2} d\mu dt \leq a_p \|f\|_p \|g\|_q,$$

with $\frac{1}{p} + \frac{1}{q} = 1$, where $a_p = C_p$ as in (3.2), if $V \neq 0$ and $a_p = p^* - 1$, if $V \equiv 0$.

We now state a result that follows from Theorem 2.4 when we have some additional information on the matrix $A = (A_{ij})$. Again, this is exactly as on \mathbb{R}^d ; see [3].

Theorem 3.1. *Suppose $V \equiv 0$. Let $A = (A_{ij})$ be symmetric and suppose that there are universal constants $-\infty < b < B < \infty$ such that for all $(t, x) \in [0, \infty) \times M$,*

$$b|\xi|^2 \leq \sum_{i,j=1}^d A_{ij}(t, x)\xi_i\xi_j \leq B|\xi|^2,$$

for all $\xi \in \mathbb{R}^d$. Then for all $1 < p < \infty$ we have

$$(3.6) \quad \|\mathcal{S}_A f\|_p \leq C_{p,b,B} \|f\|_p,$$

where $C_{p,b,B}$ is the constant in Theorem 2.4.

Proof. As above, this result will follow if we can prove the same result for the operator

$$\mathcal{S}_A^T f(x) = \mathbb{E} \left(\sum_{i,j=1}^d \int_0^T A_{ij}(T-t, Y_t) (X_j P_{T-t} f)(Y_t) dB_t^i \mid Y_T = x \right).$$

Consider the two martingales

$$Y_t = \sum_{i,j=1}^d \int_0^T A_{ij}(T-t, Y_t) (X_j P_{T-t} f)(Y_t) dB_t^i, \quad X_t = \sum_{i,j=1}^d \int_0^T (X_j P_{T-t} f)(Y_t) dB_t^i.$$

It is simple to verify (see [3]) that under our assumptions on the matrix A , these martingales satisfy the hypothesis of Theorem 2.4. From this and the contraction of the conditional expectation on L^p we obtain (3.6) for the operators \mathcal{S}_A^T . \square

3.1 An estimate of the constant in (1.3)

The above approach based on Theorem 2.5 to prove (1.3) is general but does not lead to explicit constants with useful information. We present below an alternative argument which provides explicit constants with some additional information. We use the notations introduced in the previous section.

Proposition 3.1. *Suppose that $V \neq 0$ and that $1 < p < \infty$. Then*

$$\left\| \sum_{i=1}^d \int_0^T (X_i P_{T-t} f)(Y_t) dB_t^i \right\|_p \leq \frac{2^{2-\frac{1}{p}} p^2}{p-1} \|f\|_p$$

Proof. From Itô's formula applied to $(P_{T-t} f)(Y_t)$, we find

$$(P_{T-t} f)(Y_t) = P_T f(Y_0) + \int_0^t \left(\frac{\partial}{\partial s} + \frac{1}{2} \sum_{i=1}^d X_i^2 + X_0 \right) (P_{T-s} f)(Y_s) dt + \sum_{i=1}^d \int_0^t (X_i P_{T-t} f)(Y_t) dB_t^i.$$

But,

$$\left(\frac{\partial}{\partial s} + \frac{1}{2} \sum_{i=1}^d X_i^2 + X_0 \right) (P_{T-s}f)(Y_s) = -V(Y_s)(P_{T-s}f)(Y_s)$$

and therefore

$$(P_{T-t}f)(Y_t) = P_T f(Y_0) - \int_0^t V(Y_s)(P_{T-s}f)(Y_s)dt + \sum_{i=1}^d \int_0^t X_i(P_{T-s}f)(Y_s)dB_s^i.$$

Assume now $f \geq 0$. In that case, the above computation shows that $(P_{T-t}f)(X_t)$ is a non negative submartingale. Thus from Lenglart-Lépingle-Pratelli (see Theorem 3.2 in [19]) and Doob's maximal inequality, we have

$$\left\| P_T f(Y_0) - \int_0^T V(Y_s)(P_{T-s}f)(Y_s)dt \right\|_p \leq p \left\| \sup_{0 \leq t \leq T} (P_{T-t}f)(Y_t) \right\|_p \leq \frac{p^2}{p-1} \|f\|_p.$$

We conclude

$$\left\| \sum_{i=1}^d \int_0^T (X_i P_{T-t}f)(Y_t)dB_t^i \right\|_p \leq \frac{2p^2}{p-1} \|f\|_p.$$

For a general f , we can write

$$f = f^+ - f^-$$

and we see that

$$\begin{aligned} \left\| \sum_{i=1}^d \int_0^t X_i(P_{T-s}f)(Y_s)dB_s^i \right\|_p^p &= \left\| \sum_{i=1}^d \int_0^t X_i(P_{T-s}f^+)(Y_s)dB_s^i - \sum_{i=1}^d \int_0^t X_i(P_{T-s}f^-)(Y_s)dB_s^i \right\|_p^p \\ &\leq 2^{p-1} \left\| \sum_{i=1}^d \int_0^t X_i(P_{T-s}f^+)(Y_s)dB_s^i \right\|_p^p \\ &\quad + 2^{p-1} \left\| \sum_{i=1}^d \int_0^t X_i(P_{T-s}f^-)(Y_s)dB_s^i \right\|_p^p \\ &\leq 2^{p-1} \left(\frac{2p^2}{p-1} \right)^p (\|f\|_p^p + \|f^-\|_p^p) \\ &\leq 2^{p-1} \left(\frac{2p^2}{p-1} \right)^p \|f\|_p^p. \end{aligned}$$

□

From this we obtained the following explicit bound in (1.3).

Corollary 3.2. *Let $1 < p < \infty$. For any $V \geq 0$, $f \in L_\mu^p(\mathbb{M})$,*

$$\|\mathcal{S}_A f\|_p \leq 8\|A\|(p^* - 1) \frac{p^4}{(p-1)^2} \|f\|_p$$

Proof. We first note that the martingale

$$\sum_{i,j=1}^d \int_0^T A_{ij}(T-t)(X_j P_{T-t} f)(Y_t) dB_t^i$$

is differentially subordinate to

$$\|A\| \sum_{i=1}^d \int_0^T (X_i P_{T-t} f)(Y_t) dB_t^i.$$

From Theorem 2.3 and the above proposition we obtain

$$\begin{aligned} \int_{\mathbb{M}} (\mathcal{S}_A^T f) g d\mu &= \sum_{i,j=1}^d \int_0^T \int_{\mathbb{M}} A_{ij}(t)(X_i P_t g)(X_j P_t f) d\mu dt \\ &= \mathbb{E} \left(\sum_{i,j=1}^d \int_0^T A_{ij}(T-t)(X_j P_{T-t} f)(Y_t) dB_t^i \sum_{i=1}^d \int_0^T (X_i P_{T-t} g)(Y_t) dB_t^i \right) \\ &\leq \left\| \sum_{i,j=1}^d \int_0^T A_{ij}(T-t)(X_j P_{T-t} f)(Y_t) dB_t^i \right\|_p \left\| \sum_{i=1}^d \int_0^T (X_i P_{T-t} g)(Y_t) dB_t^i \right\|_q \\ &\leq \|A\| (p^* - 1) \left\| \sum_{i=1}^d \int_0^T (X_i P_{T-t} f)(Y_t) dB_t^i \right\|_p \left\| \sum_{i=1}^d \int_0^T (X_i P_{T-t} g)(Y_t) dB_t^i \right\|_q \\ &\leq \|A\| (p^* - 1) \frac{2^{2-\frac{1}{p}} p^2 2^{2-\frac{1}{q}} q^2}{p-1} \|f\|_p \|g\|_q \\ &\leq 8 \|A\| (p^* - 1) \frac{p^4}{(p-1)^2} \|f\|_p \|g\|_q, \end{aligned}$$

which implies the corollary. \square

4 Applications

4.1 Multipliers of Laplace transform-type for Schrödinger operators on manifolds

We work in the same setting as the previous Section. We consider the following *multiplier* for the Schrödinger operator $L = -\frac{1}{2} \sum_{i=1}^d X_i^* X_i + V$. First, let $a \in L^\infty[0, \infty)$. Set

$$T_a f = \int_0^{+\infty} a(t) L P_{2t} f dt.$$

We can observe that since L is essentially self-adjoint, from spectral theorem, there is a measure space (Ω, ν) , a unitary map $U : L_\nu^2(\Omega) \rightarrow L_\mu^2(\mathbb{M})$ and a non negative measurable function on Ω such that

$$U^{-1} L U f(x) = -\lambda(x) f(x), \quad x \in \Omega.$$

The operator T_a acts on these as multiplication operator on $L^2_\nu(\Omega)$ in the sense that

$$U^{-1}T_a U f(x) = -\lambda(x) \int_0^{+\infty} a(t)e^{-t\lambda(x)} dt f(x), \quad x \in \Omega.$$

Let us also observe that in several settings the operators T_a can be interpreted as multipliers in the Fourier analysis sense. For instance, let G be a compact Lie group. Let U be a bounded operator on $L^2(G)$ which commutes with left and right translations. Then there exists a bounded function $\Phi(m)$ on \hat{G} , the space of equivalence classes of irreducible unitary representations of G , such that

$$Uf = \sum_{m \in \hat{G}} \Phi(m) d_m \chi_m * f,$$

where χ_m is the character and d_m the dimension of the representation. Conversely, for any bounded Φ , the above operator defines a bounded operator on $L^2(G)$ which commutes with left and right translations. In this framework, if L is an essentially self-adjoint diffusion operator that commutes with left and right translations (like the Laplace-Beltrami operator for a bi-invariant metric), then we have

$$T_a f = \sum_{m \in \hat{G}} -\lambda(m) \int_0^{+\infty} a(t)e^{-t\lambda(m)} dt d_m \chi_m * f$$

where $(-\lambda(m))_{m \in \hat{G}}$ is the spectrum of L .

With the notations of the previous section, we see that

$$T_a f = -\frac{1}{2} S_A f + \int_0^{+\infty} a(t) P_t V P_t f dt$$

where $A(t) = a(t)\mathbf{Id}$. In the following we denote by Q_t the Markovian semigroup with generator $-\frac{1}{2} \sum_{i=1}^d X_i^* X_i$.

Proposition 4.1. *Fix $1 < p < \infty$.*

(i) *For any nonnegative potential V that satisfies $V \leq -m$ for some $m \geq 0$, we have*

$$(4.1) \quad \|T_a f\|_p \leq \left(4\|a\|_\infty (p^* - 1) \frac{p^4}{(p-1)^2} + \int_0^{+\infty} |a(t)| e^{-2mt} \|Q_t |V|^q\|_\infty^{1/q} dt \right) \|f\|_p,$$

with $\frac{1}{q} + \frac{1}{p} = 1$.

(ii) *Suppose $V \equiv 0$. Let a be such that for all $t \in [0, \infty)$, $-\infty < b \leq a(t) \leq B < \infty$. Then*

$$(4.2) \quad \|S_a f\|_p \leq \frac{1}{2} C_{p,b,B} \|f\|_p,$$

where the $C_{p,b,B}$ is the constant of Theorem 2.4.

Proof. Since

$$T_a f = -\frac{1}{2} S_A f + \int_0^{+\infty} a(t) P_t V P_t f dt,$$

using the results of the previous section, we only need to bound in L^p the operator $\int_0^{+\infty} a(t) P_t V P_t f dt$. From Feynman-Kac formula, we have

$$P_t V P_t f(x) = \mathbb{E} \left(e^{\int_0^t V(X_s^x) ds} V(X_t^x) P_t f(X_t^x) \right),$$

where $(X_t^x)_{t \geq 0}$ is the diffusion with generator $-\frac{1}{2} \sum_{i=1}^d X_i^* X_i$ started at $x \in \mathbb{M}$. Thus

$$\begin{aligned} |P_t V P_t f(x)| &\leq \mathbb{E} \left(e^{\int_0^t V(X_s^x) ds} |V(X_t^x)|^q \right)^{1/q} \mathbb{E} (|P_t f(X_t^x)|^p)^{1/p} \\ &\leq e^{-2mt} \|Q_t |V|^q\|_\infty^{1/q} \mathbb{E} (Q_t |f|^p | (X_t^x)^p)^{1/p} \\ &\leq e^{-2mt} \|Q_t |V|^q\|_\infty^{1/q} (Q_{2t} |f|^p)(x)^{1/p} \end{aligned}$$

As a consequence,

$$\|P_t V P_t f\|_p \leq e^{-2mt} \|Q_t |V|^q\|_\infty^{1/q} \|f\|_p,$$

which yields the expected result. \square

For instance, we immediately deduce from the previous proposition:

Corollary 4.1. *Fix $1 < p < \infty$.*

- (i) *For any non-negative potential V that satisfying $-M \leq V \leq -m$, for some $m, M \geq 0$, we have*

$$\|T_a f\|_p \leq \left(4\|a\|_\infty (p^* - 1) \frac{p^4}{(p-1)^2} + M \int_0^{+\infty} |a(t)| e^{-2mt} dt \right) \|f\|_p.$$

- (ii) *Assume that the semigroup Q_t is ultracontractive. For any nonnegative potential V that satisfies $V \in L_\mu^q(\mathbb{M})$, we have*

$$\|T_a f\|_p \leq \left(4\|a\|_\infty (p^* - 1) \frac{p^4}{(p-1)^2} + \|V\|_q \int_0^{+\infty} |a(t)| \|Q_t\|_{\infty,1} dt \right) \|f\|_p,$$

with $\frac{1}{q} + \frac{1}{p} = 1$.

Remark 4.1. *Laplace transform-type operators have been extensively studied in many settings. For some of this literature, see [26], [15], [16], [2], and references contained in those papers.*

4.2 Second order Riesz transforms on Lie groups of compact type

Let G be a Lie group of compact type with Lie algebra \mathfrak{g} . We endow G with a bi-invariant Riemannian structure and consider an orthonormal basis X_1, \dots, X_d of \mathfrak{g} . In this setting the Laplace-Beltrami operator can be written as

$$L = \frac{1}{2} \sum_{i=1}^d X_i^2.$$

It is essentially self-adjoint on the space of smooth and compactly supported functions and the assumptions of the previous section are satisfied. It is remarkable here that $X_i^* = -X_i$ and the vector fields X_i do commute with the semigroup $P_t = e^{tL}$. In this case, if the matrix A only depends on time, we get therefore

$$\mathcal{S}_A f = - \sum_{i,j} \int_0^{+\infty} A_{ij}(t) (X_i X_j P_{2t} f) dt.$$

In particular, for a constant A , we obtain

$$\mathcal{S}_A f = \sum_{i,j} A_{ij} \left(\sum_{i=1}^d X_i^2 \right)^{-1} X_i X_j f.$$

Defining the Riesz transforms on G by

$$R_j f = \left(- \sum_{i=1}^d X_i^2 \right)^{-1/2} X_j f$$

we see that

$$\mathcal{S}_A f = \sum_{i,j=1}^d A_{ij} R_i R_j f.$$

We have the following result which follows from Theorems 1.1 and 3.1. These inequality are exactly as those proved in \mathbb{R}^d for the classical Riesz transforms.

Theorem 4.1. Fix $1 < p < \infty$.

(i) For any constant coefficient matrix A ,

$$\left\| \sum_{i,j=1}^d A_{ij} R_i R_j f \right\|_p \leq (p^* - 1) \|A\| \|f\|_p.$$

(ii) Assume that $A = (A_{ij})_{i,j=1}^d$ is symmetric matrix with real entries and eigenvalues $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_d$. Then

$$(4.3) \quad \left\| \sum_{i,j=1}^d A_{ij} R_i R_j f \right\|_p \leq C_{p,\lambda_1,\lambda_d} \|f\|_p$$

and this inequality is sharp. In particular, if $J \subsetneq \{1, 2, \dots, d\}$, then

$$(4.4) \quad \left\| \sum_{j \in J} R_j^2 f \right\|_p \leq C_{p,0,1} \|f\|_p = c_p \|f\|_p,$$

where c_p is the Choi constant in (2.2) and this inequality is also sharp.

The fact that the constants in this theorem are best possible follows from the fact that they are already best possible on \mathbb{R}^d . These results simply show that the construction in [3] for Riesz transforms on \mathbb{R}^d extend to Lie groups of compact type without change in their norms. We may observe that, slightly more generally, a similar statement holds on Riemannian manifolds for which the gradient ∇ commutes with the heat semigroup P_t . From the Bakry-Émery criterion (see [1]), such a commutation implies that the Ricci curvature of \mathbb{M} is non negative.

If G is a semisimple compact Lie group (see for instance the classical reference [14] for an account about structure theory and harmonic analysis on semisimple compact Lie groups), we can deduce from the above result an interesting class of multipliers.

Proposition 4.2. *Let G be a compact semisimple Lie group. Let Λ^+ be the set of highest weights and Δ be the set of all roots. For $\alpha, \beta \in \Delta$, denote*

$$U_{\alpha,\beta} f = \sum_{\lambda \in \Lambda^+} d_\lambda \frac{\langle \lambda, \alpha \rangle \langle \lambda, \beta \rangle}{\|\lambda + \rho\|^2 - \|\rho\|^2} \chi_\lambda * f,$$

where $\rho = \frac{1}{2} \sum_{\alpha \in \Delta} \alpha$, χ_λ is the character of the highest weight representation and d_λ its dimension. Then, for $1 < p < \infty$, $U_{\alpha,\beta}$ is bounded in $L^p(G)$ and

$$\|U_{\alpha,\beta} f\|_p \leq (p^* - 1) \|\alpha\| \|\beta\| \|f\|_p.$$

Proof. Let T be a maximal torus of G and let \mathfrak{t}_0 be its Lie algebra. Then, the sub-algebra \mathfrak{t} of \mathfrak{g} generated by \mathfrak{t}_0 is a Cartan sub-algebra. Let \mathfrak{t}^* denote the dual of \mathfrak{t} . If $\alpha \in \Delta$, we denote by H_α the element of \mathfrak{t} such that for every $H \in \mathfrak{t}$, $\langle H, H_\alpha \rangle = \alpha(H)$ where $\langle \cdot, \cdot \rangle$ is induced from the Killing form. With these notations, we see that

$$U_{\alpha,\beta} = C^{-1} H_\alpha H_\beta,$$

where C is the Casimir operator which is also the Laplace-Beltrami operator. □

References

- [1] D. Bakry & M. Emery: *Diffusions hypercontractives*, Sémin. de probabilités XIX, Univ. Strasbourg, Springer, 1983.
- [2] R. Bañuelos, *The foundational inequalities of D. L. Burkholder and some of their ramifications*, To appear, Illinois Journal of Mathematics, Volume in honor of D.L. Burkholder.

- [3] R. Bañuelos and A. Osękowski, *Martingales and Sharp Bounds for Fourier multipliers*, to appear in *Annales Academiae Scientiarum Fennicae Mathematica*.
- [4] R. Bañuelos and G. Wang, *Sharp inequalities for martingales with applications to the Beurling-Ahlfors and Riesz transforms*, *Duke Math. J.* **80** (1995), no. 3, 575–600.
- [5] F. Baudoin, *An introduction to the geometry of stochastic flows*. Imperial College Press, London, 2004. x+140 pp.
- [6] D.L. Burkholder, *Martingale transforms*, *Ann. Math. Statist.* **37** (1966), 1494-1504.
- [7] D.L. Burkholder, *A sharp inequality for martingale transforms*, *Ann. Prob* **7** (1979), 858-863.
- [8] D.L. Burkholder and R. F. Gundy, *Extrapolation and interpolation of quasi-linear operators on martingales*, *Acta Math.* **124** (1970), 249-304 .
- [9] K. P. Choi, *A sharp inequality for martingale transforms and the unconditional basis constant of a monotone basis in $L^p(0, 1)$* , *Trans. Amer. Math. Soc.* **330** (1992), 509–521.
- [10] C. Dellacherie and P. A. Meyer, *Probabilities and Potential B: Theory of martingales*, North Holland, Amsterdam, 1982.
- [11] O. Dragičević and A. Volberg, *Sharp estimate of the Ahlfors-Beurling operator via averaging martingale transforms*, *Michigan Math. J.* **51** (2003), 415–435.
- [12] O. Dragičević and A. Volberg, *Bellman function, Littlewood-Paley estimates and asymptotics for the Ahlfors-Beurling operator in $L^p(\mathbb{C})$* , *Indiana Univ. Math. J.* **54** (2005), no. 4, 971–995.
- [13] E. Geiss, S. Montgomery-Smith, E. Saksman, *On singular integral and martingale transforms*, *Trans. Amer. Math. Soc.* **362** (2010), 553–575.
- [14] S. Helgason, *Groups and Geometric Analysis*, Academic Press, Orlando (1984)
- [15] T.P. Hytönen, *Littlewood-Paley-Stein theory for semigroups in UMD spaces*, *Rev. Mat. Iberoam.* **23**(2007), 973–1009.
- [16] T.P. Hytönen, *Aspects of probabilistic Littlewood-Paley theory in Banach spaces*, *Banach spaces and their applications in analysis*, Walter de Gruyter, Berlin, (2007), 343–355,
- [17] D. Jerison & A. Sánchez-Calle, *Subelliptic second order differential operators*, *Lecture. Notes in Math.*, 1277 (1987), pp. 46-77.
- [18] E. Lenglart, *Relation de domination entre deux processus*, *Ann. Inst. H. Poincaré Prob. Statist.* **13** (1977), 171–179.
- [19] E. Lenglart, D. Lépingle and M. Pratelli, *Présentation unifiée de certaines inégalités de la théorie des martingales*, *Séminaire de probabilités de Strasbourg* **14** (1980), 26–48

- [20] X.-D. Li, *Martingale transform representation and L^p -estimates for Beurling-Ahlfors and Riesz transforms on Riemannian manifolds*, (**preprint**).
- [21] X.-D. Li, *Riesz transforms on forms and L^p -Hodge decomposition on complete Riemannian manifolds*, *Rev. Mat. Iberoam.* **26** (2010), 481–528
- [22] X.-D. Li, *On the weak L^p -Hodge decomposition and Beurling-Ahlfors transforms on complete Riemannian manifolds*, *Probab. Theory Relat. Fields* **150** (2011), 111–144.
- [23] X.-D. Li, *Martingale transforms and L^p -norm estimates of Riesz transforms on complete Riemannian manifolds*, *Probab. Theory Relat. Fields* **141** (2008), 247–281.
- [24] F. Nazarov and A. Volberg, *Heat extension of the Beurling operator and estimates for its norm*, *St. Petersburg Math. J.* **15**, (2004), 563-573.
- [25] D. Revuz and M. Yor, *Continuous martingales and Brownian motion*, Springer, 2005.
- [26] N. Th. Varopoulos, *Aspects of probabilistic Littlewood-Paley theory*, *J. Funct. Anal.* **38** (1980), no. 1, 25–60.
- [27] G. Wang, *Differential subordination and strong differential subordination for continuous-time martingales and related sharp inequalities*, *Ann. Probab.* **23** no. 2 (1995), 522–551.