

# Meromorphic solutions of Briot–Bouquet type differential equations

A. Eremenko

March 2011

Let  $F$  be a polynomial in two variables. Assume that the differential equation

$$F(y^{(k)}, y) = 0$$

has a solution  $y(z)$  which is meromorphic in the whole plane. What can be said about  $y$ ?

For  $k = 1$ , a complete answer is well known. Namely, all meromorphic solutions belong to the class  $W$ , which consists by definition of: all elliptic functions, all functions of the form  $R(\exp(az))$ ,  $R$  rational and  $a \neq 0$  is a complex number, and all rational functions. And vice versa, each function of the class  $W$  satisfies some differential equation of the above form with  $k = 1$ .

For  $k = 2$  Emile Picard proved in 1880 that all meromorphic solutions also belong to the class  $W$ . This result was forgotten, in 1978 E. Hille stated it as a conjecture, and in 1981 S. Bank and R. Kaufman gave a proof, more complicated than the one Picard published 101 years before.

It follows from another theorem of Picard that non-constant meromorphic solutions are possible only if the genus of the curve  $F(x, y) = 0$  is at most 1.

In the case when the genus is equal to 1 I proved for every  $k$ , that all meromorphic solutions must be elliptic functions.

Thus only the case of genus 0 remains. I also proved the following. If  $k$  is odd and a solution  $y$  is meromorphic in the plane and has at least one pole, it must belong to  $W$ .

The simplest equation not covered by these results is

$$y^{(IV)} = 24y^5.$$

It evidently has solutions of the form  $1/(z - b)$ . Does it have any other meromorphic solutions?

*Added on May 22, 2007.* L. W. Liao, T. W. Ng and the present author recently proved that for every  $k$ , all meromorphic solutions of  $F(y^{(k)}, y) = 0$  having at least one pole belong to  $W$ .

Eremenko, Alexandre E.; Liao, Liangwen; Ng, Tuen Wai, Meromorphic solutions of higher order Briot–Bouquet differential equations. *Math. Proc. Cambridge Philos. Soc.* 146 (2009), no. 1, 197–206.

This proof also implies that the only meromorphic functions that satisfy  $y^{(k)} = y^m$  with  $m > 1$  are of the form  $c(z - b)^{-n}$ .

The question of description of entire solutions remains open. Notice that non-trivial entire solutions of  $y''' = y$  do not belong to  $W$ . So it is not completely clear what is the right conjecture about entire solutions. Perhaps they can be only exponential polynomials?

*Added on June 19, 2023.* Further progress was achieved in the paper:

A. Ya. Yanchenko, One advance in the proof of the conjecture on meromorphic solutions of Briot-Bouquet type equations. (Russian) *Izv. Ross. Akad. Nauk Ser. Mat.* 86 (2022), no. 5, 197–208; translation in *Izv. Math.* 86 (2022), no. 5, 1020–1030.

This author proves that if  $F$  is an irreducible polynomial of degree  $d \geq 2$ , and the top degree homogeneous part of  $F$  has distinct roots, then all entire solutions are Laurent polynomials of  $e^{az}$  with some complex  $a$ .

## References

- [1] S. Bank and R. Kaufman, On Briot-Bouquet differential equations and a question of Einar Hille. *Math. Z.* 177 (1981), no. 4, 549–559.
- [2] A. Eremenko, Meromorphic solutions of equations of Briot–Bouquet type, *Teor. Funktsii, Funk. Anal. i Prilozh.*, 38 (1982) 48–56. English translation: *Amer. Math. Soc. Transl. (2)* Vol. 133 (1986) 15–23.
- [3] E. Picard, Sur une propriété des fonctions uniformes d'une variable et sur une classe d'équations différentielles, *C. R. Acad. Sci. Paris*, 91 (1880) 1058–1061.