

# CONNECTIVE $C^*$ -ALGEBRAS

MARIUS DADARLAT AND ULRICH PENNIG

ABSTRACT. Connectivity is a homotopy invariant property of separable  $C^*$ -algebras which has three notable consequences: absence of nontrivial projections, quasidiagonality and a more geometric realization of KK-theory for nuclear  $C^*$ -algebras using asymptotic morphisms. The purpose of this paper is to further explore the class of connective  $C^*$ -algebras. We give new characterizations of connectivity for exact and for nuclear separable  $C^*$ -algebras and show that an extension of connective separable nuclear  $C^*$ -algebras is connective. We establish connectivity or lack of connectivity for  $C^*$ -algebras associated to certain classes of groups: virtually abelian groups, linear connected nilpotent Lie groups and linear connected semisimple Lie groups.

## 1. INTRODUCTION

Connectivity of separable  $C^*$ -algebras was introduced in our earlier paper [10] under different terminology, see Definition 2.1 below. The initial motivation for studying it stemmed from our search for homotopy-symmetric  $C^*$ -algebras. By a result of Loring and the first author [9], these are precisely the separable  $C^*$ -algebras for which one can unsuspend in the E-theory of Connes and Higson [6]. Using a result of Thomsen [33], we proved in [10] that connectivity is equivalent to homotopy-symmetry for all separable nuclear  $C^*$ -algebras. Moreover, we showed that connectivity has a number of important permanence properties. These facts allowed us to exhibit new classes of homotopy-symmetric  $C^*$ -algebras.

The purpose of this paper is to further explore the class of connective  $C^*$ -algebras. We are motivated by the following three properties they share:

(i) If  $A$  is a separable nuclear  $C^*$ -algebra, then  $KK(A, B)$  is isomorphic to the homotopy classes of completely positive and contractive (cpc) asymptotic morphisms from  $A$  to  $B \otimes \mathcal{K}$  for any separable  $C^*$ -algebra  $B$ .

(ii) Connective  $C^*$ -algebras are quasidiagonal. In fact, if  $A$  is connective, then  $A \otimes B$  is quasidiagonal for any  $C^*$ -algebra  $B$ . ( $A \otimes B$  denotes the minimal tensor product.)

---

M.D. was partially supported by NSF grant #DMS-1362824.

(iii) Connective  $C^*$ -algebras do not have nonzero projections. In fact, if  $A$  is connective, then  $A \otimes B$  does not have nonzero projections for any  $C^*$ -algebra  $B$ .

Connectivity is of particular interest in the case of group  $C^*$ -algebras. A countable discrete group  $G$  is called connective if the augmentation ideal  $I(G)$  defined as the kernel of the trivial representation  $\iota: C^*(G) \rightarrow \mathbb{C}$  is a connective  $C^*$ -algebra. In view of properties (ii) and (iii) connectivity of  $G$  may be viewed as a stringent topological property that accounts simultaneously for the quasidiagonality of  $C^*(G)$  and the verification of the Kadison-Kaplansky conjecture for certain classes of groups. Examples of nonabelian connective groups were exhibited in [10] and [11].

In this paper we give new characterizations of connectivity for exact and nuclear separable  $C^*$ -algebras, see Prop. 2.2, 2.3. We prove that connectivity of separable nuclear  $C^*$ -algebras is preserved under extensions, see Thm. 2.4. This is a key permanence property which does not hold for quasidiagonal  $C^*$ -algebras.

There is a close connection between the topology of the spectrum and connectivity, which we employ to reveal an obstruction to it by using work of Blackadar and Cuntz [1] and Pasnicu and Rørdam [29]. In particular, we show that a countable discrete group  $G$  is not connective if the trivial representation  $\iota$  is a shielded point of the unitary dual of  $G$  in the sense of Def. 2.9.

Motivated by this, we give a complete description of the neighborhood of  $\iota$  in the spectrum of the Hantzsche-Wendt group  $G$ , which is a torsion free crystallographic group with holonomy  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ , in Sec. 3.1. This allows us to prove that  $G$  is not connective in this case (Cor. 3.2). Moreover, we show that this group provides a counterexample to a conjecture from [8]. Specifically, we prove that the natural map  $[[I(G), \mathcal{K}]] \rightarrow K^0(I(G))$  is not an isomorphism (Lem. 3.3). In contrast, we show that all torsion free crystallographic groups with *cyclic holonomy* are connective (Thm. 3.8).

Next, we investigate connectivity for  $C^*$ -algebras associated to Lie groups. We show that all noncompact linear connected nilpotent Lie groups have connective  $C^*$ -algebras (Thm. 4.3). Using classic results from representation theory in conjunction with permanence properties of connectivity, we show that if  $G$  is a linear connected complex semisimple Lie group, then  $C_r^*(G)$  is connective if and only if  $G$  is not compact (Thm. 4.5). Moreover, if  $G$  is a linear connected real reductive Lie group, then  $C_r^*(G)$  is connective if and only if  $G$  does not have a compact Cartan subgroup (Thm. 4.6).

A common denominator of our results concerning group  $C^*$ -algebras is that in all the cases we analyzed  $C^*(G)$  contains a large connective ideal.

2. CONNECTIVE  $C^*$ -ALGEBRAS

**2.1. Definitions and background.** For a  $C^*$ -algebra  $A$ , the *cone over  $A$*  is defined as  $CA = C_0[0, 1] \otimes A$  the *suspension of  $A$*  as  $SA = C_0(0, 1) \otimes A$ .

The first of the following two notions was introduced in [10, Def. 2.6 (i)], under a different terminology which we have abandoned. We use the abbreviation *cpc map* for a completely positive and contractive map.

**Definition 2.1.** Let  $A$  be a  $C^*$ -algebra.

(a)  $A$  is *connective* if there is a  $*$ -monomorphism

$$\Phi: A \rightarrow \prod_n CL(\mathcal{H}) / \bigoplus_n CL(\mathcal{H})$$

which is liftable to a cpc map  $\varphi: A \rightarrow \prod_n CL(\mathcal{H})$ .

(b)  $A$  is *almost connective*, if there is a (not necessarily liftable)  $*$ -monomorphism  $\Phi: A \rightarrow \prod_n CL(\mathcal{H}) / \bigoplus_n CL(\mathcal{H})$ .

For a discrete group  $G$ , we define  $I(G)$  to be the augmentation ideal, i.e. the kernel of the trivial representation  $C^*(G) \rightarrow \mathbb{C}$ . We will sometimes say that a discrete amenable group  $G$  is connective if the  $C^*$ -algebra  $I(G)$  is connective. Note that (almost) connective  $C^*$ -algebras do not have nonzero projections.

Let  $A$  and  $B$  be  $C^*$ -algebras. An *asymptotic morphism* is a family of maps  $\{\varphi_t: A \rightarrow B\}_{t \in [0, \infty)}$  such that

- a) for each  $a \in A$  the map  $t \mapsto \varphi_t(a)$  is norm-continuous and bounded,
- b) for all  $a, b \in A$  and  $\lambda \in \mathbb{C}$ , we have

$$\begin{aligned} \lim_{t \rightarrow \infty} \|\varphi_t(a + \lambda b) - (\varphi_t(a) + \lambda \varphi_t(b))\| &= 0 \\ \lim_{t \rightarrow \infty} \|\varphi_t(ab) - \varphi_t(a) \varphi_t(b)\| &= 0 \\ \lim_{t \rightarrow \infty} \|\varphi_t(a^*) - \varphi_t(a)^*\| &= 0. \end{aligned}$$

Observe that we do not assume  $\varphi_t$  to be (asymptotically) unital. A *discrete asymptotic morphism*  $(\varphi_n)_{n \in \mathbb{N}}$  between  $A$  and  $B$  is a family of maps  $\varphi_n: A \rightarrow B$  that satisfies the analogous conditions as a) and b) above with the index set replaced by  $\mathbb{N}$ . A homotopy between two (discrete) asymptotic morphisms  $(\varphi_t^0)_{t \in I}$  and  $(\varphi_t^1)_{t \in I}$  is a (discrete) asymptotic morphism  $H_t: A \rightarrow C[0, 1] \otimes B$ , such that  $\text{ev}_i \circ H_t = \varphi_t^i$  for all  $t \in I$ , where  $I$  either denotes  $[0, \infty)$  or  $\mathbb{N}$ . We will say that a (discrete) asymptotic morphism  $(\varphi_t)_{t \in I}$  is completely positive and contractive (cpc) if each of the maps  $\varphi_t$  is cpc. The corresponding homotopy classes will be denoted as follows:

- $[[A, B]]$  – homotopy classes of asymptotic morphisms,
- $[[A, B]]_{\mathbb{N}}$  – homotopy classes of discrete asymptotic morphisms,
- $[[A, B]]_{\mathbb{N}}^{\text{cpc}}$  – homotopy classes of discrete cpc asymp. morphisms

**2.2. Characterizations of connectivity.** In the following we give two more characterizations of connectivity for exact and respectively nuclear  $C^*$ -algebras.

**Proposition 2.2.** *Let  $A$  be a separable exact  $C^*$ -algebra. Then  $A$  is connective if and only if there is an injective  $*$ -homomorphism  $\pi: A \rightarrow \mathcal{O}_2$  which is null-homotopic as a discrete cpc asymptotic morphism. This means that  $[[\pi]] = 0$  in the set  $[[A, \mathcal{O}_2]]_{\mathbb{N}}^{cp}$ .*

*Proof.*  $(\Rightarrow)$  By assumption, there is a cpc discrete asymptotic morphism  $\{\varphi_n: A \rightarrow C[0, 1] \otimes \mathcal{O}_2\}_n$  such that  $\varphi_n^{(0)} = \pi$  is an injective  $*$ -homomorphism and  $\varphi_n^{(1)} = 0$ . Thus, we can view  $\{\varphi_n\}_n$  as an injective discrete asymptotic morphism  $\{\varphi_n: A \rightarrow C_0[0, 1] \otimes \mathcal{O}_2 \subset CL(\mathcal{H})\}_n$  and hence  $A$  is connective.

$(\Leftarrow)$  Suppose that  $A$  is a separable exact connective  $C^*$ -algebra. By [10, Prop. 2.11] it follows that  $[[A, \mathcal{O}_2 \otimes \mathcal{K}]]_{\mathbb{N}}^{cp}$  is an abelian group. By Kirchberg's embedding theorem, there is an injective  $*$ -homomorphism  $\pi: A \rightarrow \mathcal{O}_2 \otimes \mathcal{K}$ . Moreover  $\pi \oplus \pi: A \rightarrow \mathcal{O}_2 \otimes \mathcal{K}$  is unitarily homotopy equivalent to  $\pi$ . It follows that  $[[\pi]] \oplus [[\pi]] = [[\pi]]$  in the group  $[[A, \mathcal{O}_2 \otimes \mathcal{K}]]_{\mathbb{N}}^{cp}$  and hence  $[[\pi]] = 0$ . After embedding  $\mathcal{O}_2 \otimes \mathcal{K}$  into  $\mathcal{O}_2$  we obtain the desired conclusion.  $\square$

**Proposition 2.3.** *Let  $A$  be a separable nuclear  $C^*$ -algebra. The following properties are equivalent.*

- (i)  $A$  is connective.
- (ii)  $A \otimes B$  is connective for some  $C^*$ -algebra  $B$  that contains a nonzero projection.
- (iii)  $A \otimes B$  is connective for all  $C^*$ -algebras  $B$
- (iv)  $[[A, \mathcal{O}_2 \otimes \mathcal{K}]] = 0$ .
- (v)  $[[A, L(\mathcal{H}) \otimes \mathcal{K}]] = 0$ .

*Proof.* The equivalences  $(i) \Leftrightarrow (ii) \Leftrightarrow (iii)$  were established in [10]. (For  $(ii) \Rightarrow (i)$  observe that  $A$  is a subalgebra of  $A \otimes B$  if  $B$  contains a nonzero projection.)

$(i) \Rightarrow (iv)$  and  $(i) \Rightarrow (v)$ . Let  $B$  be a  $\sigma$ -unital  $C^*$ -algebra such that  $KK(A, B) = 0$ , for instance  $B = \mathcal{O}_2$  or  $B = L(\mathcal{H})$ . If  $A$  is connective, then  $A$  is homotopy symmetric and hence  $[[A, B \otimes \mathcal{K}]] \cong KK(A, B) = 0$  by [10, Thm. 3.1]. Note that even though [10, Thm. 3.1] was stated for separable  $C^*$ -algebras  $B$  it is routine to extend the result to general  $C^*$ -algebras using the separability of  $A$ .

$(iv) \Rightarrow (i)$  and  $(v) \Rightarrow (i)$ . Fix an embedding  $\pi: A \rightarrow \mathcal{O}_2 \subset L(\mathcal{H}) \otimes \mathcal{K}$  and regard it a constant asymptotic morphism  $\{\pi_t: A \rightarrow L(\mathcal{H}) \otimes \mathcal{K}\}_t$ . By assumption,  $[[\pi_t]] = 0$  in  $[[A, L(\mathcal{H}) \otimes \mathcal{K}]]$  and hence by restriction,  $[[\pi_n]] = 0$  in  $[[A, L(\mathcal{H}) \otimes \mathcal{K}]]_{\mathbb{N}}$ . We shall view  $L(\mathcal{H}) \otimes \mathcal{K}$  as a subalgebra of  $L(\mathcal{H})$ . The corresponding homotopy from the constant discrete morphism  $\{\pi_n\}_n$  to zero

will induce an embedding  $\Phi: A \rightarrow \prod_n CL(\mathcal{H}) / \bigoplus_n CL(\mathcal{H})$  which is liftable to a cpc map  $\varphi: A \rightarrow \prod_n CL(\mathcal{H})$  by the nuclearity of  $A$ .  $\square$

**2.3. Extensions of connective  $C^*$ -algebras.** Connectivity of  $C^*$ -algebras has a plethora of permanence properties as proven in [10, Thm. 3.3]. In particular, it is inherited by split extensions [10, Thm. 3.3 (d)]. In the following theorem this result is extended to non-split extensions as well.

**Theorem 2.4.** *Let  $0 \rightarrow J \rightarrow A \rightarrow B \rightarrow 0$  be an exact sequence of separable nuclear  $C^*$ -algebras. If  $J$  and  $B$  are connective, then  $A$  is connective.*

*Proof.* Since connectivity passes to nuclear subalgebras we may replace the given extension by

$$0 \rightarrow J \otimes \mathcal{O}_2 \otimes \mathcal{K} \rightarrow A \otimes \mathcal{O}_2 \otimes \mathcal{K} \rightarrow B \otimes \mathcal{O}_2 \otimes \mathcal{K} \rightarrow 0.$$

Adding to this extension a trivial absorbing extension, using the addition in Ext-theory we obtain an absorbing extension

$$(1) \quad 0 \rightarrow J \otimes \mathcal{O}_2 \otimes \mathcal{K} \rightarrow E \rightarrow B \otimes \mathcal{O}_2 \otimes \mathcal{K} \rightarrow 0,$$

with  $A \subset A \otimes \mathcal{O}_2 \otimes \mathcal{K} \subset E$  by construction, see for instance [4, Lemma 2.2]. Since  $\text{Ext}(B \otimes \mathcal{O}_2, J \otimes \mathcal{O}_2) = 0$  as  $\mathcal{O}_2$  is KK-contractible and we are dealing with an absorbing extension, it follows that the extension (1) splits by [23, Sec. 7] and so  $E$  is connective by [10, Thm. 3.3]. We conclude that  $A \subset E$  is connective.  $\square$

In the sequel we will need to use the following result from [10], which is based on [2].

**Theorem 2.5** ([10], Cor. 3.4.). *Let  $A$  be a separable continuous field of nuclear  $C^*$ -algebras over a compact connected metrizable space  $X$ . If one of the fibers of  $A$  is connective, then  $A$  is connective.*

**Corollary 2.6.** *Let  $A$  be a separable continuous field of nuclear  $C^*$ -algebras over a locally compact metrizable space  $X$  that has no compact open subsets. Then  $A$  is connective.*

*Proof.* Let  $Y = X \cup \{y_0\}$  be the one-point compactification of  $X$ . Then  $Y$  is a compact metrizable space which must be connected. Indeed, arguing by contradiction, say that  $Y = U \cup V$  with  $U, V$  open and nonempty with  $y_0 \in U$  and  $U \cap V = \emptyset$ . Then  $V = V \cap X$  is both an open and compact subset of  $X$ .

We can view  $A$  as a continuous field over  $Y$  (see the remark on page 145 of [2]) and the fiber over  $y_0$  satisfies  $A(y_0) = \{0\}$ . It follows that  $A$  is connective by Thm. 2.5.  $\square$

**2.4. Obstructions to connectivity.** If  $A$  is a  $C^*$ -algebra, we denote by  $\widehat{A}$  the spectrum of  $A$ , which consists of all unitary equivalence classes of irreducible representations and by  $\text{Prim}(A)$  the primitive spectrum of  $A$  consisting of kernels of irreducible representations. The unitary dual  $\widehat{G}$  of a group  $G$  identifies with  $\widehat{C^*(G)}$ . Recall that  $\widehat{A}$  is topologized by pulling-back the Jacobson topology of  $\text{Prim}(A)$  under the natural map  $\widehat{A} \rightarrow \text{Prim}(A)$ ,  $\pi \mapsto \ker \pi$ , [14]. Let  $\pi$  and  $(\pi_n)_n$  be irreducible representations of  $A$  acting on the same separable Hilbert space  $H$ . Suppose that  $\|\pi_n(a)\xi - \pi(a)\xi\| \rightarrow 0$  for all  $a \in A$  and  $\xi \in H$ . Then the sequence  $(\pi_n)_n$  converges to  $\pi$  in the topology of  $\widehat{A}$ , see [14, Sec. 3.5].

**Proposition 2.7.** *Let  $A$  be a separable  $C^*$ -algebra.*

- (i) *If  $\text{Prim}(A)$  contains a non-empty compact open subset, then  $A$  is not connective.*
- (ii) *If  $A$  is nuclear and  $\text{Prim}(A)$  is Hausdorff, then  $A$  is connective if and only if  $\text{Prim}(A)$  does not contain a non-empty compact open subset.*

*Proof.* (i) Set  $X = \text{Prim}(A)$ . If  $X$  has a non-empty compact open subset, then  $A \otimes \mathcal{O}_2$  contains a nonzero projection by [29, Prop. 2.7] and hence  $A$  cannot be connective.

(ii) One implication follows from (i). For the other implication suppose that  $X = \text{Prim}(A)$  does not contain a non-empty compact open subset. Since  $X$  is Hausdorff by assumption,  $A$  is a nuclear separable continuous field over the locally compact space  $X$ , [16]. This is explained in detail in [3, Sec. 2.2.2]. Now we apply Cor. 2.6.  $\square$

We would like thank Gabor Szabo for pointing out the following invariance property of connectivity.

**Proposition 2.8.** *Let  $A$  and  $B$  be separable nuclear  $C^*$ -algebra with homeomorphic primitive spectra. Then  $A$  is connective if and only if  $B$  is connective.*

*Proof.* Kirchberg's classification theorem [25] implies that if  $A$  and  $B$  are as in the statement, then  $A \otimes \mathcal{O}_2 \otimes \mathcal{K} \cong B \otimes \mathcal{O}_2 \otimes \mathcal{K}$ . The desired conclusion follows now from Proposition 2.3.  $\square$

**Definition 2.9.** Let  $A$  be a separable  $C^*$ -algebra. A point  $\pi \in \widehat{A}$  is called *shielded*, if any sequence  $(\pi_n)_n$  in  $\widehat{A} \setminus \{\pi\}$  which converges  $\pi$  also converges to another point  $\eta \in \widehat{A} \setminus \{\pi\}$ .

**Lemma 2.10.** *Let  $A$  be a unital separable  $C^*$ -algebra. If  $\pi \in \widehat{A}$  is closed and shielded, then  $I = \ker \pi$  is not connective.*

*Proof.* It suffices to show that  $I \otimes \mathcal{O}_2$  contains a non-trivial projection, since this implies that  $I$  can not be connective. By [29, Prop. 2.7] this will follow if we can show that  $I \otimes \mathcal{O}_2 \subset A \otimes \mathcal{O}_2$  corresponds to a compact open subset of  $\text{Prim}(A \otimes \mathcal{O}_2) \cong \text{Prim}(A)$ . To this goal we will show that  $\text{Prim}(A) \setminus \{I\}$  is a compact-open subset of  $\text{Prim}(A)$ . The quotient map  $q: \widehat{A} \rightarrow \text{Prim}(A)$  is open. Since  $\pi$  is closed it follows that  $q^{-1}(I) = \{\pi\}$ . Therefore the lemma follows if  $\widehat{A} \setminus \{\pi\}$  is compact and open.  $\widehat{A} \setminus \{\pi\}$  is open because  $\{\pi\}$  is closed by assumption. Since  $\widehat{A}$  is compact and satisfies the second axiom of countability [14], it suffices to show that  $\widehat{A} \setminus \{\pi\}$  is sequentially compact, [24, p. 138]. Let  $(\pi_n)_n$  be a sequence in  $\widehat{A} \setminus \{\pi\}$ . By compactness of  $\widehat{A}$  it contains a subsequence  $(\pi_{n_k})_k$  converging in  $\widehat{A}$ . If it converges to  $\pi \in S$ , then it also converges to some other point  $\eta \in \widehat{A} \setminus \{\pi\}$ , because  $\pi$  is shielded. Hence  $\widehat{A} \setminus \{\pi\}$  is also compact.  $\square$

**Corollary 2.11.** *Let  $G$  be a countable discrete group. If the trivial representation  $\iota \in \widehat{G}$  is shielded, then  $I(G)$  is not connective.*

*Proof.* Since  $\iota$  is a one-dimensional representation, it follows that  $\{\iota\}$  is closed in  $\widehat{G}$ . Thus, the statement follows from Lemma 2.10.  $\square$

### 3. CONNECTIVITY OF CRYSTALLOGRAPHIC GROUPS

It is known that there are precisely 10 closed flat 3-dimensional manifolds. Conway and Rossetti [7] call these manifolds platycosms (“flat universes”). The Hantzsche-Wendt manifold [17], or the didicosm in the terminology of [7], is the only platycosm with finite homology. Its fundamental group  $G$ , known as the Hantzsche-Wendt group, is generated by two elements  $x$  and  $y$  subject to two relations:

$$x^2yx^2 = y, \quad y^2xy^2 = x.$$

The group  $G$  is one of the classic torsion free 3-dimensional crystallographic groups, [17, 7]. It is useful to introduce the notation  $z = (xy)^{-1}$ .

A concrete realization of  $G$  as rigid motions of  $\mathbb{R}^3$  is given by the following transformations  $X, Y, Z$  that correspond to the group elements  $x, y$  and  $z$ .

$$X(\xi) = A\xi + a, \quad Y(\xi) = B\xi + b, \quad Z(\xi) = C\xi + c, \quad \xi \in \mathbb{R}^3,$$

where

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad B = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad C = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

and

$$a = \begin{pmatrix} 1/2 \\ 1/2 \\ 0 \end{pmatrix}, \quad b = \begin{pmatrix} 0 \\ 1/2 \\ 1/2 \end{pmatrix}, \quad c = \begin{pmatrix} 1/2 \\ 0 \\ 1/2 \end{pmatrix}.$$

The transformations  $X^2$ ,  $Y^2$  and  $Z^2$  are just translations by unit vectors in the positive directions of the coordinate axes.

One shows (independently of the previous concrete realization) that the elements  $x^2$ ,  $y^2$  and  $z^2$  commute. Moreover one has the following relations in  $G$ :

$$\begin{aligned} xx^2x^{-1} &= x^2, & xy^2x^{-1} &= y^{-2}, & xz^2x^{-1} &= z^{-2} \\ yx^2y^{-1} &= x^{-2}, & yy^2y^{-1} &= y^2, & yz^2y^{-1} &= z^{-2} \\ zx^2z^{-1} &= x^{-2}, & zy^2z^{-1} &= y^{-2}, & zz^2z^{-1} &= z^2 \end{aligned}$$

The subgroup  $N$  of  $G$  generated by  $x^2$ ,  $y^2$  and  $z^2$  is normal in  $G$  and it is isomorphic to  $\mathbb{Z}^3 \cong \mathbb{Z}x^2 \oplus \mathbb{Z}y^2 \oplus \mathbb{Z}z^2$ . Let  $q : G \rightarrow H = G/N$  denote the quotient map.

$$1 \longrightarrow N \longrightarrow G \xrightarrow{q} H \longrightarrow 1$$

$H$  is isomorphic to  $\mathbb{Z}/2 \oplus \mathbb{Z}/2$  with generators are  $q(x)$  and  $q(y)$ .

For later use, we will need the following identities that hold in  $G$ .

$$(2) \quad \begin{aligned} x^{-1}y &= yxy^2z^{-2} = zx^2z^{-2}, & x^{-1}z &= yz^2, \\ y^{-1}x &= zx^2, & y^{-1}z &= x(x^{-2}y^2). \end{aligned}$$

**3.1. Induced representations and the unitary dual of  $G$ .** Based on Corollary 2.11 we will show that  $I(G)$  for the Hantzsche-Wendt group  $G$  is not connective. This requires a thorough analysis of the spectrum  $\widehat{G}$ .

Our basic reference for this section is the book of Kaniuth and Taylor [22]. The unitary dual of  $G$  consists of unitary equivalence classes of irreducible unitary representations of  $G$  and is denoted by  $\widehat{G}$ .  $G$  acts on  $\widehat{N} \cong \mathbb{T}^3$  by  $g \cdot \chi \mapsto \chi(g \cdot g^{-1})$ . If we identify the character  $\chi \in \widehat{N}$  with the point  $(\chi(x^2), \chi(y^2), \chi(z^2)) = (u, v, w) \in \mathbb{T}^3$ , then the action of  $G$  is described as follows:

$$x \cdot (u, v, w) = (u, \bar{v}, \bar{w}), \quad y \cdot (u, v, w) = (\bar{u}, v, \bar{w}), \quad z \cdot (u, v, w) = (\bar{u}, \bar{v}, w).$$

The stabilizer of a character  $\chi$  is the subgroup  $G_\chi$  of  $G$  defined by  $G_\chi = \{g \in G : \chi(g \cdot g^{-1}) = \chi(\cdot)\}$ . It is clear that  $N \subset G_\chi$  and that there is a bijection from  $G/G_\chi$  onto the orbit of  $\chi$ . In particular, the orbits of the action of  $G$  on  $\widehat{N}$  can only have length 1, 2 or 4. Mackey has shown that each irreducible representation  $\pi \in \widehat{G}$  is supported by the orbit of some character  $\chi \in \widehat{N}$ , in the sense that the restriction of  $\pi$  to  $N$  is unitarily equivalent to some



multiple  $m_\pi$  of the direct sum of the characters in the orbit of  $\chi$ .

$$\pi|_N \sim m_\pi \bigoplus_{g \in G/G_\chi} \chi(g \cdot g^{-1}).$$

In the sum above  $g$  runs through a set of coset representatives.

Mackey's theory has a particularly nice form for virtually abelian discrete groups. Let  $\Omega \subset \widehat{N}$  be a subset which intersects each orbit of  $G$  exactly once. For each  $\chi \in \widehat{N}$ , let  $\widehat{G}_\chi$  be the unitary dual of the stabilizer group  $G_\chi$  and denote by  $\widehat{G}_\chi^{(\chi)}$  the subset of  $\widehat{G}_\chi$  consisting of classes of irreducible representations  $\sigma$  of  $G_\chi$  such the restriction of  $\sigma$  to  $N$  is unitarily equivalent to a multiple of  $\chi$ . Then, according to [22, Thm. 4.28]

**Theorem 3.1.**  $\widehat{G} = \left\{ \text{ind}_{G_\chi}^G(\sigma) : \sigma \in \widehat{G}_\chi^{(\chi)}, \chi \in \Omega \right\}$ .

Let  $\iota$  be the trivial representation of  $G$ . We will prove that  $\iota \in \widehat{G}$  is shielded by showing that any sequence  $(\pi_n)_n$  of points in  $\widehat{G} \setminus \{\iota\}$  that converges to  $\iota$  has a subsequence which is convergent to a point  $\eta \neq \iota$ .

Let  $R_\ell \subset \widehat{G}$  consist of those classes of irreducible representations which lie over  $\ell$ -orbits, i.e. the orbits of length  $\ell$ . Write  $\widehat{G}$  as the disjoint union  $\widehat{G} = R_1 \cup R_2 \cup R_4$ . It suffices to assume that all the elements  $\pi_n$  belong to the same subset  $R_\ell$ . We distinguish the three possible cases for  $\ell$ :

**1-orbits.** Consider the characters of  $N$  of the form  $\chi = (\varepsilon_1, \varepsilon_2, \varepsilon_3)$ , where  $\varepsilon_1, \varepsilon_2, \varepsilon_3 \in \{\pm 1\}$ . These are precisely the points in  $\widehat{N}$  which are fixed under the action of  $G$ . In other words  $G_\chi = G$ . Let  $(\pi_n)_n$  be a sequence of elements in  $R_1 \subset \widehat{G}$  and such that  $(\pi_n)_n$  is convergent to  $\iota$ . Since the restriction of  $\pi_n$  to  $N$  is a multiple of a character  $\chi_n = (\varepsilon_1(n), \varepsilon_2(n), \varepsilon_3(n))$ , it follows that  $\varepsilon_1(n) = \varepsilon_2(n) = \varepsilon_3(n) = 1$  for all sufficiently large  $n$  and hence since  $\pi_n$  is irreducible, there is  $m$  such that  $\pi_n = \iota$  for  $n \geq m$ . Hence there is no sequence in  $R_1 \setminus \{\iota\}$  which converges to  $\iota$ .

**2-orbits.** The characters  $\chi \in \widehat{N}$  with orbits of length two are those  $\chi = (u, v, w)$  where precisely only one of the coordinates is not equal to  $\pm 1$ . Let us argue first that if  $(\pi_n)_n$  is a sequence of elements in  $R_2 \subset \widehat{G}$  such that  $\pi_n$  lies over the orbit of  $\chi_n = (u_n, v_n, w_n)$  and  $(\pi_n)_n$  is convergent to  $\iota$ , then two of the coordinates  $u_n, v_n, w_n$  must be equal to 1 for all sufficiently large  $n$ .

Suppose that each  $\pi_n$  lies over the orbit of a character  $\chi_n$  of the form  $\chi_n = (u_n, \varepsilon_2(n), \varepsilon_3(n))$  where  $u_n \neq \pm 1$  and  $\varepsilon_2(n), \varepsilon_3(n) \in \{\pm 1\}$ . Then  $G_{\chi_n}$  is generated by  $x, y^2, z^2$  and  $\{e, y\}$  are coset representatives for  $G/G_{\chi_n}$ .

Since  $\pi_n|_N \sim m_n(\chi_n(\cdot) \oplus \chi_n(y \cdot y^{-1}))$ , it follows that

$$\pi_n(y^2) \sim m_n \begin{pmatrix} \varepsilon_2(n) & 0 \\ 0 & \varepsilon_2(n) \end{pmatrix}, \quad \pi_n(z^2) \sim m_n \begin{pmatrix} \varepsilon_3(n) & 0 \\ 0 & \varepsilon_3(n) \end{pmatrix}$$

and hence if  $(\pi_n)_n$  converges to  $\iota$ , then we must have  $\varepsilon_2(n) = \varepsilon_3(n) = 1$  for all sufficiently large  $n$ . The cases  $\chi_n = (\varepsilon_1, v_n, \varepsilon_3)$  and  $\chi_n = (\varepsilon_1, \varepsilon_2, w_n)$  are treated similarly.

In view of the discussion above, it suffices to focus on characters of  $N$  the form  $\chi = (u, 1, 1)$ . The orbit of  $\chi$  consists of two points,  $(u, 1, 1)$  and  $(\bar{u}, 1, 1)$ . The corresponding stabilizer  $G_\chi$  is generated by  $x, y^2$  and  $z^2$ . In particular  $G_\chi = N \cup xN$  and  $G = G_\chi \cup yG_\chi$ . The exact sequence

$$1 \longrightarrow N \longrightarrow G_\chi \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 1$$

does not split since  $G$  is torsion free. The quotient  $G/G_\chi$  is generated by the coset of  $y$ . Let  $\sigma \in \widehat{G}_\chi$  be an irreducible representation of  $G_\chi$  whose restriction to  $N$  is a multiple of  $\chi$ . Since  $\chi(y^2) = \chi(z^2) = 1$ , it follows that  $\sigma$  factors through  $G_\chi/\mathbb{Z}^2$ . Moreover we have a nontrivial central extension

$$1 \longrightarrow \mathbb{Z} \longrightarrow G_\chi/\mathbb{Z}^2 \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 1$$

where the normal subgroup is generated by the image of  $x^2$  under the map  $N \rightarrow N/\langle y^2, z^2 \rangle$  and the quotient group is generated by the image of  $q(x)$  under the map  $H \rightarrow H/\langle q(y) \rangle$ . Since  $G_\chi/\mathbb{Z}^2$  is an abelian group,  $\sigma$  must be a character such that  $\sigma(x)^2 = \sigma(x^2) = \chi(x^2) = u$ . Thus  $\sigma(x) = a \in \mathbb{T}$  with  $a^2 = u$ . Let us compute the representation  $\pi = \text{ind}_{G_\chi}^G(\sigma)$  of  $G$  induced by  $\sigma$ . It acts on the Hilbert space

$$H_\pi = \{\xi : G \rightarrow \mathbb{C} : \xi(gh) = \sigma(h^{-1})\xi(g), \quad g \in G, h \in G_\chi\}.$$

Since  $G = G_\chi \cup yG_\chi$ , we can identify  $H_\pi$  with  $\mathbb{C}^2$  via the isometry  $\xi \mapsto (\xi(e), \xi(y))$ . Then  $\pi(g)\xi = \xi(g^{-1}\cdot)$  can be described using (2) as follows:

$$\begin{aligned} \pi(x)\xi(e) &= \xi(x^{-1}) = \sigma(x)\xi(e) = a\xi(e) \\ \pi(x)\xi(y) &= \xi(x^{-1}y) = \xi(yxy^2z^{-2}) = \sigma(z^2y^{-2}x^{-1})\xi(y) = \bar{a}\xi(y) \\ \pi(y)\xi(e) &= \xi(y^{-1}) = \xi(y \cdot y^{-2}) = \sigma(y^2)\xi(y) = \xi(y) \\ \pi(y)\xi(y) &= \xi(y^{-1} \cdot y) = \xi(e) \end{aligned}$$

which produces the following matrices with respect to the basis given above:

$$(3) \quad \pi(x) = \begin{pmatrix} a & 0 \\ 0 & \bar{a} \end{pmatrix}, \quad \pi(y) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \pi(z) = \begin{pmatrix} 0 & a \\ \bar{a} & 0 \end{pmatrix}$$

Corresponding to the characters  $(1, v, 1)$  and  $(1, 1, w)$  we obtain the irreducible representations, where we use the isometries  $\xi \mapsto (\xi(e), \xi(x))$  and  $\xi \mapsto (\xi(e), \xi(y))$  respectively:

$$(4) \quad \pi(x) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \pi(y) = \begin{pmatrix} b & 0 \\ 0 & \bar{b} \end{pmatrix}, \quad \pi(z) = \begin{pmatrix} 0 & \bar{b} \\ b & 0 \end{pmatrix}, \quad b^2 = v,$$

and

$$(5) \quad \pi(x) = \begin{pmatrix} 0 & \bar{c} \\ c & 0 \end{pmatrix}, \quad \pi(y) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \pi(z) = \begin{pmatrix} c & 0 \\ 0 & \bar{c} \end{pmatrix}, \quad c^2 = w.$$

Let  $(\pi_n)_n$  be a sequence in  $R_2$  that converges to  $\iota$  in  $\widehat{G}$ . Arguing by symmetry, we may assume that each  $\pi_n$  is given by the formulas (3) corresponding to a sequence of points  $u_n \in \mathbb{T}$  with  $u_n \notin \{\pm 1\}$ . Since  $\pi_n \rightarrow \iota$  it follows from the equation (3) that  $u_n \rightarrow 1$ . Again from (3) we can compute the limits of the sequences  $\pi_n(x)$  and  $\pi_n(y)$  in  $U(2)$ . This gives the representation  $\pi: G \rightarrow U(2)$ :

$$\pi(x) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \pi(y) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

It is clear that  $\pi$  is a representation of  $G$  that factors through the left regular representation of  $\mathbb{Z}/2$ . Decompose  $\pi$  into a direct sum of characters  $\pi \sim \iota \oplus \eta$ . Then  $\eta$  is not equivalent to  $\iota$  and  $\pi_n \rightarrow \eta$  in  $\widehat{G}$ .

**4-orbits.** Let  $\chi = (u, v, w) \in \mathbb{T}^3$  be a character of  $N$  with  $u, v, w \notin \{\pm 1\}$ . Its orbit under the action of  $G$  consists of four points and  $G_\chi = N$ . Let us compute the representation  $\pi = \text{ind}_N^G(\chi)$  of  $G$  induced by  $\sigma$ . It acts on the Hilbert space  $H_\pi = \{\xi : G \rightarrow \mathbb{C} : \xi(gh) = \chi(h^{-1})\xi(g), \quad g \in G, h \in N\}$ . Thus one can identify  $H_\pi$  with  $\mathbb{C}^4$  via the isometry  $\xi \mapsto (\xi(e), \xi(x), \xi(y), \xi(z))$ . Using the identities (2), we verify that  $\pi(g)\xi = \xi(g^{-1}\cdot)$  is described as follows:

$$\begin{aligned} \pi(x)\xi(e) &= \xi(x^{-1}) = \xi(x \cdot x^{-2}) = \chi(x^2)\xi(x) = u\xi(x) \\ \pi(x)\xi(x) &= \xi(e) \\ \pi(x)\xi(y) &= \xi(x^{-1}y) = \xi(zx^2z^{-2}) = \chi(z^2x^{-2})\xi(z) = w\bar{u}\xi(z) \\ \pi(x)\xi(z) &= \xi(x^{-1}z) = \xi(yz^2) = \chi(z^{-2})\xi(y) = \bar{w}\xi(y) \\ \\ \pi(y)\xi(e) &= \xi(y^{-1}) = \xi(y \cdot y^{-2}) = \chi(y^2)\xi(y) = v\xi(y) \\ \pi(y)\xi(x) &= \xi(y^{-1}x) = \xi(zx^2) = \chi(x^{-2})\xi(z) = \bar{u}\xi(z) \\ \pi(y)\xi(y) &= \xi(e) \\ \pi(y)\xi(z) &= \xi(y^{-1}z) = \xi(x \cdot x^{-2}y^2) = \chi(y^{-2}x^2)\xi(x) = \bar{v}u\xi(x) \end{aligned}$$

producing the matrices:

$$(6) \quad \pi(x) = \begin{pmatrix} 0 & u & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & \bar{u}w \\ 0 & 0 & \bar{w} & 0 \end{pmatrix}, \quad \pi(y) = \begin{pmatrix} 0 & 0 & v & 0 \\ 0 & 0 & 0 & \bar{u} \\ 1 & 0 & 0 & 0 \\ 0 & \bar{v} & 0 & 0 \end{pmatrix}$$

It follows that

$$(7) \quad \pi(x^2) = \begin{pmatrix} u & 0 & 0 & 0 \\ 0 & u & 0 & 0 \\ 0 & 0 & \bar{u} & 0 \\ 0 & 0 & 0 & \bar{u} \end{pmatrix}, \quad \pi(y^2) = \begin{pmatrix} v & 0 & 0 & 0 \\ 0 & \bar{v} & 0 & 0 \\ 0 & 0 & v & 0 \\ 0 & 0 & 0 & \bar{v} \end{pmatrix}, \quad \pi(z^2) = \begin{pmatrix} w & 0 & 0 & 0 \\ 0 & \bar{w} & 0 & 0 \\ 0 & 0 & \bar{w} & 0 \\ 0 & 0 & 0 & w \end{pmatrix}$$

Let  $(\pi_n)_n$  be a sequence in  $R_4$  converging to  $\iota$  in  $\widehat{G}$ . Each  $\pi_n$  is given by the formulas (6) corresponding to a sequence of points  $(u_k, v_k, w_k) \in \mathbb{T}^3$  with  $u_k, v_k, w_k \notin \{\pm 1\}$ . Since  $\pi_n \rightarrow \iota$  it follows from equation (7) that  $u_k, v_k, w_k \rightarrow 1$ . Again from (6) we can compute the limits of the sequences  $\pi_n(x)$  and  $\pi_n(y)$  in the space of unitary operators  $U(4)$ . This gives the representation  $\pi: G \rightarrow U(4)$ :

$$\pi(x) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad \pi(y) = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

It is clear that  $\pi$  is a representation of  $G$  that factors through  $H = \mathbb{Z}/2 \times \mathbb{Z}/2$ . Decompose  $\pi$  into a direct sum of characters  $\eta_i$ . Since  $\pi$  is not equivalent to a multiple of the trivial representation, it follows that at least one of these characters is not equivalent to  $\iota$ . On the other hand  $\pi_n \rightarrow \eta_i$  in  $\widehat{G}$  for all  $i$ . Combining the above analysis with Corollary 2.11 we obtain immediately

**Corollary 3.2.** *If  $G$  is the Hantzsche-Wendt group, then  $I(G)$  is not connective.*

It was conjectured in [8] that if  $G$  is a torsion free discrete amenable group, then  $[[I(G), \mathcal{K}] \cong KK(I(G), \mathbb{C})$ . We argue now that this conjecture fails for the Hantzsche-Wendt group. Indeed this follows from the previous corollary in conjunction with the following lemma.

**Lemma 3.3.** *Let  $G$  be a residually finite torsion free discrete amenable group which admits a classifying space with finitely generated  $K$ -homology group  $K_1(BG)$ . Then  $[[I(G), \mathcal{K}] \cong KK(I(G), \mathbb{C})$  if and only if  $I(G)$  is connective.*

*Proof.* Suppose first that  $[[I(G), \mathcal{K}] \cong KK(I(G), \mathbb{C})$ . Since  $G$  is amenable and residually finite it follows that  $C^*(G)$  is residually finite dimensional. Since  $G$  is amenable,  $G$  satisfies the Baum-Connes conjecture and  $C^*(G)$  satisfies the UCT. In particular we have a short exact sequence

$$0 \rightarrow Ext^1(K_1(C^*(G)), \mathbb{Z}) \rightarrow KK(C^*(G), \mathbb{C}) \rightarrow \text{Hom}(K_0(C^*(G)), \mathbb{Z}) \rightarrow 0$$

Let  $\pi_n: C^*(G) \rightarrow M_{d(n)}(\mathbb{C})$  be a separating sequence of finite dimensional representations. The restriction of  $\pi_n$  to  $I(G)$  will also be denoted by  $\sigma_n$ . By [8, Prop. 3.2]  $(\pi_n)_* = d(n)\iota_*: K_0(C^*(G)) \rightarrow \mathbb{Z}$  and hence  $[\sigma_n] \in Ext^1(K_1(I(G)), \mathbb{Z}) \subset KK(I(G), \mathbb{C})$  is a torsion element since  $K_1(I(G))$  is finitely generated. After replacing  $\pi_n$  by a suitable multiple of itself we have

arranged that  $[\sigma_n] = 0$  in  $KK(I(G), \mathbb{C})$  and hence  $[[\sigma_n]] = 0$  in  $[[I(G), \mathcal{K}]]$ . Since the sequence  $(\sigma_n)$  separates the elements of  $I(G)$  it follows that  $I(G)$  is connective.

The converse is contained in the main result of [10] which shows that if  $A$  is a separable nuclear connective  $C^*$ -algebra, then  $[[A, \mathcal{K}]] \cong KK(A, \mathbb{C})$ .  $\square$

**3.2. Crystallographic groups with cyclic holonomy.** In this section we are going to show that torsion free crystallographic groups with cyclic holonomy are connective. Apart from this we isolate a lemma, which proves that  $I(G)$  for a group  $G$  which is a finite extension of a connective group always contains a “big” connective ideal. In particular, the lemma also holds for the Hantzsche-Wendt group.

The proof of both results uses some tools from the index theory of  $C^*$ -subalgebras. A reference is [37]. Let  $\Gamma$  and  $G$  be discrete groups and let  $H$  be a finite group. Suppose that they fit into an exact sequence

$$1 \longrightarrow \Gamma \longrightarrow G \xrightarrow{q} H \longrightarrow 1 .$$

Let  $E: C^*(G) \rightarrow C^*(\Gamma)$  be the faithful conditional expectation [37, Ex. 1.2.3] given on group elements by

$$E(g) = \begin{cases} g & \text{if } g \in \Gamma \\ 0 & \text{else} \end{cases} .$$

Choose a lift  $g_h \in G$  for each  $h \in H$ . The pairs  $(g_h^{-1}, g_h)$  form a quasi-basis in the sense of [37, Def. 1.2.2]. Let  $\mathcal{E} = C^*(G)$  considered as a right Hilbert  $C^*(\Gamma)$ -module, where the right action is induced by the inclusion  $C^*(\Gamma) \rightarrow C^*(G)$  and the inner product is given by  $\langle a, b \rangle = E(a^*b)$  [37, Sec. 2.1]. Note that  $\mathcal{E}$  is complete [37, Prop. 2.1.5]. The quasi-basis induces an isometric isomorphism of right Hilbert  $C^*(\Gamma)$ -modules  $u: \mathcal{E} \rightarrow \ell^2(H) \otimes C^*(\Gamma)$  with

$$u(a) = \sum_h \delta_h \otimes E(g_h a)$$

and inverse  $u^*: \ell^2(H) \otimes C^*(\Gamma) \rightarrow \mathcal{E}$  with  $u^*(\delta_h \otimes b) = g_h^{-1} b$ . Let  $\mathcal{L}_{C^*(\Gamma)}(\mathcal{E})$  be the bounded adjointable operators on  $\mathcal{E}$  and denote by  $\mathcal{K}_{C^*(\Gamma)}(\mathcal{E})$  the compact ones. Then we have  $\mathcal{L}_{C^*(\Gamma)}(\mathcal{E}) \cong \mathcal{K}_{C^*(\Gamma)}(\mathcal{E}) \cong \mathcal{K}(\ell^2(H)) \otimes C^*(\Gamma)$ . The left multiplication of  $C^*(G)$  on  $\mathcal{E}$  induces a  $*$ -homomorphism

$$\psi: C^*(G) \rightarrow \mathcal{K}(\ell^2(H)) \otimes C^*(\Gamma)$$

with matrix entries  $\psi_{h',h}(a) = E(g_{h'} a g_h^{-1})$ . Suppose we have  $a \in C^*(G)$  with  $\psi(a) = 0$ . Then

$$a = \frac{1}{|H|} \sum_{h,h'} g_{h'}^{-1} E(g_{h'} a g_h^{-1}) g_h = 0 .$$

Hence,  $\psi$  is injective.

**Lemma 3.4.** *Let  $\Gamma$  be a connective group and let  $H$  be a finite group. Suppose that the group  $G$  fits into a short exact sequence of the form*

$$1 \longrightarrow \Gamma \longrightarrow G \xrightarrow{q} H \longrightarrow 1 .$$

Then  $I(G, H) = \ker(I(q): I(G) \rightarrow I(H))$  is connective as well.

*Proof.* Let  $\iota: C^*(\Gamma) \rightarrow \mathbb{C}$  be the trivial representation and let  $\psi$  be the injective  $*$ -homomorphism constructed above. For all  $b \in C^*(\Gamma) \subset C^*(G)$  we have  $\psi_{h',h}(b) = \delta_{h',h} g_h b g_h^{-1}$ . In particular,  $\psi$  embeds the ideal  $J$  generated by  $I(\Gamma)$  into  $\ker(\text{id} \otimes \iota) = \mathcal{K}(\ell^2(H)) \otimes I(\Gamma)$ , which is connective. Hence  $J$  is connective as well. It is clear that  $J \subseteq I(G, H)$ . Let  $x \in I(G, H)$ . By the property of the quasi-basis

$$0 = q(x) = \sum_{h \in H} q(E(x g_h^{-1})) h \quad \Rightarrow \quad q(E(x g_h^{-1})) = 0 \quad \forall h \in H .$$

Since  $I(G, H) \cap C^*(\Gamma) = I(\Gamma)$  we obtain that  $E(x g_h^{-1}) \in I(\Gamma)$  for all  $h \in H$  and therefore  $x = \sum_h E(x g_h^{-1}) g_h \in J$ , hence  $J = I(G, H)$ .  $\square$

The proof of the second result uses an induction over the rank of the free abelian subgroup based on the following observation.

**Lemma 3.5.** *Let  $m > 1$  and let  $\Gamma$  and  $G$  be countable discrete groups that fit into the following short exact sequence*

$$1 \longrightarrow \Gamma \longrightarrow G \xrightarrow{\pi} \mathbb{Z}/m\mathbb{Z} \longrightarrow 1 .$$

Suppose that  $\Gamma$  is connective and there are group homomorphisms  $\varphi: G \rightarrow \mathbb{Z}$  and  $q: \mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$ , such that  $\pi = q \circ \varphi$ . Then  $G$  is connective as well.

*Proof.* Let  $\psi: C^*(G) \rightarrow \mathcal{K}(\ell^2(H)) \otimes C^*(\Gamma)$  be the injective  $*$ -homomorphism constructed above and let  $\iota: C^*(\Gamma) \rightarrow \mathbb{C}$  be the trivial representation. Observe that  $\rho = (\text{id} \otimes \iota) \circ \psi$  satisfies

$$\rho(g \gamma)_{h',h} = \iota(E(g_{h'} g \gamma g_h^{-1})) = \iota(E(g_{h'} g g_h^{-1}) g_h \gamma g_h^{-1}) = \iota(E(g_{h'} g g_h^{-1}))$$

for all  $g \in G$  and  $\gamma \in \Gamma$ . In particular,  $\rho$  factors through the  $*$ -homomorphism  $C^*(G) \rightarrow C^*(\mathbb{Z}/m\mathbb{Z})$  induced by  $\pi$ . By assumption this decomposes as

$$C^*(G) \xrightarrow{\varphi} C^*(\mathbb{Z}) \xrightarrow{q} C^*(\mathbb{Z}/m\mathbb{Z}) .$$

Altogether we obtain that  $\rho$  decomposes into a direct sum of one-dimensional representations, each of which is homotopic through representations to the trivial one. Hence, to show that  $G$  is connective, it suffices to construct a path through discrete asymptotic morphisms connecting a faithful morphism with a direct sum of copies of  $\rho$ .

Choose a path witnessing the connectivity of  $\Gamma$ , i.e. a discrete asymptotic morphism

$$H_n: C^*(\Gamma) \rightarrow C([0, 1]) \otimes M_n(\mathbb{C})$$

such that for  $H_n^{(t)} = \text{ev}_t \circ H_n: C^*(\Gamma) \rightarrow M_n(\mathbb{C})$  we have that  $H_n^{(0)}$  is faithful and  $H_n^{(1)}$  is a multiple of  $\iota$ . Then  $(\text{id}_{M_m(\mathbb{C})} \otimes H_n) \circ \psi$  has the desired properties. Hence,  $G$  is connective.  $\square$

We need the following elementary fact:

**Lemma 3.6.** *Let  $a, b > 1$  be integers and consider the exact sequence*

$$0 \longrightarrow \mathbb{Z}/a\mathbb{Z} \xrightarrow{\cdot b} \mathbb{Z}/ab\mathbb{Z} \xrightarrow{\pi} \mathbb{Z}/b\mathbb{Z} \longrightarrow 0$$

with  $\pi(x) = x \pmod{b}$ . Any generator of  $\mathbb{Z}/b\mathbb{Z}$  lifts to a generator of  $\mathbb{Z}/ab\mathbb{Z}$ .

*Proof.* Let  $\bar{y} \in \mathbb{Z}/b\mathbb{Z}$  be a generator and let  $y \in \{0, \dots, b-1\}$  be a representative. Let  $p_1, \dots, p_s$  be the distinct prime factors of  $ab$ , such that  $p_1, \dots, p_r$  for  $r \leq s$  are the ones not dividing  $y$  and  $p_{r+1}, \dots, p_s$  divide  $y$ . Since  $\gcd(y, b) = 1$ , the primes  $p_{r+1}, \dots, p_s$  do not divide  $b$ . Let  $x = y + p_1 \dots p_r b$ . We have for  $i \in \{1, \dots, r\}$  and  $j \in \{r+1, \dots, s\}$

$$\begin{aligned} x &\equiv y \not\equiv 0 \pmod{p_i}, \\ x &\equiv p_1 \dots p_r b \not\equiv 0 \pmod{p_j}. \end{aligned}$$

Hence  $\gcd(x, ab) = 1$  and  $x \in \mathbb{Z}/ab\mathbb{Z}$  is a generator with  $\pi(x) = \bar{y}$ .  $\square$

To start the induction we need the following lemma.

**Lemma 3.7.** *Let  $G$  be a countable torsion free discrete group, which fits into an exact sequence*

$$0 \longrightarrow \mathbb{Z} \longrightarrow G \longrightarrow \mathbb{Z}/m\mathbb{Z} \longrightarrow 0$$

Then  $G$  is isomorphic to  $\mathbb{Z}$ , hence in particular connective.

*Proof.* This can be proven by calculating  $H^2(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z})$  for all  $\mathbb{Z}/m\mathbb{Z}$ -module structures on  $\mathbb{Z}$ , but we give a direct argument here.

Let  $x \in G$  be a lift of  $1 \in \mathbb{Z}/m\mathbb{Z}$ . Then  $G$  is generated by  $x$  and  $\mathbb{Z}$ . Moreover,  $x^m \in \mathbb{Z} \cap Z(G)$ , where  $Z(G)$  denotes the center of  $G$ . We have  $\text{Aut}(\mathbb{Z}) \cong GL_1(\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$ . If  $t \in \mathbb{Z}$  denotes a generator, we therefore can only have  $xtx^{-1} = t^{-1}$  or  $xt = tx$ . Suppose the first is true, then

$$x^m = x x^m x^{-1} = x^{-m} \Rightarrow x^{2m} = e$$

contradicting that  $G$  is torsion free. Thus,  $t$  and  $x$  commute and  $x^m = t^n$  for some  $n \in \mathbb{Z}$ . Without loss of generality we can assume  $\gcd(m, n) = 1$ . Indeed, if  $m = m'\ell$  and  $n = n'\ell$  with  $\ell > 1$ , then  $(x^{m'}t^{-n'})^\ell = e$  and

therefore  $x^{m'} = t^{n'}$  also holds in  $G$ . Consider

$$\alpha: G \rightarrow \mathbb{Z} \quad ; \quad x^k t^\ell \mapsto kn + \ell m .$$

This is a well-defined group homomorphism, which is easily seen to be bijective as a consequence of  $\gcd(m, n) = 1$ .  $\square$

**Theorem 3.8.** *Let  $G$  be a countable, torsion free, discrete group, which fits into an exact sequence of the form*

$$0 \longrightarrow \mathbb{Z}^n \longrightarrow G \xrightarrow{\pi} \mathbb{Z}/m\mathbb{Z} \longrightarrow 0$$

for some  $n, m \in \mathbb{N}$ . Then  $G$  is connective.

*Proof.* This will be proven by induction over the rank of the free abelian subgroup. The case  $n = 1$  follows from Lemma 3.7.

Observe that  $Z(G) \neq \{e\}$ . Indeed, let  $x \in G$  be a lift of the generator of  $\mathbb{Z}/m\mathbb{Z}$ . Then  $G$  is generated by  $\mathbb{Z}^n$  and  $x$ . Moreover,  $x^m \neq e$  since  $G$  is torsion free and  $\pi(x^m)$  is trivial, hence  $x^m \in \mathbb{Z}^n$ . Thus,  $x^m$  commutes with  $\mathbb{Z}^n$  and  $x$ , hence with all elements of  $G$ , i.e.  $x^m \in Z(G)$ .

This implies that the transfer homomorphism  $T: G \rightarrow \mathbb{Z}^n$  associated to the finite index subgroup  $\mathbb{Z}^n$  is non-trivial. Therefore there exists a surjective group homomorphism  $\varphi: G \rightarrow \mathbb{Z}$ . Let  $q: \mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$  be the canonical quotient homomorphism and let  $\bar{\varphi} = q \circ \varphi$ . Let  $H = \ker(\varphi)$ . We have the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & H & \longrightarrow & \mathbb{Z}/a\mathbb{Z} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathbb{Z}^n & \longrightarrow & G & \xrightarrow{\pi} & \mathbb{Z}/m\mathbb{Z} \longrightarrow 0 \\ & & \downarrow & & \downarrow \varphi & & \downarrow \\ 0 & \longrightarrow & b\mathbb{Z} & \longrightarrow & \mathbb{Z} & \xrightarrow{\pi'} & \mathbb{Z}/b\mathbb{Z} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & \end{array}$$

The value of  $a$  is chosen in such a way that  $\text{Im}(\pi|_H) \cong \mathbb{Z}/a\mathbb{Z}$  and  $b$  satisfies  $m = ab$ . The homomorphism  $\pi'$  is surjective since  $\pi$  and  $\mathbb{Z}/m\mathbb{Z} \rightarrow \mathbb{Z}/b\mathbb{Z}$  are. The vertical arrow on the left hand side is induced by  $\varphi|_{\mathbb{Z}^n}$ .

Suppose  $H \subset \ker(\pi) = \mathbb{Z}^n$ . Then  $a = 1$ ,  $b = m$  and  $\pi = \pi' \circ \varphi$ . By Lemma 3.5,  $G$  is then connective. So we may assume  $a > 1$ .



We claim that there is an element  $g \in G$  such that  $\varphi(g) = 1$  and  $\pi(g)$  is a generator of  $\mathbb{Z}/m\mathbb{Z}$ . This is constructed as follows: If  $b = 1$ , we can choose  $g \in G$ , such that  $\varphi(g) = 1$  and modify it by an element in  $H$  to achieve that  $\pi(g)$  becomes a generator. Otherwise, choose  $g' \in G$  such that  $\varphi(g') = 1$  and note that  $\pi'(\varphi(g'))$  is a generator of  $\mathbb{Z}/b\mathbb{Z}$  by surjectivity. We can lift  $\pi'(\varphi(g'))$  to a generator  $x \in \mathbb{Z}/m\mathbb{Z}$  by Lemma 3.6. Note that  $\pi(g') - x \in \mathbb{Z}/a\mathbb{Z}$  and lift this difference to an element  $h \in H$ . Let  $g = g' h^{-1}$ . Then  $\varphi(g) = \varphi(g') = 1$  and  $\pi(g) = \pi(g') - \pi(h) = \pi(g') - \pi(g') + x = x$ .

Let  $G' = \ker(\bar{\varphi}) = \{g \in G \mid \varphi(g) = \ell \cdot m \text{ for } \ell \in \mathbb{Z}\} \supset H$ . Hence, the following diagram has exact rows.

$$(8) \quad \begin{array}{ccccccc} 1 & \longrightarrow & G' & \longrightarrow & G & \xrightarrow{\bar{\varphi}} & \mathbb{Z}/m\mathbb{Z} \longrightarrow 0 \\ & & \downarrow \varphi|_{G'} & & \downarrow \varphi & & \downarrow = \\ 1 & \longrightarrow & m\mathbb{Z} & \longrightarrow & \mathbb{Z} & \xrightarrow{q} & \mathbb{Z}/m\mathbb{Z} \longrightarrow 0 \end{array}$$

The group  $G$  is generated by  $\mathbb{Z}^n$  and the element  $g$  constructed above. We have  $g^m \in \mathbb{Z}^n \cap Z(G)$  and  $\varphi(g^m) = m$ . In particular,  $g^m \in Z(G')$ . Let

$$\psi: H \times \mathbb{Z} \rightarrow G' \quad ; \quad (h, k) \mapsto h \cdot g^{mk}.$$

This is a group homomorphism, since  $g^m$  is central and it fits into the commutative diagram with exact upper and lower part

$$\begin{array}{ccccc} & & G' & & \\ & \nearrow & \uparrow & \searrow \varphi|_{G'}/m & \\ 1 & \longrightarrow & H & & \mathbb{Z} \longrightarrow 0 \\ & \searrow i & \downarrow \psi & \nearrow \text{pr} & \\ & & H \times \mathbb{Z} & & \end{array}$$

proving that  $\psi$  is in fact an isomorphism. By the upper row in diagram (8) and Lemma 3.5, the connectivity of  $G$  follows if  $H \times \mathbb{Z}$ , hence  $H$ , is connective [10, Thm. 4.1]. But  $H$  fits into a short exact sequence of the form

$$0 \rightarrow A \rightarrow H \rightarrow \mathbb{Z}/a\mathbb{Z} \rightarrow 0$$

where  $A$  is the free abelian kernel of the nonzero homomorphism  $\mathbb{Z}^n \rightarrow b\mathbb{Z}$  from above, which has rank  $(n - 1)$ . This completes the induction step.  $\square$

#### 4. CONNECTIVITY OF LIE GROUP $C^*$ -ALGEBRAS

In this section we determine which linear connected nilpotent Lie groups and which linear connected reductive Lie groups have connective reduced  $C^*$ -algebras. Let us recall that nilpotent connected Lie groups are liminary

as shown by Dixmier [13] and Kirillov [26] and semisimple connected Lie groups are liminary as shown by Harish-Chandra [18].

**4.1. Solvable and nilpotent Lie groups.** A locally compact group  $N$  is compactly generated if  $N = \bigcup_n V^n$  for some compact subset  $V$  of  $N$ . Every connected locally compact group is automatically compactly generated. The structure of abelian compactly generated locally compact groups is known. If  $N$  is such a group, then  $N \cong \mathbb{R}^n \times \mathbb{Z}^m \times K$  for integers  $n, m \geq 0$  and  $K$  a compact group, [12, Thm. 4.4.2].

**Proposition 4.1.** *If  $G$  is a second countable locally compact amenable group (for example a solvable Lie group) whose center contains a noncompact closed connected subgroup, then  $C^*(G)$  is connective.*

*Proof.* Let  $N$  be a noncompact closed connected subgroup of  $Z(G)$ . Then, by the structure theorem quoted above,  $N$  must have a closed subgroup isomorphic to  $\mathbb{R}$ . Consider the central extension:

$$0 \rightarrow \mathbb{R} \rightarrow G \rightarrow H \rightarrow 0.$$

Since  $G$  is amenable, by [28, Thm. 1.2] (as explained in [15, Lemma 6.3]),  $C^*(G)$  has the structure of a continuous field of  $C^*$ -algebras over  $\widehat{\mathbb{R}} \cong \mathbb{R}$ . The desired conclusion follows from Cor. 2.6 since  $\mathbb{R}$  has no compact open subsets.  $\square$

**Example 4.2.** We give here two examples that complement Proposition 4.1.

- (i) Simply connected solvable Lie groups can have discrete noncompact centers. This is case for  $G = \mathbb{C} \rtimes_{\alpha} \mathbb{R}$  where  $\alpha : \mathbb{R} \rightarrow \text{Aut}(\mathbb{C})$  is defined by  $\alpha(t)(z) = e^{it}z$  for  $t \in \mathbb{R}$  and  $z \in \mathbb{C}$ . In this case  $Z(G) = \{0\} \times 2\pi\mathbb{Z}$ . Nevertheless in this case  $C^*(G)$  is connective. Consider the extension

$$0 \rightarrow Z(G) \rightarrow G \rightarrow G/Z(G) \cong \mathbb{C} \times \mathbb{T} \rightarrow 0.$$

Then  $C^*(G)$  is a continuous  $C(\mathbb{T})$ -algebra whose fiber at 1 is the algebra  $C^*(\mathbb{C} \times \mathbb{T})$ . Since  $C^*(\mathbb{C} \times \mathbb{T}) \cong C_0(\mathbb{R}^2) \otimes c_0(\mathbb{Z})$  is connective, so is  $C^*(G)$ .

- (ii) Both the real and the complex “ax+b” groups

$$G = \left\{ \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} : a \in F^{\times}, b \in F \right\}$$

where  $F = \mathbb{R}$  or  $F = \mathbb{C}$  are solvable with trivial center and their  $C^*$ -algebras contain a copy of the compacts  $\mathcal{K}$ , see [32], and so they are not connective.

**Theorem 4.3.** *Let  $G$  be a (real or complex) linear connected nilpotent Lie group. Then  $C^*(G)$  is connective if and only if  $G$  is not compact.*

*Proof.* We view  $G$  as a real Lie group. By [35, Chap. 2, Thm. 7.3], if  $G$  is a linear connected nilpotent Lie group, then  $G$  decomposes as a direct product  $G = T \times N$  of a torus  $T$  and a simply connected nilpotent group  $N$ . If  $G$  is noncompact, then  $N$  is nontrivial and so the center of  $G$  is given by  $Z(G) = T \times Z(N)$ , where the center  $Z(N)$  of  $N$  is isomorphic to  $\mathbb{R}^n$  for some  $n \geq 1$ . We conclude the proof by applying Proposition 4.1.  $\square$

**Remark 4.4.** It is not true that a liminary (CCR)  $C^*$ -algebra is connective if and only if does not have nonzero projections. Indeed

$$A = \left\{ f \in C([0, 1], M_2(\mathbb{C})) : f(0) = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix}, f(1) = \begin{pmatrix} \lambda & 0 \\ 0 & 0 \end{pmatrix}, \lambda \in \mathbb{C} \right\}$$

does not contain nonzero projections but is not connective since  $\text{Prim}(A)$  is homeomorphic to a circle  $S^1$  and hence it is compact (and open in itself).

**4.2. Reductive Lie groups.** A linear connected reductive group  $G$  is a closed group of linear or complex matrices that is closed under conjugate transpose. In other words  $G$  is a closed and selfadjoint subgroup of the general linear group over either  $\mathbb{R}$  or  $\mathbb{C}$ . A linear connected semisimple group is a linear connected reductive group with finite center [27].

Say  $G \subset GL(n, \mathbb{R})$  or  $GL(n, \mathbb{C})$ . Define  $K = G \cap O(n)$ , or  $K = G \cap U(n)$  in the complex case. If  $G$  is linear connected reductive, then  $K$  is compact, connected and is a maximal compact subgroup of  $G$  [27, Prop.1.2].

Let  $G = KAN$  be the Iwasawa decomposition of the linear connected semisimple Lie group  $G$ .  $A$  is abelian and  $N$  is nilpotent and both are closed simply connected subgroups of  $G$ , [27, Thm. 5.12].

First we consider the case of complex Lie groups.

**Theorem 4.5.** *If  $G$  is a linear connected complex semisimple Lie group, then  $C_r^*(G)$  is connective if and only if  $G$  is not compact.*

*Proof.* If  $G$  is compact,  $C_r^*(G)$  is isomorphic with a direct sum of matrix algebras and hence it is not connective as it contains nonzero projections.

Conversely, suppose now that  $G$  is a non-compact linear connected semisimple complex Lie group. Note that from the Cartan decomposition  $G = KAK$  [27, Thm. 5.20] it follows that if  $G$  is non-compact, then so is  $A$  and therefore  $A \cong \mathbb{R}^n$ , for some  $n \geq 1$ .

Let  $M$  be the centralizer of  $A$  in  $K$ . By Lemma 3.3 and Proposition 4.1 of [30], if  $G$  is a linear connected complex semisimple Lie group, then

$$C_r^*(G) \subset C_0(\widehat{M} \times \widehat{A}, \mathcal{K}).$$

Since  $\widehat{M} \times \widehat{A} \cong \widehat{M} \times \mathbb{R}^n$  does not have nonempty compact open subsets, it follows from Proposition 2.7(ii) that  $C_0(\widehat{M} \times \widehat{A}, \mathcal{K})$  is connective. This completes the proof since connectivity passes to  $C^*$ -subalgebras.  $\square$

Next we consider the case of linear connected real reductive Lie groups. An element  $g \in G$  is semisimple if it can be diagonalized over  $\mathbb{C}$  when viewed as a matrix  $g \in M_n(\mathbb{C})$ .

A closed subgroup  $H$  of  $G$  is called a Cartan subgroup if it is a maximal abelian subgroup consisting of semisimple elements, [21, p.67]. If  $G$  is either compact or a complex Lie group, then all Cartan subgroups of  $G$  are connected and they are conjugated inside  $G$ . In the general case  $G$  has finitely many Cartan subgroups up to conjugacy and Cartan subgroups can have finitely many connected components.

We denote by  $\widehat{G}_d \subset \widehat{G}$  the discrete series representations. It consists of unitary equivalence classes of square-integrable representations

$$\sigma : G \rightarrow U(H_\sigma).$$

Harish-Chandra has shown that the discrete series representations of a semi-simple Lie group  $G$  are parametrized by compact Cartan subgroups and in particular  $G$  has discrete series representations if and only if it has a compact Cartan subgroup, [19, 20].

We recall the following facts from [21, p.72]. Let  $H$  be a Cartan subgroup of  $G$ . Then  $H$  decomposes as a direct product  $H = TA = T \times A$ , where  $T$  is a compact group and  $A$  is a vector group isomorphic to  $\mathbb{R}^n$  for  $n \geq 0$ . The case  $n = 0$  occurs when  $H$  is a compact Cartan subgroup. The centralizer of  $A$  in  $G$  denoted by

$$L = C_G(A) = \{g \in G : ga = ag, \forall a \in A\}$$

is a Levi subgroup of  $G$ . This means that there is there is a parabolic subgroup of  $G$  of the form  $P = LN$  (not unique) with  $L$  as a Levi subgroup. Since  $A$  is central in  $L$ ,  $H$  is a relatively compact Cartan subgroup of  $L$ , i.e.  $H/Z(L)$  is compact. This implies that  $L$  has discrete series representations. Thus  $P$  is a cuspidal parabolic subgroup of  $G$ .

We can further decompose  $L = MA$  and so, one has a Langlands decomposition

$$P = MAN = MA \ltimes N,$$

with  $N$  a unipotent group. If  $H$  is a compact Cartan subgroup, then  $L = P = G$  by [21, p.72].

We will write  $P = M_P A_P N_P$  whenever we want emphasize the components of  $P$ .

Given  $\sigma \in \widehat{M}_d$  and  $\omega \in \widehat{A}_d$  one considers the unitary principal series representations of  $G$  associated to  $P$  (also called the  $P$ -principal series):

$$\text{Ind}_P^G(\sigma \otimes \omega \otimes 1_N)$$

where  $1_N$  is the trivial representation of  $N$ .

Consider two pairs  $(P_i, \sigma_i)$ ,  $i = 1, 2$  consisting of parabolic subgroups of  $G$  and irreducible square-integrable unitary representations of the subgroups  $M_i$ ,  $i = 1, 2$ . We say that the pairs are *associated* if there is  $g \in G$  such that  $gP_1g^{-1} = P_2$  and  $\sigma_1(g \cdot g^{-1})$  is unitarily equivalent to  $\sigma_2$ . This is an equivalence relation [5][Def.5.2]. We denote by  $[P, \sigma]$  the equivalence class of the pair  $(P, \sigma)$ .

The following statement is based on the calculation of  $C_r^*(G)$  by A. Wassermann although we don't really use the full strength of his results. An expanded treatment of the structure of  $C_r^*(G)$  appears in [5].

Let  $J(G) = \bigcap_{\pi \in \widehat{G}_d} \ker(\pi) \subset C_r^*(G)$  be the common kernel of the discrete series representations. It follows that the  $K$ -homology of  $C_r^*(G)$  can be described in terms of homotopy classes asymptotic morphisms  $C_r^*(G) \rightarrow \mathcal{K}$  which factor through  $J(G)$  and discrete series representations.

**Theorem 4.6.** *Let  $G$  be a linear connected real reductive Lie group. Then  $C_r^*(G) \cong J(G) \oplus \bigoplus_{\sigma \in \widehat{G}_d} K(H_\sigma)$  and  $J(G)$  is a connective liminary  $C^*$ -algebra. Moreover, the following assertions are equivalent:*

- (i)  $C_r^*(G)$  is connective
- (ii)  $G$  does not have discrete series representations
- (iii)  $G$  does not have a compact Cartan subgroup
- (iv) There are no nonzero projections in  $C_r^*(G)$ .

*Proof.* As explained in [36, p.560], [5, p.1306] the reduced  $C^*$ -algebra of a linear reductive connected Lie group admits an embedding

$$C_r^*(G) \hookrightarrow \bigoplus_{[P, \sigma]} C_0(\widehat{A}_P, \mathcal{K}(H_\sigma)).$$

It is important to emphasize that if  $G$  has a compact Cartan subgroup, then  $G$  is itself one of the parabolic subgroups and we have:

$$C_r^*(G) \hookrightarrow \bigoplus_{\sigma \in \widehat{G}_d} \mathcal{K}(H_\sigma) \oplus \bigoplus_{[P, \sigma]} C_0(\widehat{A}_P, \mathcal{K}(H_\sigma)),$$

where the second direct sum involves proper parabolic subgroups and hence  $\dim(\widehat{A}_P) > 0$ . Moreover by [36], [5]:

$$(9) \quad C_r^*(G) \cong J(G) \oplus \bigoplus_{\sigma \in \widehat{G}_d} \mathcal{K}(H_\sigma)$$

where

$$J(G) \hookrightarrow \bigoplus_{[P, \sigma]} C_0(\widehat{A}_P, \mathcal{K}(H_\sigma)).$$

Hence,  $J(G)$  is connective being a subalgebra of a connective  $C^*$ -algebra. The first part of the statement follows now from the decomposition (9).

The equivalence  $(ii) \Leftrightarrow (iii)$  is Harish-Chandra's result mentioned earlier. In view of the decomposition (9),  $(ii)$  implies that  $C_r^*(G) = J(G)$  and hence  $(i)$  since  $J(G)$  is always connective. Connective  $C^*$ -algebras do not contain nonzero projections and hence  $(i) \Rightarrow (iv)$ . Finally by using (9) again, we see that  $(iv) \Rightarrow (ii)$  since  $\mathcal{K}(H_\sigma)$  contains nonzero projections if  $H_\sigma \neq 0$ .  $\square$

**4.3. A remark on full  $C^*$ -algebras of Lie groups.** The full  $C^*$ -algebra  $C^*(G)$  of a property (T) Lie group  $G$  contains nonzero projections and hence it is not connective, see [34]. Nevertheless, inspection of several classes of examples indicates that  $C^*(G)$  has interesting connective ideals that arise naturally from the representation theory of  $G$ . We postpone a detailed discussion of what is known for another time but would like to mention two examples.

If  $G$  is a connected semisimple Lie group with finite center, then  $C^*(G)$  is liminary (or CCR), see [38, p.115].

**Proposition 4.7.** (a)  $C^*(SL_2(\mathbb{C}))$  is connective.

(b)  $C^*(SL_3(\mathbb{C})) = I(SL_3(\mathbb{C})) \oplus \mathbb{C}$  and  $I(SL_3(\mathbb{C}))$  is connective.

*Proof.* (a)  $C^*(SL_2(\mathbb{C}))$  was computed by Fell [16, Thm. 5.4]. We describe now his result. Let  $Z$  be the subspace of  $\mathbb{R}^2$  defined by  $Z = \bigcup_{n=0}^{\infty} \{n\} \times L_n$  where  $L_0 = (-1, \infty)$  and  $L_n = (-\infty, \infty)$  for all  $n \geq 1$ . We endow  $Z$  with the induced topology from  $\mathbb{R}^2$ . Let  $H_0$  be a separable infinite dimensional Hilbert space, let  $H = H_0 \oplus \mathbb{C}$  and fix a unitary operator  $V : H_0 \rightarrow H$ . Then  $C^*(SL_2(\mathbb{C}))$  is isomorphic to

$$\{F \in C_0(Z, \mathcal{K}(H)) : F(0, -1) = V^*F(2, 0)V \oplus \lambda, \text{ for some } \lambda \in \mathbb{C}\}$$

Since  $Z$  has no nonempty open compact subsets it follows that  $C_0(Z, \mathcal{K}(H))$  is connective and therefore so is its subalgebra  $C^*(SL_2(\mathbb{C}))$ .

(b) This will be obtained as a consequence of the following result on the structure of  $C^*(SL_3(\mathbb{C}))$  obtained by Pierrot [31]. Let  $G = SL_3(\mathbb{C})$  and denote by  $\lambda_G : C^*(G) \rightarrow C_r^*(G)$  the morphism induced by the left regular representation and by  $\iota_G : C^*(G) \rightarrow \mathbb{C}$  the trivial representation. Pierrot proved that the kernel  $J$  of the morphism  $\lambda_G \oplus \iota_G : C^*(G) \rightarrow C_r^*(G) \oplus \mathbb{C}$  is a contractible  $C^*$ -algebra. The representation  $\iota_G$  is isolated since  $G$  has property (T). Therefore we there is an exact sequence

$$0 \rightarrow J \rightarrow I(G) \rightarrow C_r^*(G) \rightarrow 0$$

where  $J$  is contractible and  $C_r^*(G)$  is connective by Theorem 4.5. We conclude that  $I(G)$  is connective by applying Theorem 2.4.  $\square$

## REFERENCES

- [1] B. Blackadar and J. Cuntz. The structure of stable algebraically simple  $C^*$ -algebras. *American J. Math.*, 104:813–822, 1982. [2](#)
- [2] É. Blanchard. Subtriviality of continuous fields of nuclear  $C^*$ -algebras. *J. Reine Angew. Math.*, 489:133–149, 1997. [5](#)
- [3] É. Blanchard and E. Kirchberg. Non-simple purely infinite  $C^*$ -algebras: the Hausdorff case. *J. Funct. Anal.*, 207:461–513, 2004. [6](#)
- [4] N. P. Brown and M. Dadarlat. Extensions of quasidiagonal  $C^*$ -algebras and  $K$ -theory. In *Operator algebras and applications*, volume 38 of *Adv. Stud. Pure Math.*, pages 65–84. Math. Soc. Japan, Tokyo, 2004. [5](#)
- [5] P. Clare, T. Crisp, and N. Higson. Parabolic induction and restriction via  $C^*$ -algebras and Hilbert  $C^*$ -modules. *Compos. Math.*, 152(6):1286–1318, 2016. ISSN 0010-437X. doi: 10.1112/S0010437X15007824. URL <http://dx.doi.org/10.1112/S0010437X15007824>. [21](#)
- [6] A. Connes and N. Higson. Déformations, morphismes asymptotiques et  $K$ -théorie bivariante. *C. R. Acad. Sci. Paris Sér. I Math.*, 311(2): 101–106, 1990. [1](#)
- [7] J. H. Conway and J. P. Rossetti. Describing the platycosms. Preprint 2003, arXiv:math/0311476v1. [7](#)
- [8] M. Dadarlat. Group quasi-representations and almost flat bundles. *J. Noncommut. Geom.*, 8(1):163–178, 2014. [2](#), [12](#)
- [9] M. Dadarlat and T. A. Loring.  $K$ -homology, asymptotic representations, and unsuspending  $E$ -theory. *J. Funct. Anal.*, 126(2):367–383, 1994. [1](#)
- [10] M. Dadarlat and U. Pennig. Deformations of nilpotent groups and homotopy symmetric  $C^*$ -algebras. *Mathem. Ann.*, 2016. ISSN 1432-1807. doi: 10.1007/s00208-016-1379-0. URL <http://dx.doi.org/10.1007/s00208-016-1379-0>. [1](#), [2](#), [3](#), [4](#), [5](#), [13](#), [17](#)
- [11] M. Dadarlat, U. Pennig, and A. Schneider. Deformations of wreath products. to appear in *Bull. London. Math. Soc.*, Preprint (2016), arXiv:1609.00604, 2016. [2](#)
- [12] A. Deitmar and S. Echterhoff. *Principles of harmonic analysis*. Universitext. Springer, Cham, second edition, 2014. ISBN 978-3-319-05791-0; 978-3-319-05792-7. doi: 10.1007/978-3-319-05792-7. URL <http://dx.doi.org/10.1007/978-3-319-05792-7>. [18](#)
- [13] J. Dixmier. Sur les représentations unitaires des groupes de Lie nilpotents. V. *Bull. Soc. Math. France*, 87:65–79, 1959. ISSN 0037-9484. [18](#)
- [14] J. Dixmier.  *$C^*$ -algebras*. North Holland, Amsterdam, 1982. [6](#), [7](#)

- [15] S. Echterhoff and D. P. Williams. Crossed products by  $C_0(X)$ -actions. *J. Funct. Anal.*, 158(1):113–151, 1998. ISSN 0022-1236. doi: 10.1006/jfan.1998.3295. URL <http://dx.doi.org/10.1006/jfan.1998.3295>. 18
- [16] J. M. G. Fell. The structure of algebras of operator fields. *Acta Math.*, 106:233–280, 1961. 6, 22
- [17] W. Hantzsche and H. Wendt. Dreidimensionale euklidische Raumformen. *Math. Ann.*, 110(1):593–611, 1935. ISSN 0025-5831. doi: 10.1007/BF01448045. URL <http://dx.doi.org/10.1007/BF01448045>. 7
- [18] Harish-Chandra. Representations of semisimple Lie groups. III. *Trans. Amer. Math. Soc.*, 76:234–253, 1954. ISSN 0002-9947. 18
- [19] Harish-Chandra. Discrete series for semisimple Lie groups. I. Construction of invariant eigendistributions. *Acta Math.*, 113:241–318, 1965. ISSN 0001-5962. 20
- [20] Harish-Chandra. Discrete series for semisimple Lie groups. II. Explicit determination of the characters. *Acta Math.*, 116:1–111, 1966. ISSN 0001-5962. 20
- [21] R. A. Herb. An elementary introduction to Harish-Chandra’s work. In *The mathematical legacy of Harish-Chandra (Baltimore, MD, 1998)*, volume 68 of *Proc. Sympos. Pure Math.*, pages 59–75. Amer. Math. Soc., Providence, RI, 2000. doi: 10.1090/pspum/068/1767892. URL <http://dx.doi.org/10.1090/pspum/068/1767892>. 20
- [22] E. Kaniuth and K. F. Taylor. *Induced representations of locally compact groups*, volume 197 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2013. ISBN 978-0-521-76226-7. 8, 9
- [23] G. G. Kasparov. The operator  $K$ -functor and extensions of  $C^*$ -algebras. *Izv. Akad. Nauk SSSR Ser. Mat.*, 44(3):571–636, 719, 1980. ISSN 0373-2436. 5
- [24] J. L. Kelley. *General topology*. D. Van Nostrand Company, Inc., Toronto-New York-London, 1955. 7
- [25] E. Kirchberg. Das nicht-kommutative Michael-Auswahlprinzip und die Klassifikation nicht-einfacher Algebren. In  *$C^*$ -algebras*, pages 92–141, Berlin, 2000. Springer. (Münster, 1999). 6
- [26] A. A. Kirillov. Unitary representations of nilpotent Lie groups. *Uspehi Mat. Nauk*, 17(4 (106)):57–110, 1962. ISSN 0042-1316. 18
- [27] A. W. Knap. *Representation theory of semisimple groups*, volume 36 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1986. ISBN 0-691-08401-7. An overview based on examples. 19
- [28] J. A. Packer and I. Raeburn. On the structure of twisted group  $C^*$ -algebras. *Trans. Amer. Math. Soc.*, 334(2):685–718, 1992. ISSN 0002-9947. doi: 10.2307/2154478. URL <http://dx.doi.org/10.2307/>



2154478. 18

- [29] C. Pasnicu and M. Rørdam. Purely infinite  $C^*$ -algebras of real rank zero. *J. Reine Angew. Math.*, 613:51–73, 2007. ISSN 0075-4102. doi: 10.1515/CRELLE.2007.091. URL <http://dx.doi.org/10.1515/CRELLE.2007.091>. 2, 6, 7
- [30] M. G. Pennington and R. J. Plymen. The Dirac operator and the principal series for complex semisimple Lie groups. *J. Funct. Anal.*, 53(3): 269–286, 1983. ISSN 0022-1236. doi: 10.1016/0022-1236(83)90035-6. URL [http://dx.doi.org/10.1016/0022-1236\(83\)90035-6](http://dx.doi.org/10.1016/0022-1236(83)90035-6). 19
- [31] F. Pierrot. La  $K$ -théorie de la  $C^*$ -algèbre pleine de  $SL_3(\mathbb{C})$ . *J. Funct. Anal.*, 193(2):261–277, 2002. ISSN 0022-1236. doi: 10.1006/jfan.2001.3907. URL <http://dx.doi.org/10.1006/jfan.2001.3907>. 22
- [32] J. Rosenberg. The  $C^*$ -algebras of some real and  $p$ -adic solvable groups. *Pacific J. Math.*, 65(1):175–192, 1976. ISSN 0030-8730. 18
- [33] K. Thomsen. Discrete asymptotic homomorphisms in  $E$ -theory and  $KK$ -theory. *Math. Scand.*, 92(1):103–128, 2003. ISSN 0025-5521. 1
- [34] A. Valette. Projections in full  $C^*$ -algebras of semisimple Lie groups. *Math. Ann.*, 294(2):277–287, 1992. ISSN 0025-5831. doi: 10.1007/BF01934326. URL <http://dx.doi.org/10.1007/BF01934326>. 22
- [35] È. B. Vinberg, editor. *Lie groups and Lie algebras, III*, volume 41 of *Encyclopaedia of Mathematical Sciences*. 19
- [36] A. Wassermann. Une démonstration de la conjecture de Connes-Kasparov pour les groupes de Lie linéaires connexes réductifs. *C. R. Acad. Sci. Paris Sér. I Math.*, 304(18):559–562, 1987. ISSN 0249-6291. 21
- [37] Y. Watatani. Index for  $C^*$ -subalgebras. *Mem. Amer. Math. Soc.*, 83 (424):vi+117, 1990. ISSN 0065-9266. doi: 10.1090/memo/0424. URL <http://dx.doi.org/10.1090/memo/0424>. 13
- [38] J. A. Wolf, M. Cahen, and M. De Wilde, editors. *Harmonic analysis and representations of semisimple Lie groups*, volume 5 of *Mathematical Physics and Applied Mathematics*. 22

MD: DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, USA

*E-mail address:* mdd@purdue.edu

UP: SCHOOL OF MATHEMATICS, CARDIFF UNIVERSITY, SENGHENNYDD ROAD, CARDIFF CF24 4AG, UK

*E-mail address:* PennigU@cardiff.ac.uk