

# COMPUTING COHOMOLOGY GROUPS THAT CLASSIFY BUNDLES OF STRONGLY SELF-ABSORBING $C^*$ -ALGEBRAS

MARIUS DADARLAT, JAMES E. MCCLURE, AND ULRICH PENNIG

ABSTRACT. Locally trivial bundles of  $C^*$ -algebras with fibre  $D \otimes \mathcal{K}$  for a strongly self-absorbing  $C^*$ -algebra  $D$  over a finite CW-complex  $X$  form a group  $E_D^1(X)$  that is the first group of a cohomology theory  $E_D^*(X)$ . In this paper we compute these groups by expressing them in terms of ordinary cohomology and connective  $K$ -theory. To compare the  $C^*$ -algebraic version of  $gl_1(KU)$  with its classical counterpart we also develop a uniqueness result for the unit spectrum of complex periodic topological  $K$ -theory.

## CONTENTS

1. Introduction	1
2. Symmetric ring spectra representing $K$ -theory and its localisations	4
2.1. Strongly self-absorbing $C^*$ -algebras	4
2.2. The symmetric spectra $KU^D$	5
2.3. The unit spectrum of $KU^D$	7
2.4. Connective $K$ -theory	8
3. Units of $K$ -theory and the classification of bundles	9
3.1. Splitting results	10
3.2. Comparing $E_{\mathbb{Z}}^1(X)$ and $bsu_{\otimes}^1(X)$ and proof of Theorem A	15
4. $K$ -theory as a commutative $S$ -algebra and its units	16
4.1. Localizations, connective $K$ -theory, $bsu_{\oplus}$ and $bsu_{\otimes}$	17
5. Uniqueness of $gl_1$ for $K$ -theory spectra	18
5.1. Change of categories.	19
5.2. Obstruction theory.	20
5.3. The $S$ -module $\Sigma^{\infty}CP_+^{\infty}[b^{-1}]$	21
5.4. Equivalences from $\Sigma^{\infty}CP_+^{\infty}[b^{-1}]$ to $KU$ and $G$ .	22
5.5. A product map for $CP_+^{\infty}[b^{-1}]$	23
5.6. Proof of Lemma 5.6	23
References	25

## 1. INTRODUCTION

Continuous fields of  $C^*$ -algebras occur naturally as they correspond to bundles of  $C^*$ -algebras in the sense of topology [9]. Any  $C^*$ -algebra with Hausdorff primitive spectrum  $X$  is a continuous field of simple  $C^*$ -algebras over  $X$ , [25], [22]. More importantly, continuous fields are used as versatile tools in several areas:  $E$ -theory [12], deformations of the tangent groupoid

---

J.E.McC. thanks the Lord for making his work possible.

of manifolds [11], [1], strict deformation quantization [52], [39], the Novikov conjecture and the Baum-Connes conjecture, [37], [30], [60], representation theory and index theory [28], [29].

Particularly well-behaved examples of continuous fields over a compact space  $X$  can be obtained as the continuous sections  $C(X, E)$  of a locally trivial bundle  $E \rightarrow X$  with fibres isomorphic to a fixed  $C^*$ -algebra  $A$ . Such bundles are constructed by gluing together trivial bundles  $U_i \times A$  over an open cover  $(U_i)_{i \in I}$  of the base space  $X$  using 1-cocycles  $\gamma_{ij}: U_i \cap U_j \rightarrow \text{Aut}(A)$ . Two such locally trivial bundles  $E$  and  $F$  over a compact Hausdorff space  $X$  are isomorphic if and only if their section  $C^*$ -algebras  $C(X, E)$  and  $C(X, F)$  are isomorphic via a  $C(X)$ -linear  $*$ -isomorphism. The proof of this property is an elementary exercise in bundle theory [14].

As the description by 1-cocycles indicates, the classification of isomorphism classes of locally trivial bundles with fibre  $A$  can be reduced to the classification of principal  $\text{Aut}(A)$ -bundles over  $X$  up to isomorphism, and hence to the computation of the homotopy classes of maps from  $X$  to the classifying space  $B\text{Aut}(A)$ . Without further structure the homotopy type of the classifying space is typically very difficult to determine and the computation of homotopy sets such as  $[X, B\text{Aut}(A)]$  is out of reach. This changes quite drastically, however, if  $A = D \otimes \mathcal{K}$ , where  $D$  belongs to the class of strongly self-absorbing  $C^*$ -algebras (see Sec. 2.1) and  $\mathcal{K}$  denotes the compact operators on an infinite-dimensional separable Hilbert space. In this case  $\text{Aut}(D \otimes \mathcal{K})$  turns out to be an infinite loop space, which not only implies that  $E_D^1(X) := [X, B\text{Aut}(D \otimes \mathcal{K})]$  is an abelian group, the 1-group of the generalized cohomology theory  $E_D^*(X)$ , but also that it is amenable to methods from stable homotopy theory, [19], [18].

Remarkably, the group law on  $E_D^1(X)$  coming from the infinite loop space structure of  $B\text{Aut}(D \otimes \mathcal{K})$  coincides with the operation induced by the tensor product of  $D \otimes \mathcal{K}$ -bundles. This led to a generalized Dixmier-Douady theory. Let us note that if  $D = \mathbb{C}$ , then  $E_{\mathbb{C}}^1(X) \cong H^3(X, \mathbb{Z})$  is the home of the classic Dixmier-Douady class. Strongly self-absorbing  $C^*$ -algebras [57] are separable unital  $C^*$ -algebras  $D$  defined by a crucial property that they share with the complex numbers  $\mathbb{C}$ . Namely, there exists an isomorphism  $D \rightarrow D \otimes D$  which is unitarily homotopic to the map  $d \mapsto d \otimes 1_D$  [21], [61]. Any strongly self-absorbing  $C^*$ -algebra  $D$  is either stably finite or purely infinite. The latter condition is equivalent to  $D \cong D \otimes \mathcal{O}_\infty$ , where  $\mathcal{O}_\infty$  is the infinite Cuntz algebra. Due to recent progress in classification theory [62] we now have a complete list of all the strongly self-absorbing  $C^*$ -algebras that satisfy the Universal Coefficient Theorem (abbreviated UCT) in KK-theory.

The main goal of this paper is to compute the group  $E_D^1(X)$  for all strongly self-absorbing  $C^*$ -algebras  $D$  in the UCT class. More precisely, we express  $E_D^1(X)$  and its variants using connective  $K$ -theory and ordinary cohomology groups. While this question is interesting in itself and in view of direct connections with higher twisted K-theory [50], we draw additional motivation from the following recent development: a conjecture of Izumi from [33], [34] has been recently proved due to combined work of Meyer [48] and Gabe and Szabó [26]. It asserts that for a countable torsion free amenable group  $G$  and for a Kirchberg algebra  $D$ , there is a bijection between the cocycle conjugacy classes of outer actions of  $G$  on  $D \otimes K$  and the isomorphism classes of principal  $\text{Aut}(D \otimes K)$ -bundles over the classifying space  $BG$ , i.e. the set of homotopy classes  $[BG, B\text{Aut}(D \otimes K)]$ .

It follows that if  $BG$  admits a model as a finite CW-complex and if  $D$  is a strongly self-absorbing Kirchberg algebra that satisfies the UCT, then the set of cocycle conjugacy classes of outer actions of  $G$  on  $D \otimes K$  forms a group with respect to the tensor product operation, and this group is isomorphic to  $E_D^1(BG)$ . Moreover, this group can be computed as explained below.

Let  $k^*(X)$  denote the complex connective  $K$ -theory of the space  $X$ . For a finite based CW-complex  $X$  with skeleta  $X_i$ ,  $\tilde{k}^i(X) \cong \tilde{K}^i(X, X_{i-2})$  and in particular  $k^5(X) \cong K^1(X, X_3)$ , see Proposition 2.2. For a set of primes  $P \neq \emptyset$ , consider the  $C^*$ -algebra  $M_P = \bigotimes_{p \in P} M_p(\mathbb{C})^{\otimes \infty}$  and its  $K$ -theory ring  $\mathbb{Z}_P = \bigotimes_{p \in P} \mathbb{Z}[\frac{1}{p}]$  viewed as a subring of  $\mathbb{Q}$ . Our main result is the following:

**Theorem A.** *Let  $X$  be a finite CW-complex and let  $P$  be a nonempty set of prime numbers. There are isomorphisms*

- (a)  $E_{\mathbb{Z}}^1(X) \cong H^3(X, \mathbb{Z}) \oplus k^5(X)$ .
- (b)  $E_{M_P}^1(X) \cong H^1(X, (\mathbb{Z}_P)_+^{\times}) \oplus H^3(X, \mathbb{Z}_P) \oplus k^5(X, \mathbb{Z}_P)$
- (c)  $E_{\mathcal{O}_{\infty}}^1(X) \cong (H^1(X, \mathbb{Z}/2) \times_{tw} H^3(X, \mathbb{Z})) \oplus k^5(X)$ .
- (d)  $E_{M_P \otimes \mathcal{O}_{\infty}}^1(X) \cong H^1(X, (\mathbb{Z}_P)_+^{\times}) \oplus (H^1(X, \mathbb{Z}/2) \times_{tw} H^3(X, \mathbb{Z}_P)) \oplus k^5(X, \mathbb{Z}_P)$

The (twisted) multiplication on  $H^1(X; \mathbb{Z}/2) \times H^3(X, \mathbb{Z}_P)$  is given by

$$(1) \quad (w, \tau) \cdot (w', \tau') = (w + w', \tau + \tau' + \beta_P(w \cup w'))$$

for  $w, w' \in H^1(X, \mathbb{Z}/2)$  and  $\tau, \tau' \in H^3(X, \mathbb{Z}_P)$ , where  $\beta_P: H^2(X, \mathbb{Z}/2) \rightarrow H^3(X, \mathbb{Z}_P)$  is the composition of the Bockstein homomorphism  $\beta$  with the coefficient map  $H^3(X, \mathbb{Z}) \rightarrow H^3(X, \mathbb{Z}_P)$ . The multiplication in (c) is just like in (1) with  $\beta_P$  replaced by  $\beta: H^2(X, \mathbb{Z}/2) \rightarrow H^3(X, \mathbb{Z})$ .

The isomorphisms above are not natural. The article is structured as follows: In Section 2.1 we recall the definition of strongly self-absorbing  $C^*$ -algebras, discuss their classification and give their  $K$ -theory groups. To each such algebra one can associate a commutative symmetric ring spectrum  $KU^D$  (see [18]). Its definition is recalled in Section 2.2. Just like a commutative ring has a group of invertible elements (or units), the spectrum  $KU^D$  has an associated unit spectrum, whose definition is based on commutative  $\mathcal{I}$ -monoids, see [53]. All of these notions are reviewed in Section 2.3. As we will see in later chapters the groups classifying bundles of stabilised strongly self-absorbing  $C^*$ -algebras can be expressed in terms of connective  $K$ -theory  $ku$ . Therefore we recall the main properties of  $ku$  needed in the rest of the paper in Section 2.4.

In Section 3 we discuss the classification of  $C^*$ -algebra bundles with fibre  $D \otimes \mathcal{K}$  for a strongly self-absorbing  $C^*$ -algebra  $D$ . We review the construction of the cohomology theories  $E_D^*(X)$  and its variants at the beginning of Section 3. As shown in Section 3.1 these theories all split off a low-degree summand that can be expressed via ordinary cohomology groups with a “twisted multiplication”. The complement of this summand, called  $h(X)$  in Section 3, is identified in Section 3.2 with  $bsu_{\otimes}^1(X)$ . Due to the Adams-Priddy result on the uniqueness of  $bsu$ , [4], we can express  $bsu_{\otimes}^1(X)$  in terms of connective  $K$ -theory (which we review in Sect.2.4). The subtle point in this last part is that we have to deal with two constructions of the units of topological  $K$ -theory:  $gl_1(KU^{\mathbb{C}})$  and  $gl_1(KU)$  for a commutative  $S$ -algebra  $KU$  representing  $K$ -theory. We will show in Sections 4 and 5 that both of these give the same cohomology theory.

Since the proof of the uniqueness result in Section 5 requires a lot of heavy machinery from stable homotopy theory, Section 4 outlines the preliminaries that we need: We discuss the definition of commutative  $S$ -algebras and their unit spectra in Section 4.1, which also contains the definition of the commutative  $S$ -algebra  $KU$  that will play a crucial role in Section 5.

Section 5 of this paper is devoted entirely to the proof of the uniqueness result which gives a natural isomorphism  $gl_1(KU^{\mathbb{C}})^*(X) \cong gl_1(KU)^*(X)$ . In Section 5.1 we explain how to move from units of commutative symmetric ring spectra to units of commutative  $S$ -algebras. We use the obstruction theory of Goerss and Hopkins in Section 5.2 to show that uniqueness follows if we have a unital and multiplicative isomorphism between a commutative  $S$ -algebra model  $G$  for  $K$ -theory and  $KU$  in the homotopy category. In the remaining Sections 5.3 to 5.6 we construct

an intermediate spectrum  $\Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}]$ , which has the benefit that maps from it to  $G$  and  $KU$  can be constructed from maps on  $\mathbb{C}P^n$  that are easily obtained from the multiplicative natural transformation.

## 2. SYMMETRIC RING SPECTRA REPRESENTING $K$ -THEORY AND ITS LOCALISATIONS

Topological  $K$ -theory is a multiplicative cohomology theory and can be modelled as a commutative symmetric ring spectrum [36]. In this form it has a natural extension  $KU^D$  that takes a strongly self-absorbing  $C^*$ -algebra  $D$  as input. For any such  $D$  the group  $K_0(D)$  is actually a ring and if  $D$  satisfies the UCT, then  $K_0(D) \subseteq \mathbb{Q}$  as a subring. The spectrum  $KU^D$  represents topological  $K$ -theory with coefficients in  $K_0(D)$ , i.e. a localisation of  $K$ -theory. We recall the definition of strongly self-absorbing  $C^*$ -algebras, the construction of  $KU^D$  and its unit spectrum in the sense of stable homotopy theory in the next sections.

**2.1. Strongly self-absorbing  $C^*$ -algebras.** A  $C^*$ -algebra  $A$  absorbs another  $C^*$ -algebra  $D$  tensorially if  $A \otimes D \cong A$ . As was already observed in [38, 51], tensorial absorption properties are crucial in the classification programme for separable simple nuclear  $C^*$ -algebras. This prompted an analysis of strongly self-absorbing  $C^*$ -algebras [58]. By definition a unital  $C^*$ -algebra  $D$  belongs to this class if there exists an isomorphism  $D \rightarrow D \otimes D$  that is approximately unitarily equivalent to  $d \mapsto d \otimes 1$ .

The two Cuntz algebras  $\mathcal{O}_2$  and  $\mathcal{O}_\infty$  with two, respectively infinitely many generators (see [13]) are strongly self-absorbing. Another prominent example in this class is the Jiang-Su algebra  $\mathcal{Z}$  (see [35]), which can be viewed as the infinite-dimensional stably finite counterpart of  $\mathbb{C}$ , while  $\mathcal{O}_\infty$  is its purely infinite version. We have an isomorphism  $\mathcal{O}_\infty \otimes \mathcal{Z} \cong \mathcal{O}_\infty$ . The unital  $*$ -homomorphisms  $\mathbb{C} \rightarrow \mathcal{Z} \rightarrow \mathcal{O}_\infty$  are  $KK$ -equivalences.  $\mathcal{O}_2$  is  $KK$ -contractible.

All self-absorbing  $C^*$ -algebras that satisfy the UCT, with the exception of  $\mathbb{C}$  and  $\mathcal{O}_2$ , can be obtained as tensor products of either  $\mathcal{Z}$  or  $\mathcal{O}_\infty$  with an infinite UHF-algebra. The construction of these is a  $C^*$ -algebraic version of the localisation at a set of primes: For a prime  $p$ , let  $M_p$  denote the infinite tensor product  $M_p = M_p(\mathbb{C})^{\otimes \infty}$ . For a set of primes  $P$  define  $M_P$  to be

$$M_P = \bigotimes_{p \in P} M_p .$$

If  $P = \emptyset$ , then we set  $M_P = \mathbb{C}$ . There is a dichotomy for strongly self-absorbing  $C^*$ -algebras: they are either stably finite or purely infinite. In the first case the algebra is isomorphic to either  $\mathbb{C}$ ,  $\mathcal{Z}$ , or  $M_P$  for some nonempty set  $P$  of primes, in the second to  $\mathcal{O}_\infty$ ,  $M_P \otimes \mathcal{O}_\infty$  for some nonempty set  $P$  of primes or to  $\mathcal{O}_2$ , [62]. If  $D$  is purely infinite, then  $D \otimes \mathcal{O}_\infty \cong D$ .

If  $D$  is strongly self-absorbing and  $D \neq \mathbb{C}$ , then  $D \cong D \otimes \mathcal{Z}$ . In particular, we have  $M_P \otimes \mathcal{Z} \cong M_P$ . If  $P \subset P'$  is an inclusion of subsets of prime numbers, then  $M_{P'} \otimes M_P \cong M_{P'}$ . In case  $P$  is the set of all primes, then we denote  $M_P$  by  $\mathcal{Q}$ . The  $C^*$ -algebra  $\mathcal{Q}$  is called the universal UHF-algebra. The relationship between the various strongly self-absorbing  $C^*$ -algebras is illustrated in the following diagram:

$$(2) \quad \begin{array}{ccccccc} & & \mathcal{Z} & \longrightarrow & M_P & \longrightarrow & \mathcal{Q} & & \\ & \nearrow & \downarrow & & \downarrow & & \downarrow & \searrow & \\ \mathbb{C} & & \mathcal{O}_\infty & \longrightarrow & \mathcal{O}_\infty \otimes M_P & \longrightarrow & \mathcal{O}_\infty \otimes \mathcal{Q} & \longrightarrow & \mathcal{O}_2 \end{array}$$

An arrow  $D \rightarrow D'$  in this diagram not only indicates a unital embedding, but also the property  $D' \otimes D \cong D'$ .

For strongly self-absorbing  $C^*$ -algebras  $D$  in the UCT class the  $K$ -theory groups can be computed as follows, [58]:  $K_1(D) = 0$  and as mentioned above, the group  $K_0(D) \cong K_0(\mathcal{O}_\infty \otimes D)$  has a natural unital commutative ring structure with multiplication induced by the isomorphism  $D \otimes D \cong D$ . Let  $\mathbb{Z}_p = \mathbb{Z}[\frac{1}{p}]$  denote the localization of  $\mathbb{Z}$  away from  $p$  and  $\mathbb{Z}_{(p)}$  the localization of  $\mathbb{Z}$  at  $p$ . Then we have natural ring isomorphisms:

$$\begin{aligned} K_0(\mathbb{C}) &\cong K_0(\mathcal{Z}) \cong K(\mathcal{O}_\infty) \cong \mathbb{Z} , \\ K_0(M_P) &\cong K_0(M_P \otimes \mathcal{O}_\infty) \cong \bigotimes_{p \in P} \mathbb{Z}_p =: \mathbb{Z}_P , \\ K_0(\mathcal{O}_2) &= 0 . \end{aligned}$$

Mirroring the localisation of integers, we denote the algebra  $M_P$  by  $M_{(p)}$  in case  $P$  is the set of all primes different from  $p$ . Thus:  $K_0(M_p) \cong \mathbb{Z}_p$ ,  $K_0(M_{(p)}) \cong \mathbb{Z}_{(p)}$ , and  $K_0(\mathcal{Q}) \cong \mathbb{Q}$ .

We will denote the invertible elements of the commutative ring  $K_0(D)$  either by  $K_0(D)^\times$  or by  $GL_1(K_0(D))$ . Note that  $K_0(D)$  is an ordered group with positive cone given by the elements represented by classes of projections in  $D \otimes \mathcal{K}$  (as opposed to formal differences). The subgroup of positive elements of  $K_0(D)^\times$  is then denoted by  $K_0(D)_+^\times$ . Since the order structure is trivial if  $D$  is purely infinite, we have  $K_0(D)_+^\times = K_0(D)^\times$  in this case.

**2.2. The symmetric spectra  $KU^D$ .** A  $\mathbb{Z}/2\mathbb{Z}$ -grading on a  $C^*$ -algebra  $A$  is an automorphism  $\gamma \in \text{Aut}(A)$  with  $\gamma^2 = \text{id}_A$ . We call the pair  $(A, \gamma)$  a graded  $C^*$ -algebra and will often suppress the grading in the notation. Any  $\mathbb{Z}/2\mathbb{Z}$ -graded  $C^*$ -algebra  $A$  has a Banach space decomposition  $A \cong A^0 \oplus A^1$  with

$$A^0 = \{a \in A \mid \gamma(a) = a\} \quad \text{and} \quad A^1 = \{a \in A \mid \gamma(a) = -a\}$$

such that the even part  $A^0$  is a  $C^*$ -subalgebra and  $A^i \cdot A^j \subset A^{i+j}$ , where the supscript is taken modulo 2 here. The elements  $a \in A^i$  are said to be homogeneous and have degree  $i$ , which we denote by  $\text{deg}(a) = i$ . If  $\gamma = \text{id}_A$ , then we call  $A$  trivially graded. The (minimal) graded tensor product of two graded  $C^*$ -algebras  $A$  and  $B$  is a completion of the algebraic tensor product  $A \odot B$  with the multiplication and involution defined on homogeneous elements by

$$(a \otimes b) \cdot (a' \otimes b') = (-1)^{\text{deg}(a') \cdot \text{deg}(b)} a a' \otimes b b' \quad \text{and} \quad (a \otimes b)^* = (-1)^{\text{deg}(a) \cdot \text{deg}(b)} a^* \otimes b^*$$

The tensor flip  $A \otimes B \rightarrow B \otimes A$  has to be decorated with the sign  $(-1)^{\text{deg}(a) \cdot \text{deg}(b)}$  as well to be a  $*$ -isomorphism.

Two graded  $C^*$ -algebras will play a central role in the following construction: The Clifford algebra  $\mathcal{C}\ell_n$  is the unital  $C^*$ -algebra with self-adjoint generators  $\{e_1, \dots, e_n\}$  and relations

$$e_i e_j + e_j e_i = \delta_{ij} 1 ,$$

where  $\delta_{ii}$  is the Kronecker delta. The grading  $\gamma$  is defined on generators by  $\gamma(e_i) = -e_i$ , i.e. the elements  $e_i$  are odd. These structures turn  $\mathcal{C}\ell_n$  into a finite-dimensional graded  $C^*$ -algebra.

The other non-trivially graded  $C^*$ -algebra that we will encounter is the graded suspension algebra  $\mathcal{S} = C_0(\mathbb{R})$  equipped with the grading by odd and even functions. This algebra can be equipped with a coassociative and cocommutative comultiplication  $\Delta: \mathcal{S} \rightarrow \mathcal{S} \otimes \mathcal{S}$  that has a counit  $\epsilon: \mathcal{S} \rightarrow \mathbb{C}$  [36, p. 94].

Let  $D$  be a strongly self-absorbing  $C^*$ -algebra, considered as a trivially graded algebra. It was shown in [18] that the sequence of spaces  $(KU_n^D)_{n \in \mathbb{N}_0}$  given by

$$KU_n^D = \text{hom}_{\text{gr}}(\mathcal{S}, (\mathcal{C}l_1 \otimes D \otimes \mathcal{K})^{\otimes n})$$

and equipped with the point-norm topology forms a commutative symmetric ring spectrum representing the cohomology theory  $X \mapsto K_*(C(X) \otimes D)$ . The multiplicative structure on  $KU^D$  is induced by

$$KU_n^D \wedge KU_m^D \rightarrow KU_{n+m}^D \quad , \quad \varphi \wedge \psi \mapsto (\varphi \otimes \psi) \circ \Delta .$$

Bott periodicity gives an element in  $\text{hom}_{\text{gr}}(\mathcal{S}, C_0(\mathbb{R}) \otimes \mathcal{C}l_1)$  and hence by extension with a rank 1-projection  $1 \otimes e \in D \otimes \mathcal{K}$  an element in

$$\text{Map}_*(S^1, KU_1^D) \cong \text{hom}_{\text{gr}}(\mathcal{S}, C_0(\mathbb{R}) \otimes \mathcal{C}l_1 \otimes D \otimes \mathcal{K})$$

(see [18, Sec. 4.1] for details). The unit map  $\eta_n: S^n \rightarrow KU_n^D$  of the ring spectrum is constructed as an  $n$ -fold power of  $\eta_1$  (using the above multiplication). As usual combining the unit maps with the multiplication gives the structure maps

$$(3) \quad S^1 \wedge KU_n^D \rightarrow KU_{n+1}^D$$

Note that the space  $KU_0^D = \text{hom}_{\text{gr}}(\mathcal{S}, \mathbb{C}) \simeq S^0$  contains only two points. One is given by the zero homomorphism, the other one is given by the evaluation at 0, which is the only evaluation that respects the grading. Nevertheless, as we will see in Lem. 2.1 all adjoints of the structure maps in degree  $\geq 1$  are weak homotopy equivalences. Symmetric spectra with this property are called *positive  $\Omega$ -spectra*. In fact, more is true:

**Lemma 2.1.** *Let  $D$  be a strongly self-absorbing  $C^*$ -algebra. The spectrum  $KU^D$  is a positive  $\Omega$ -spectrum that represents  $X \mapsto K_*(C(X) \otimes D)$  as a multiplicative cohomology theory.*

*Proof.* For  $n \geq 1$  the adjoint  $KU_n^D \rightarrow \Omega KU_{n+1}^D$  of (3) is a weak equivalence by [18, Thm. 4.2]. Thus,  $KU^D$  is a positive  $\Omega$ -spectrum. By [59, Thm. 4.7] there are natural isomorphisms

$$[X, KU_n^D] \cong [\mathcal{S}, C(X) \otimes \mathcal{C}l_n \otimes D \otimes \mathcal{K}]_{\text{gr}} \cong KK(\mathbb{C}, C(X) \otimes \mathcal{C}l_n \otimes D) \cong K_n(C(X) \otimes D) ,$$

for  $n \geq 1$ , where  $[\cdot, \cdot]_{\text{gr}}$  denotes the homotopy classes of graded homomorphisms. It remains to be checked that this is compatible with the multiplicative structure. Using Bott periodicity and a suspension argument this can be reduced to the question whether  $KU_2^D \wedge KU_2^D \rightarrow KU_4^D$  implements the multiplication in  $K_0(C(X) \otimes D) \cong K_2(C(X) \otimes D)$ .

Note that  $\mathcal{C}l_2 \cong M_2(\mathbb{C})$  if the right hand side is equipped with the diagonal/off-diagonal grading. Let  $p_1, p_2, q_1, q_2 \in C(X) \otimes D \otimes \mathcal{K}$  be projections and consider

$$\varphi_{p_i, q_i}: \mathcal{S} \rightarrow \mathcal{C}l_2 \otimes C(X) \otimes D \otimes \mathcal{K} \quad , \quad f \mapsto \epsilon(f) \begin{pmatrix} p_i & 0 \\ 0 & q_i \end{pmatrix} .$$

By [59, p. 303] this represents the element  $[p_i] - [q_i] \in K_0(C(X) \otimes D)$  in  $[X, KU_2^D]$ . We have  $\mathcal{C}l_2 \otimes \mathcal{C}l_2 \cong \mathcal{C}l_2 \otimes M_2(\mathbb{C})$  with  $M_2(\mathbb{C})$  trivially graded. Moreover,  $(\epsilon \otimes \epsilon) \circ \Delta = \epsilon$ , since  $\epsilon$  is a counit. Hence, we obtain

$$((\varphi_{p_1, q_1} \otimes \varphi_{p_2, q_2}) \circ \Delta)(f) = \epsilon(f) \begin{pmatrix} p_1 \otimes p_2 \oplus q_1 \otimes q_2 & 0 \\ 0 & p_1 \otimes q_2 \oplus q_1 \otimes p_2 \end{pmatrix} ,$$

which indeed represents the class  $([p_1] - [q_1]) \cdot ([p_2] - [q_2]) \in K_0(C(X) \otimes D)$ .  $\square$



**2.3. The unit spectrum of  $KU^D$ .** Following Schlichtkrull [53], we give a brief outline of how to define the units of a commutative symmetric ring spectrum. This is based on the following model for  $E_\infty$ -spaces: Let  $\mathcal{I}$  be the category with objects the sets  $\mathbf{n} = \{1, \dots, n\}$  for  $n \in \mathbb{N}_0$  (with  $\mathbf{0} = \emptyset$ ) and injective maps as morphisms. This is a symmetric monoidal category, where the tensor product is given by  $\mathbf{n} \sqcup \mathbf{m} = \{1, \dots, n+m\}$  on objects and on morphisms by identifying  $\mathbf{n}$  with  $\{1, \dots, n\} \subset \mathbf{n} \sqcup \mathbf{m}$  and  $\mathbf{m}$  with  $\{n+1, \dots, n+m\} \subset \mathbf{n} \sqcup \mathbf{m}$ . The object  $\mathbf{0}$  is the monoidal unit and the symmetry  $\mathbf{n} + \mathbf{m} \rightarrow \mathbf{m} + \mathbf{n}$  is given by a block permutation.

An  $\mathcal{I}$ -space is a functor  $X: \mathcal{I} \rightarrow \mathcal{T}$  from  $\mathcal{I}$  to the category of based topological spaces. A morphism of  $\mathcal{I}$ -spaces is a natural transformation. The category of  $\mathcal{I}$ -spaces also has a symmetric monoidal structure: For two given  $\mathcal{I}$ -spaces  $X$  and  $Y$ , the tensor product  $X \boxtimes Y$  is defined as the left Kan extension of the  $\mathcal{I}^2$ -space  $X \times Y$  along  $\sqcup: \mathcal{I} \times \mathcal{I} \rightarrow \mathcal{I}$ . This definition implies that a morphism  $X \boxtimes Y \rightarrow Z$  of  $\mathcal{I}$ -spaces  $X, Y, Z$  is the same as a natural transformation

$$X(\mathbf{n}) \times Y(\mathbf{m}) \rightarrow Z(\mathbf{n} \sqcup \mathbf{m})$$

of  $\mathcal{I}^2$ -spaces. A (commutative)  $\mathcal{I}$ -monoid is a (commutative) monoid in the category of  $\mathcal{I}$ -spaces. Note that the category  $\mathcal{I}$  is denoted by  $\mathbb{I}$  in [40] and  $\mathcal{I}$ -monoids are called  $\mathbb{I}$ -FCPs. If  $X$  is a commutative  $\mathcal{I}$ -monoid, then

$$X_{h\mathcal{I}} = \text{hocolim}_{\mathcal{I}} X$$

is an  $E_\infty$ -space [40, Rem. 4.3]. One should think of the homotopy colimit of an  $\mathcal{I}$ -space as the homotopy type modelled by it.

Let  $(R_n)_{n \in \mathbb{N}_0}$  be a commutative symmetric ring spectrum and define  $(\Omega^\bullet R)(\mathbf{n}) = \Omega^n R_n$ . This extends to an  $\mathcal{I}$ -space  $\Omega^\bullet R$ , which should model the infinite loop space underlying the spectrum  $R$ . However, there is a caveat here, since the homotopy groups of  $(\Omega^\bullet R)_{h\mathcal{I}}$  and  $R$  do not necessarily agree (see also [40, Rem. 2.1]). If  $R$  is represented by a positive  $\Omega$ -spectrum, which we will assume for the rest of this paragraph, then this problem does not arise. In analogy to (23) we define the  $\mathcal{I}$ -space of units  $(\Omega^\bullet R)^\times(\mathbf{n})$  of  $R$  to be the subspace of  $\Omega^n R_n$  consisting of those based maps  $f: S^n \rightarrow R_n$  which are stably invertible in the sense that there exists  $g: S^m \rightarrow R_m$  such that

$$\mu \circ (f \wedge g): S^{n+m} \rightarrow R_{n+m}$$

is homotopic to the unit map of the ring spectrum, where  $\mu: R_n \wedge R_m \rightarrow R_{n+m}$  denotes its multiplication. The smash product  $(f, g) \mapsto \mu \circ (f \wedge g)$  gives  $(\Omega^\bullet R)^\times$  the structure of a commutative  $\mathcal{I}$ -monoid and

$$(\Omega^\bullet R)_{h\mathcal{I}}^\times \simeq GL_1(R) .$$

Thus, the space underlying  $(\Omega^\bullet R)^\times$  is  $GL_1(R)$ , but the commutative  $\mathcal{I}$ -monoid reveals an  $E_\infty$ -structure on it. To obtain the spectrum  $gl_1(R)$  from this, we can turn  $(\Omega^\bullet R)^\times$  into a  $\Gamma$ -space and apply an infinite loop space machine to it turning it into a weak  $\Omega$ -spectrum. For details we refer the reader to [53] or to [40, Construction 12.1 and Def. 12.5].

Applying this functor to  $KU^C$  gives a first way of defining the unit spectrum of topological  $K$ -theory and which we denote by  $gl_1(KU^C)$ . We will define the unit spectrum of topological  $K$ -theory in a second second way in Section 4 and denote the corresponding object by  $gl_1(KU)$ . The two spectra  $gl_1(KU^C)$  and  $gl_1(KU)$  will be compared in Section 5.

We finish this part by listing some comparison results among the spectra  $gl_1(KU^D)$ . Recall that a *level equivalence* is a map of symmetric spectra which induces a weak equivalence of the  $n$ -th spaces of the spectra for all  $n \geq 0$  ([43, Definition 6.1(i)]). A level equivalence is a stable equivalence ([43, bottom of page 466]). The converse is not true in general but it is true if the spectra are  $\Omega$ -spectra ([43, Lemma 8.11]).

The spectra  $KU^D$  are positive  $\Omega$ -spectra by Lem. 2.1 and therefore fibrant objects in the model category of commutative symmetric ring spectra by [43, Thm. 14.2]. By [40, Lem. 13.5] the functor  $gl_1$  preserves weak equivalences between fibrant objects. The  $*$ -homomorphisms  $\mathbb{C} \rightarrow \mathcal{Z}$  and  $\mathbb{C} \rightarrow \mathcal{O}_\infty$  induce level and therefore stable equivalences of commutative symmetric ring spectra  $KU^{\mathbb{C}} \rightarrow KU^{\mathcal{Z}}$  and  $KU^{\mathbb{C}} \rightarrow KU^{\mathcal{O}_\infty}$ . Given any set of primes  $P$  the unital  $*$ -homomorphism  $M_P \rightarrow M_P \otimes \mathcal{O}_\infty$  gives a stable equivalence  $KU^{M_P} \rightarrow KU^{M_P \otimes \mathcal{O}_\infty}$ . From these we therefore obtain equivalences of weak  $\Omega$ -spectra

$$(4) \quad gl_1(KU^{\mathbb{C}}) \simeq gl_1(KU^{\mathcal{Z}}) \simeq gl_1(KU^{\mathcal{O}_\infty}) ,$$

$$(5) \quad gl_1(KU^{M_P}) \simeq gl_1(KU^{M_P \otimes \mathcal{O}_\infty}) .$$

**2.4. Connective K-theory.** Let  $ku$  be a spectrum representing connective  $K$ -theory. It is the connective cover i.e. the  $(-1)$ -connected cover of the spectrum  $KU$  of complex periodic  $K$ -theory. There is a map  $ku \rightarrow KU$  of spectra which induces isomorphisms on homotopy groups in non-negative degrees while  $\pi_i(ku) = 0$  for  $i < 0$ . Thus, the homotopy types of the first few spaces forming the spectrum  $ku$  are:

$$BU \times \mathbb{Z}, U, BU, SU, BSU, BBSU \dots$$

The spectrum  $bsu_\oplus$  is the 3-connected cover of  $ku$  (or of  $KU$ ) (see [4]). Hence, we can identify  $bsu_\oplus$  with  $\Sigma^4 ku$ . Let  $k^*$  be the cohomology theory associated to  $ku$ . By our observations there is a natural isomorphism

$$(6) \quad bsu_\oplus^n(X) \cong k^{n+4}(X), \text{ for all } n \geq 0.$$

For our purposes, it is useful to address the question of computing  $bsu_\oplus^1(X) \cong k^5(X)$ . Let  $H\mathbb{Z}$  denote the usual Eilenberg-Mac Lane spectrum representing ordinary cohomology. The Bott operation  $b$  induces a cofiber sequence of spectra

$$\Sigma^2 ku \xrightarrow{b} ku \longrightarrow H\mathbb{Z}$$

and hence a long exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & k^3(X) & \xrightarrow{b} & k^1(X) & \longleftarrow & H^1(X, \mathbb{Z}) \\ & & & & & \swarrow & \\ 0 & \longrightarrow & k^4(X) & \xleftarrow{b} & k^2(X) & \xrightarrow{c_1} & H^2(X, \mathbb{Z}) \\ & & & & & \swarrow & \\ 0 & \longrightarrow & k^5(X) & \xleftarrow{b} & k^3(X) & \longrightarrow & H^3(X, \mathbb{Z}) \\ & & & & & \swarrow & \\ & & & & & \swarrow & \\ & & & & k^6(X) & \xleftarrow{b} & k^4(X) \longrightarrow H^4(X, \mathbb{Z}) \end{array}$$

We see that  $k^1(X) \cong K^1(X)$  and  $k^3(X) \cong \ker(K^1(X) \rightarrow H^1(X, \mathbb{Z}))$  so that

$$(7) \quad k^5(X) \cong \ker(k^3(X) \rightarrow H^3(X, \mathbb{Z})) .$$

The maps  $k^i(X) \rightarrow H^i(X, \mathbb{Z})$  are induced by the deloopings of determinant map  $\det: U \rightarrow U(1)$ . The map  $c_1$  identifies with the first Chern class and is surjective.

For a different description of  $k^5(X)$ , one verifies that if  $X$  is a finite CW-complex with  $i$ -skeleta  $X_i$ ,  $i \geq 0$ , then  $k^3(X) \cong K^1(X, X_1)$ ,  $k^5(X) \cong K^1(X, X_3)$  is isomorphic to the kernel of



the restriction map  $K^1(X, X_1) \rightarrow K^1(X_3, X_1)$ . Indeed, for the benefit of the reader we include the following proposition known to the experts:

**Proposition 2.2.** *If  $X$  is a finite CW-complex with base point, the pair  $X_{i-2} \subset X$  induces an isomorphism of reduced theories  $\tilde{k}^i(X) \cong \tilde{k}^i(X, X_{i-2}) \cong \tilde{K}^i(X, X_{i-2})$ .*

*Proof.* The long exact sequence

$$\longrightarrow \tilde{k}^{i-1}(X_{i-2}) \longrightarrow \tilde{k}^i(X, X_{i-2}) \longrightarrow \tilde{k}^i(X) \longrightarrow \tilde{k}^i(X_{i-2}) \longrightarrow$$

gives the isomorphism  $\tilde{k}^i(X) \cong \tilde{k}^i(X, X_{i-2})$  since  $\tilde{k}^{i-1}(X_{i-2}) = \tilde{k}^i(X_{i-2}) = 0$  as the space  $bu(r)$  is  $(r-1)$ -connected by construction and hence  $\tilde{k}^r(Y) = [Y, bu(r)] = 0$  if  $\dim(Y) \leq r-1$ . Next, the exact sequence

$$\tilde{H}^{j-1}(Y, \mathbb{Z}) \longrightarrow \tilde{k}^{j+2}(Y) \longrightarrow \tilde{k}^j(Y) \longrightarrow \tilde{H}^j(Y, \mathbb{Z})$$

shows that  $\tilde{k}^{j+2}(Y) \cong \tilde{k}^j(Y)$  if  $Y$  is  $j$ -connected and hence

$$\tilde{k}^{j+2}(Y) \cong \tilde{k}^j(Y) \cong \tilde{k}^{j-2}(Y) \cong \dots \cong \tilde{K}^{j-2m}(Y) \cong \tilde{K}^j(Y)$$

for  $2m \geq j$ . Applying this isomorphism for  $Y = X/X_{i-2}$  (which is  $(i-2)$ -connected) one obtains that  $\tilde{k}^i(X, X_{i-2}) \cong \tilde{K}^i(X, X_{i-2})$ .  $\square$

We refer the reader to [55], [16] and [15] for other applications of connective K-theory in operator algebras.

### 3. UNITS OF $K$ -THEORY AND THE CLASSIFICATION OF BUNDLES

Let  $D$  be a stably finite strongly self-absorbing  $C^*$ -algebra satisfying the UCT. From (2) we know that  $D \cong M_P$  for some set  $P$  (possibly empty) of prime numbers. It was shown in [19, 18] that the isomorphism classes of locally trivial  $C^*$ -algebra bundles with fibre  $D \otimes \mathcal{K}$  form a group with respect to the fibrewise tensor product. Each of these groups is part of a cohomology theory closely related to the one represented by the spectrum  $gl_1(KU^D)$ . The reason for the existence of these theories is an infinite loop space structure on the spaces

$$\text{Aut}_0(D \otimes \mathcal{K}), \text{Aut}(D \otimes \mathcal{K}), \text{Aut}(D \otimes \mathcal{O}_\infty \otimes \mathcal{K}) \quad \text{and} \quad \text{Aut}_{\text{gr}}(\mathbb{C}\ell_1 \otimes D \otimes \mathcal{K}),$$

where  $\text{Aut}_0(\cdot)$  denotes the identity component of the automorphism group and  $\text{Aut}_{\text{gr}}(\cdot)$  are the automorphisms that preserve the grading. In fact, each of these spaces is an  $E_\infty$ -space that can be modelled by a commutative  $\mathcal{I}$ -monoid naturally associated to  $D$ . These  $\mathcal{I}$ -monoids and our notation for the associated cohomology theory are listed in Table 1.

commutative $\mathcal{I}$ -monoid		cohomology theory
$\bar{G}_D(\mathbf{n})$	$= \text{Aut}_0((D \otimes \mathcal{K})^{\otimes n})$	$\bar{E}_D^*(X)$
$G_D(\mathbf{n})$	$= \text{Aut}((D \otimes \mathcal{K})^{\otimes n})$	$E_D^*(X)$
$G_{D \otimes \mathcal{O}_\infty}(\mathbf{n})$	$= \text{Aut}((D \otimes \mathcal{O}_\infty \otimes \mathcal{K})^{\otimes n})$	$E_{D \otimes \mathcal{O}_\infty}^*(X)$
$G_D^{\text{gr}}(\mathbf{n})$	$= \text{Aut}_{\text{gr}}((\mathbb{C}\ell_1 \otimes D \otimes \mathcal{K})^{\otimes n})$	$\hat{E}_D^*(X)$

TABLE 1. Cohomology theories associated to the automorphism groups

We refer the reader to [18, Sec. 4.2] and to [20] for further details about their construction. Note that each of these  $\mathcal{I}$ -monoids takes values in topological groups, and if  $G$  denotes one

of them, then  $\mathbf{n} \rightarrow BG(\mathbf{n})$  (defined by taking classifying spaces levelwise) is a commutative  $\mathcal{T}$ -monoid as well. It was shown in [18, Thm. 3.6] that the spectrum obtained from  $BG$  is in fact the shifted version of the spectrum associated to  $G$  itself. This is not obvious, since the classifying space delooping could a priori be different from the one produced by the infinite loop space machine. As a consequence we have  $[X, B\text{Aut}(D \otimes \mathcal{K})] \cong E_D^1(X)$  and similarly for the other variations listed in the table above.

We will need the following result contained in Theorem 5.4 from [20]. Recall that the unit spectrum of  $KU^D$ , denoted by  $gl_1(KU^D)$  was discussed in subsec. 2.3.

**Theorem 3.1.** *Let  $D$  be a stably finite strongly self-absorbing  $C^*$ -algebra that satisfies the UCT. There is a natural action of  $G_D^{gr}$  on the ring spectrum  $KU^D$  which induces a map*

$$(8) \quad \Gamma(G_D^{gr}) \rightarrow \Gamma((\Omega^\bullet KU^D)^\times)$$

of the associated  $\Gamma$ -spaces which in its turn induces an equivalence of the underlying connective spectra in the stable homotopy category. In particular we obtain a natural isomorphism

$$(9) \quad \hat{E}_D^*(X) \cong gl_1(KU^D)^*(X)$$

for  $X$  a finite CW-complex.

**3.1. Splitting results.** The main result in this section is Prop. 3.9, in which we show that each of the cohomology theories defined above splits off a low-degree ordinary cohomology group. It suffices to treat the case where  $D$  is stably finite, because a key result of [20] establishes the isomorphism

$$E_{D \otimes \mathcal{O}_\infty}^1(X) \cong \hat{E}_D^1(X).$$

We have revisited in [20] the following result from [23], [49].

**Proposition 3.2.** *For any finite CW-complex  $X$ ,  $\hat{E}_\mathbb{C}^1(X) \cong H^1(X, \mathbb{Z}/2) \times_{tw} H^3(X, \mathbb{Z})$  with group structure:*

$$(w, \tau) \cdot (w', \tau') = (w + w', \tau + \tau' + \beta(w \cup w'))$$

for  $w, w' \in H^1(X, \mathbb{Z}/2)$  and  $\tau, \tau' \in H^3(X, \mathbb{Z})$ , where  $\beta: H^2(X, \mathbb{Z}/2) \rightarrow H^3(X, \mathbb{Z})$  is the Bockstein homomorphism.

Let us recall the following two theorems from [20]:

**Theorem 3.3.** *Let  $X$  be a finite CW-complex and let  $D$  be a stable finite strongly self-absorbing  $C^*$ -algebra satisfying the UCT. The groups  $E_D^1(X)$  and  $\hat{E}_D^1(X)$  fit into a short exact sequence*

$$0 \longrightarrow E_D^1(X) \longrightarrow \hat{E}_D^1(X) \xrightarrow{\delta_0} H^1(X, \mathbb{Z}/2) \longrightarrow 0.$$

If  $L$  is a real line vector bundle on  $X$  with associated Clifford bundle  $\mathcal{C}l_L$ , then  $\delta_0(\mathcal{C}l_L \otimes D \otimes \mathcal{K}) = w_1(L)$ , where  $w_1(L)$  is the first Stiefel-Whitney class of  $L$ .

**Theorem 3.4.** *Let  $X$  be a finite CW-complex and let  $D$  be a stably finite strongly self-absorbing  $C^*$ -algebra satisfying the UCT. Then there is an isomorphism of groups*

$$\hat{E}_D^1(X) \cong H^1(X; \mathbb{Z}/2) \times_{tw} E_D^1(X)$$

with multiplication on the direct product  $H^1(X; \mathbb{Z}/2) \times E_D^1(X)$  given by

$$(w, \tau) \cdot (w', \tau') = (w + w', \tau + \tau' + j_P \circ \beta(w \cup w'))$$

for  $w, w' \in H^1(X, \mathbb{Z}/2)$  and  $\tau, \tau' \in E_D^1(X)$ , where  $j_P: E_\mathbb{C}^1(X) \rightarrow E_D^1(X)$  is the map induced by the unital  $*$ -homomorphism  $\mathbb{C} \rightarrow D$  and we identify  $E_\mathbb{C}^1(X) \cong H^3(X, \mathbb{Z})$ .

**Remark 3.5.** Just like in [20], we use here the following basic fact, [10, p.93]. Suppose that

$$0 \longrightarrow A \xrightarrow{i} E \xrightarrow{\pi} G \longrightarrow 0$$

is an extension of abelian groups and  $\sigma : G \rightarrow E$  is a map such that  $\sigma(0) = 0$  and  $\pi \circ \sigma = \text{id}_E$ . Let  $c : G \times G \rightarrow A$  be the normalized 2-cocycle defined by  $i(c(g, h)) = \sigma(gh)\sigma(h)^{-1}\sigma^{-1}(g)$ ,  $g, h \in G$ . Then the group  $E$  is isomorphic to  $G \times A$  endowed with the group law:

$$(g, a)(g', a') = (g + g', a + a' + c(g, g')).$$

The coefficients of the cohomology theory  $E_D^*(X)$  are given by the homotopy groups of  $\text{Aut}(D \otimes \mathcal{K})$  computed in [19, Thm.2.18]:

$$(10) \quad E_D^{-i}(\ast) \cong \pi_i(\text{Aut}(D \otimes \mathcal{K})) = \begin{cases} K_0(D)_+^\times, & \text{if } i = 0 \\ K_i(D), & \text{if } i \geq 1. \end{cases}$$

If  $D$  satisfies the UCT, then  $K_i(D) = 0$  if  $i$  is odd and in particular if  $P \neq \emptyset$  is a set of prime numbers, then

$$(11) \quad E_{M_P}^{-i}(\ast) = \begin{cases} (\mathbb{Z}_P)_+^\times, & \text{if } i = 0 \\ \mathbb{Z}_P, & \text{if } i = 2k > 0, \\ 0, & \text{if } i = 2k + 1. \end{cases}$$

Let  $P$  be a set of prime numbers and let  $\mathcal{A} \rightarrow X$  be a locally trivial  $C^*$ -algebra bundle with fibre  $M_P \otimes \mathcal{K}$ . Let  $(U_i)_{i \in I}$  be a trivialising open cover of  $X$  for  $\mathcal{A}$  and denote the transition maps by  $\varphi_{ij} : U_i \cap U_j \rightarrow \text{Aut}(M_P \otimes \mathcal{K})$ . If we apply the  $K_0$ -functor to  $\varphi_{ij}$  we obtain a 1-cocycle

$$K_0(\varphi_{ij}) : U_i \cap U_j \rightarrow \text{Aut}(K_0(M_P \otimes \mathcal{K})) \cong GL_1(\mathbb{Z}_P),$$

which factors through the group of positive invertible elements  $(\mathbb{Z}_P)_+^\times \subset GL_1(\mathbb{Z}_P)$ . The cocycle only depends on the isomorphism class of  $\mathcal{A}$  and is compatible with tensor products, which gives us a group homomorphism (see also [17, Prop. 2.3])

$$\delta_0 : E_{M_P}^1(X) \rightarrow H^1(X, (\mathbb{Z}_P)_+^\times).$$

**Proposition 3.6.** *Let  $X$  be a finite CW-complex and let  $P \neq \emptyset$  be a set of prime numbers. The group  $E_{M_P}^1(X)$  fits into a short exact sequence*

$$0 \longrightarrow \bar{E}_{M_P}^1(X) \longrightarrow E_{M_P}^1(X) \xrightarrow{\delta_0} H^1(X, (\mathbb{Z}_P)_+^\times) \longrightarrow 0$$

and this sequence splits (not naturally). In particular, there is a non-natural isomorphism

$$E_{M_P}^1(X) \cong H^1(X, (\mathbb{Z}_P)_+^\times) \oplus \bar{E}_{M_P}^1(X).$$

*Proof.* By (11),  $\pi_0(\text{Aut}(M_P \otimes \mathcal{K})) \cong K_0(M_P)_+^\times \cong (\mathbb{Z}_P)_+^\times$ . By [18, Cor.2.19], there is an exact sequence

$$1 \rightarrow \text{Aut}_0(M_P \otimes \mathcal{K}) \rightarrow \text{Aut}(M_P \otimes \mathcal{K}) \rightarrow K_0(M_P)_+^\times \rightarrow 1.$$

We will first show that this sequence splits. We will do so by constructing a homomorphism  $\gamma : (\mathbb{Z}_P)_+^\times \rightarrow \text{Aut}(M_P \otimes \mathcal{K})$  that lifts  $\text{Aut}(M_P \otimes \mathcal{K}) \rightarrow \pi_0(\text{Aut}(M_P \otimes \mathcal{K}))$ . Note that

$$(12) \quad (\mathbb{Z}_P)_+^\times \cong \bigoplus_{p \in P} \mathbb{Z}$$

by the prime factor decomposition. Thus, we need to find for each  $p \in P$  an automorphism  $\alpha_p \in \text{Aut}(M_P \otimes \mathcal{K})$  such that all of them commute and  $[\alpha_p(1 \otimes e)] \in K_0(M_P)$  corresponds to  $p$  under the isomorphism  $K_0(M_P) \cong \mathbb{Z}_P$  induced by the trace. Fix  $p$ . We define  $\bar{\alpha}_p$  as follows

$$M_p \otimes \mathcal{K} \xrightarrow{\varphi_1 \otimes \text{id}_{\mathcal{K}}} M_p \otimes M_p(\mathbb{C}) \otimes \mathcal{K} \xrightarrow{\text{id}_{M_p} \otimes \varphi_2} M_p \otimes \mathcal{K}$$

where  $\varphi_1: M_p \rightarrow M_p \otimes M_p(\mathbb{C})$  and  $\varphi_2: M_p(\mathbb{C}) \otimes \mathcal{K} \rightarrow \mathcal{K}$  are isomorphisms. If  $\tau$  is the trace on the finite rank projections of  $M_p \otimes \mathcal{K}$  with  $\tau(1 \otimes e) = 1$ , then  $\tau(\bar{\alpha}_p(1 \otimes e)) = p$ . We may view  $M_P \otimes \mathcal{K}$  as the tensor product over all  $M_p \otimes \mathcal{K}$  for all  $p \in P$ . In case  $P$  is an infinite set we choose

$$(M_{p_1} \otimes \mathcal{K}) \otimes \cdots \otimes (M_{p_i} \otimes \mathcal{K}) \rightarrow (M_{p_1} \otimes \mathcal{K}) \otimes \cdots \otimes (M_{p_i} \otimes \mathcal{K}) \otimes (M_{p_{i+1}} \otimes \mathcal{K})$$

to be given by  $a \mapsto a \otimes (1 \otimes e)$ . The automorphism  $\alpha_p \in \text{Aut}(M_P \otimes \mathcal{K})$  is defined to act via  $\bar{\alpha}_p$  on the appropriate tensor factor and the identity on the rest. These clearly commute.

Set  $D = M_P$  and recall that  $\bar{G}_D(\mathbf{n}) = \text{Aut}_0((D \otimes \mathcal{K})^{\otimes n})$ .

There are maps of  $\mathcal{I}$ -commutative monoids [20, Lem.6.2]:

$$(13) \quad \bar{G}_D(\mathbf{n}) \rightarrow G_D(\mathbf{n}) \rightarrow K_0(D)_+^{\times}$$

and a group homomorphism

$$K_0(D)_+^{\times} \times \text{Aut}_0(D \otimes \mathcal{K}) \rightarrow \text{Aut}((D \otimes \mathcal{K}) \otimes (D \otimes \mathcal{K})) \cong \text{Aut}(D \otimes \mathcal{K}),$$

given by  $(x, \alpha) \mapsto \gamma(x) \otimes \alpha$ . From our previous discussion, this is a homotopy equivalence. We obtain a homotopy equivalence

$$B(K_0(D)_+^{\times}) \times B\text{Aut}_0(D \otimes \mathcal{K}) \rightarrow B\text{Aut}(D \otimes \mathcal{K}).$$

In conjunction with (13), this gives the exact sequence from the statement (and a split as a sequence of pointed sets). To see that the exact sequence of groups splits, one observes that if  $X$  is a finite CW-complex, then  $H^1(X, (Z_P)_+^{\times})$  is a free group since it is isomorphic to  $\text{Hom}(H_1(X, \mathbb{Z}), (Z_P)_+^{\times}) \cong ((Z_P)_+^{\times})^r$  where  $r$  is the rank of  $H_1(X, \mathbb{Z})$ .  $\square$

**Proposition 3.7.** *Let  $P$  be a nonempty set of primes and let  $X$  be a finite CW-complex. The natural maps  $\bar{E}_{M_P}^*(X) \rightarrow \bar{E}_{M_P}^*(X) \otimes \mathbb{Z}_P$  and  $\bar{E}_{\mathbb{Z}}^*(X) \otimes \mathbb{Z}_P \rightarrow \bar{E}_{M_P}^*(X) \otimes \mathbb{Z}_P$  are isomorphisms of groups. It follows that  $\bar{E}_{M_P}^*(X) \cong \bar{E}_{\mathbb{Z}}^*(X) \otimes \mathbb{Z}_P$ .*

*Proof.* Let  $D$  be a strongly self-absorbing  $C^*$ -algebra satisfying the UCT. Since  $\mathbb{Z}_P$  is flat, it follows that  $X \mapsto \bar{E}_D^*(X) \otimes \mathbb{Z}_P$  still satisfies all axioms of a generalized cohomology theory on finite CW-complexes. We have natural transformations of cohomology theories:

$$\bar{E}_{M_P}^*(X) \rightarrow \bar{E}_{M_P}^*(X) \otimes \mathbb{Z}_P, \quad \bar{E}_{\mathbb{Z}}^*(X) \otimes \mathbb{Z}_P \rightarrow \bar{E}_{M_P}^*(X) \otimes \mathbb{Z}_P.$$

These transformations induce isomorphisms of groups on coefficients by (11) and therefore they induce isomorphisms of cohomology theories.  $\square$

Let us recall from [19] that the natural map  $D \rightarrow D \otimes \mathcal{O}_{\infty}$  induces an isomorphism

$$(14) \quad \bar{E}_D^*(X) \xrightarrow{\cong} \bar{E}_{D \otimes \mathcal{O}_{\infty}}^*(X)$$

since one checks that it induces an isomorphism on coefficients in view of (10).

**Proposition 3.8.** *Let  $X$  be a finite CW complex. The canonical maps  $\hat{\rho}: \hat{E}_{\mathbb{C}}^1(X) \rightarrow \hat{E}_{\mathbb{Z}}^1(X)$  and  $\rho: E_{\mathbb{C}}^1(X) \rightarrow E_{\mathbb{Z}}^1(X)$  induced by the unital  $*$ -homomorphism  $\mathbb{C} \rightarrow \mathbb{Z}$  split naturally. Consequently we have natural splittings*

$$(15) \quad \hat{E}_{\mathbb{Z}}^1(X) \cong \hat{E}_{\mathbb{C}}^1(X) \oplus h(X) \quad \text{and} \quad E_{\mathbb{Z}}^1(X) \cong E_{\mathbb{C}}^1(X) \oplus h(X).$$

*Proof.* By the CW-approximation theorem, there are the natural diagrams

$$(16) \quad \begin{array}{ccc} \hat{E}_{\mathbb{C}}^1(X) & \xrightarrow{\hat{\rho}} & \hat{E}_{\mathbb{Z}}^1(X) & & E_{\mathbb{C}}^1(X) & \xrightarrow{\rho} & E_{\mathbb{Z}}^1(X) \\ \downarrow \hat{r}_{\mathbb{C}} & & \downarrow \hat{r}_{\mathbb{Z}} & & \downarrow r_{\mathbb{C}} & & \downarrow r_{\mathbb{Z}} \\ \hat{E}_{\mathbb{C}}^1(X_4) & \xrightarrow{\hat{\rho}_4} & \hat{E}_{\mathbb{Z}}^1(X_4) & & E_{\mathbb{C}}^1(X_4) & \xrightarrow{\rho_4} & E_{\mathbb{Z}}^1(X_4) \end{array}$$

induced by the inclusion of skeleta  $X_4 \hookrightarrow X$ . We will verify that the maps  $\hat{\rho}_4, \hat{r}_{\mathbb{C}}, \rho_4$  and  $r_{\mathbb{C}}$  are bijections. This will clearly imply (15) with

$$h(X) := \ker \hat{r}_{\mathbb{Z}} \cong \ker r_{\mathbb{Z}}.$$

The isomorphism  $\ker \hat{r}_{\mathbb{Z}} \cong \ker r_{\mathbb{Z}}$  follows from the naturality of (3.3).

Consider the commutative diagram from the proof of [20, Thm. 6.7]:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & E_{\mathbb{Z}}^1(X) & \longrightarrow & \hat{E}_{\mathbb{Z}}^1(X) & \longrightarrow & H^1(X; \mathbb{Z}/2) & \longrightarrow & 0 \\ & & \rho \uparrow & & \hat{\rho} \uparrow & & \parallel & & \\ 0 & \longrightarrow & E_{\mathbb{C}}^1(X) & \longrightarrow & \hat{E}_{\mathbb{C}}^1(X) & \longrightarrow & H^1(X; \mathbb{Z}/2) & \longrightarrow & 0 \\ & & \parallel & & \parallel & & \parallel & & \\ 0 & \longrightarrow & H^3(X; \mathbb{Z}) & \longrightarrow & H^3(X; \mathbb{Z}) \times_{tw} H^1(X; \mathbb{Z}/2) & \longrightarrow & H^1(X; \mathbb{Z}/2) & \longrightarrow & 0 \end{array}$$

If we show that  $\rho_4 : E_{\mathbb{C}}^1(X_4) \rightarrow E_{\mathbb{Z}}^1(X_4)$  is bijective so is  $\hat{\rho}_4$ . The map  $\rho$  is induced by the map  $B\text{Aut}(\mathcal{K}) \rightarrow B\text{Aut}(\mathcal{Z} \otimes \mathcal{K})$  which is 4-connected by the computations of [17], see (10). By Whitehead's theorem this shows that  $\rho_4$  is surjective and that  $\rho_3 : E_{\mathbb{C}}^1(X_3) \rightarrow E_{\mathbb{Z}}^1(X_3)$  is bijective. The restriction map  $r'_{\mathbb{C}} : E_{\mathbb{C}}^1(X_4) \rightarrow E_{\mathbb{C}}^1(X_3)$  in the commutative diagram below

$$(17) \quad \begin{array}{ccc} E_{\mathbb{C}}^1(X_4) & \xrightarrow{\rho_4} & E_{\mathbb{Z}}^1(X_4) \\ \downarrow r'_{\mathbb{C}} & & \downarrow r'_{\mathbb{Z}} \\ E_{\mathbb{C}}^1(X_3) & \xrightarrow[\cong]{\rho_3} & E_{\mathbb{Z}}^1(X_3) \end{array}$$

is injective since it identifies with the map  $H^3(X_4, \mathbb{Z}) \rightarrow H^3(X_3, \mathbb{Z})$  which is injective since  $H^3(X_4/X_3, \mathbb{Z}) = 0$ . It follows that  $\rho_4$  is also injective. Next we show that the map  $\hat{r}_{\mathbb{C}}$  is bijective. Using the naturality of the exact sequence from Theorem 3.3, since the restriction map  $H^1(X; \mathbb{Z}/2) \rightarrow H^1(X_4; \mathbb{Z}/2)$  is bijective, it suffices to show that the restriction map  $r_{\mathbb{C}} : E_{\mathbb{C}}^1(X) \rightarrow E_{\mathbb{C}}^1(X_4)$  is bijective. This follows from the exact sequence

$$H^3(X/X_4, \mathbb{Z}) \rightarrow H^3(X, \mathbb{Z}) \rightarrow H^3(X_4, \mathbb{Z}) \rightarrow H^4(X/X_4, \mathbb{Z}),$$

since  $X/X_4$  is 4-connected. □

**Proposition 3.9.** *Let  $P \neq \emptyset$  be a set of prime numbers and let  $X$  be a finite CW complex. Then there are isomorphisms of groups*

- (a)  $E_{\mathbb{Z}}^1(X) \cong \bar{E}_{\mathbb{Z}}^1(X) \cong H^3(X, \mathbb{Z}) \oplus h(X)$ .
- (b)  $\hat{E}_{\mathbb{Z}}^1(X) \cong (H^1(X, \mathbb{Z}/2) \times_{tw} H^3(X, \mathbb{Z})) \oplus h(X)$ .
- (c)  $E_{M_P}^1(X) \cong H^1(X, (\mathbb{Z}_P)_+^{\times}) \oplus \bar{E}_{M_P}^1(X) \cong H^1(X, (\mathbb{Z}_P)_+^{\times}) \oplus H^3(X, \mathbb{Z}) \otimes \mathbb{Z}_P \oplus h(X) \otimes \mathbb{Z}_P$

- (d)  $\hat{E}_{M_P}^1(X) \cong H^1(X, (Z_P)_+^\times) \oplus (H^1(X; \mathbb{Z}/2) \times_{tw} H^3(X, \mathbb{Z}_P)) \oplus h(X) \otimes \mathbb{Z}_P$  with multiplication on  $H^1(X; \mathbb{Z}/2) \times H^3(X, \mathbb{Z}_P)$

$$(w, \tau) \cdot (w', \tau') = (w + w', \tau + \tau' + \beta_P(w \cup w'))$$

for  $w, w' \in H^1(X, \mathbb{Z}/2)$  and  $\tau, \tau' \in H^3(X, \mathbb{Z}_P)$ , where  $\beta_P: H^2(X, \mathbb{Z}/2) \rightarrow H^3(X, \mathbb{Z}_P)$  is the composition of the Bockstein homomorphism with the coefficient map  $H^3(X, \mathbb{Z}) \rightarrow H^3(X, \mathbb{Z}_P)$ .

*Proof.* The isomorphism (a) follows from (15). Recall that  $E_{\mathbb{Z}}^1(X) \cong \bar{E}_{\mathbb{Z}}^1(X)$  since  $\text{Aut}(\mathcal{Z} \otimes \mathcal{K})$  is path connected. The isomorphism (c) follows from (a), Proposition 3.6 and Proposition 3.7.

It remains to deal with (b) and (d). Along the way we shall review the proof of Theorem 3.4. If  $D' \mapsto D$  is a unital  $*$ -monomorphism of strongly self-absorbing  $C^*$ -algebras, by [20, Lem. 6.3] there is a commutative diagram of commutative  $\mathcal{I}$ -monoids:

$$\begin{array}{ccccc} G_D(\mathbf{n}) & \longrightarrow & G_D^{\text{gr}}(\mathbf{n}) & \longrightarrow & \mathbb{Z}/2 \\ \uparrow & & \uparrow & & \parallel \\ G_{D'}(\mathbf{n}) & \longrightarrow & G_{D'}^{\text{gr}}(\mathbf{n}) & \longrightarrow & \mathbb{Z}/2 \end{array}$$

which induces a commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & E_{M_P}^1(X) & \longrightarrow & \hat{E}_{M_P}^1(X) & \longrightarrow & H^1(X; \mathbb{Z}/2) & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \parallel & & \\ 0 & \longrightarrow & E_{\mathbb{Z}}^1(X) & \longrightarrow & \hat{E}_{\mathbb{Z}}^1(X) & \longrightarrow & H^1(X; \mathbb{Z}/2) & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \parallel & & \\ 0 & \longrightarrow & E_{\mathbb{C}}^1(X) & \longrightarrow & \hat{E}_{\mathbb{C}}^1(X) & \longrightarrow & H^1(X; \mathbb{Z}/2) & \longrightarrow & 0 \\ & & \parallel & & \parallel & & \parallel & & \\ 0 & \longrightarrow & H^3(X; \mathbb{Z}) & \longrightarrow & H^3(X; \mathbb{Z}) \times H^1(X; \mathbb{Z}/2) & \longrightarrow & H^1(X; \mathbb{Z}/2) & \longrightarrow & 0 \end{array}$$

From Proposition 3.2, Remark 3.5 and the diagram above we obtain :

$$\hat{E}_{\mathbb{Z}}^1(X) \cong H^1(X; \mathbb{Z}/2) \times E_{\mathbb{Z}}^1(X) \cong (H^1(X; \mathbb{Z}/2) \times_{tw} H^3(X, \mathbb{Z})) \oplus h(X),$$

with multiplication on the first two factors as in Proposition 3.2. This proves part (b).

Let  $j : H^3(X, \mathbb{Z}) = E_{\mathbb{C}}^1(X) \rightarrow E_{M_P}^1(X)$  be the map induced by unital  $*$ -homomorphism  $\mathbb{C} \rightarrow M_P$ . From the diagram above, Proposition 3.2 and Remark 3.5 we obtain that

$$(18) \quad \hat{E}_{M_P}^1(X) \cong H^1(X; \mathbb{Z}/2) \times_{tw} E_{M_P}^1(X)$$

where the group structure is given by  $(w, x) \cdot (w', x') = (w + w', x + x' + j(\beta(w \cup w')))$  for  $w, w' \in H^1(X, \mathbb{Z}/2)$  and  $x, x' \in E_{M_P}^1(X)$ . Note that the image of  $j$  is contained in  $\bar{E}_{M_P}^1(X)$  since  $E_{\mathbb{C}}^1(X) = \bar{E}_{\mathbb{C}}^1(X)$ . Using the isomorphism:

$$E_{M_P}^1(X) \cong H^1(X, (Z_P)_+^\times) \oplus \bar{E}_{M_P}^1(X) \cong H^1(X, (Z_P)_+^\times) \oplus \bar{E}_{\mathbb{Z}}^1(X) \otimes \mathbb{Z}_P$$

and the previous discussion, we can identify that map  $j : E_{\mathbb{C}}^1(X) \rightarrow E_{M_P}^1(X)$  with the map

$$H^3(X, \mathbb{Z}) \rightarrow H^1(X, (Z_P)_+^\times) \oplus H^3(X, \mathbb{Z}) \otimes \mathbb{Z}_P \oplus h(X) \otimes \mathbb{Z}_P,$$

induced by  $H^3(X, \mathbb{Z}) \rightarrow H^3(X, \mathbb{Z}) \otimes \mathbb{Z}_P$ ,  $h \mapsto h \otimes 1$ . Part (d) follows now from (18).  $\square$



**3.2. Comparing  $E_{\mathbb{Z}}^1(X)$  and  $bsu_{\otimes}^1(X)$  and proof of Theorem A.** In this section we will identify the summand  $h(X)$  of  $E_{\mathbb{Z}}^1(X)$  from Prop. 3.9 with the first group of the generalised cohomology theory  $bsu_{\otimes}^*(X)$ . To understand this group note that the spaces  $BU$  and  $BSU$  both have two  $H$ -space structures: one arising from the direct sum and another one from the tensor product. To distinguish them we will denote the second one by  $BU_{\otimes}$  and  $BSU_{\otimes}$ , respectively. It was first observed by Segal in [54] that  $BU_{\otimes}$  is in fact an infinite loop space. In particular, there is a cohomology theory  $X \mapsto bu_{\otimes}^*(X)$ , such that  $bu_{\otimes}^0(X) = [X, BU_{\otimes}]$ . This was later understood by May in [44] to fit into a much richer theory of units for  $E_{\infty}$ -ring spectra.  $K$ -theory provides such an  $E_{\infty}$ -ring spectrum  $KU$ , which has a unit spectrum  $gl_1(KU)$ , whose 0-connected cover is  $sl_1(KU) \simeq bu_{\otimes}$  and whose 2-connected cover gives  $bsu_{\otimes}$ .

The results in [44] are phrased in the language of  $S$ -modules and  $S$ -algebras and not in terms of symmetric spectra. Thus, in principle the cohomology theories represented by the spectrum  $gl_1(KU^{\mathbb{C}})$  constructed in Sec. 2.3 could differ from  $gl_1(KU)$  for an  $S$ -algebra  $KU$  representing topological  $K$ -theory. We will see in Sections 5 that this is not the case by proving a strong uniqueness result. In particular, Corollary 5.3 establishes a natural isomorphism

$$(19) \quad gl_1(KU^{\mathbb{C}})^1(X) \cong gl_1(KU)^1(X),$$

which implies the following proposition:

**Proposition 3.10.** *There is a natural isomorphism  $\hat{E}_{\mathbb{Z}}^1(X) \cong \hat{E}_{\mathbb{C}}^1(X) \oplus bsu_{\otimes}^1(X)$ .*

*Proof.* Consider the commutative diagram

$$(20) \quad \begin{array}{ccccc} \hat{E}_{\mathbb{Z}}^1(X) & \xrightarrow{\cong} & gl_1(KU^{\mathbb{Z}})^1(X) & \xleftarrow{\cong} & gl_1(KU^{\mathbb{C}})^1(X) \\ \downarrow \hat{r}_{\mathbb{Z}} & & \downarrow & & \downarrow \\ \hat{E}_{\mathbb{Z}}^1(X_4) & \xrightarrow{\cong} & gl_1(KU^{\mathbb{Z}})^1(X_4) & \xleftarrow{\cong} & gl_1(KU^{\mathbb{C}})^1(X_4) \end{array}$$

The horizontal arrows are isomorphisms by (4) and (9). It follows that the vertical arrows have isomorphic kernels. Thus  $h(X) = \ker \hat{r}_{\mathbb{Z}}$  is isomorphic to the kernel of the map  $gl_1(KU^{\mathbb{C}})^1(X) \rightarrow gl_1(KU^{\mathbb{C}})^1(X_4)$  and which in its turn, by Corollary 5.3, is isomorphic to the kernel of the map  $gl_1(KU)^1(X) \rightarrow gl_1(KU)^1(X_4)$  which is the map  $bu_{\otimes}^1(X) \rightarrow bu_{\otimes}^1(X_4)$  and hence isomorphic to  $bsu_{\otimes}^1(X)$ .  $\square$

As a consequence of the Adams-Priddy result on the uniqueness of  $bsu$  which we review in Sec. 4.1, for a finite CW-complex  $X$  there is a (nonnatural) isomorphism

$$(21) \quad bsu_{\otimes}^*(X) \cong bsu_{\oplus}^*(X).$$

Using this in conjunction with the results from the previous sections, we derive our main result (Theorem A from introduction) which we restate here for the convenience of the reader.

**Theorem 3.11.** *Let  $X$  be a finite CW-complex and let  $P$  be a nonempty set of prime numbers. There are (not natural) isomorphisms*

- (a)  $E_{\mathbb{Z}}^1(X) \cong H^3(X, \mathbb{Z}) \oplus k^5(X)$ .
- (b)  $E_{M_P}^1(X) \cong H^1(X, (\mathbb{Z}_P)_+^{\times}) \oplus H^3(X, \mathbb{Z}_P) \oplus k^5(X, \mathbb{Z}_P)$
- (c)  $E_{\mathcal{O}_{\infty}}^1(X) \cong (H^1(X, \mathbb{Z}/2) \times_{tw} H^3(X, \mathbb{Z})) \oplus k^5(X)$ .
- (d)  $E_{M_P \otimes \mathcal{O}_{\infty}}^1(X) \cong H^1(X, (\mathbb{Z}_P)_+^{\times}) \oplus (H^1(X, \mathbb{Z}/2) \times_{tw} H^3(X, \mathbb{Z}_P)) \oplus k^5(X, \mathbb{Z}_P)$

The (twisted) multiplication on  $H^1(X; \mathbb{Z}/2) \times H^3(X, \mathbb{Z}_P)$  is given by

$$(22) \quad (w, \tau) \cdot (w', \tau') = (w + w', \tau + \tau' + \beta_P(w \cup w'))$$

for  $w, w' \in H^1(X, \mathbb{Z}/2)$  and  $\tau, \tau' \in H^3(X, \mathbb{Z}_P)$ , where  $\beta_P: H^2(X, \mathbb{Z}/2) \rightarrow H^3(X, \mathbb{Z}_P)$  is the composition of the Bockstein homomorphism  $\beta$  with the coefficient map  $H^3(X, \mathbb{Z}) \rightarrow H^3(X, \mathbb{Z}_P)$ . The multiplication in (c) is just like in (22) with  $\beta_P$  replaced by  $\beta: H^2(X, \mathbb{Z}/2) \rightarrow H^3(X, \mathbb{Z})$ .

*Proof.* By (21) we have  $bsu_{\otimes}^*(X) \otimes \mathbb{Z}_P \cong bsu_{\oplus}^*(X) \otimes \mathbb{Z}_P$  for any set of primes  $P$ . It follows by (6) that  $bsu_{\otimes}^1(X) \otimes \mathbb{Z}_P \cong k^5(X) \otimes \mathbb{Z}_P \cong k^5(X, \mathbb{Z}_P)$ . Thus the statement follows now from Propositions 3.10 and 3.4 as  $h(X) \cong bsu_{\otimes}^1(X) \cong k^5(X)$ .  $\square$

The map  $\beta_P$  vanishes if  $H^3(X, \mathbb{Z})$  has no 2-torsion or if  $2 \in P$ .

The proof of Theorem 3.11 relies on the isomorphisms (21) and (19) that will be discussed in the next two sections.

#### 4. $K$ -THEORY AS A COMMUTATIVE $S$ -ALGEBRA AND ITS UNITS

The tensor product gives topological  $K$ -theory the structure of a multiplicative cohomology theory. The graded commutative multiplication lifts to the level of spectra in the sense of stable homotopy theory. In fact,  $K$ -theory can be represented by an  $E_{\infty}$ -ring spectrum. There are several approaches to make this precise. We have already met the commutative symmetric ring spectrum  $KU^{\mathbb{C}}$ . As we have seen in the previous sections,  $KU^{\mathbb{C}}$  is closely linked to the classification of  $C^*$ -algebra bundles. In this section we give a brief overview of  $K$ -theory as a commutative  $S$ -algebra. A lot of the results about the infinite loop spaces  $BU_{\oplus}$  and  $BU_{\otimes}$  are easiest to prove using this approach. We will only give a brief overview of  $S$ -modules and  $S$ -algebras here and refer the reader to [24] for a complete reference.

Constructing a symmetric monoidal category of spectra with the sphere spectrum  $S$  as its unit object is quite an intricate endeavour. The category  $\mathcal{S}[\mathbb{L}]$  of (coordinate free) spectra that are also algebras over the linear isometries operad  $\mathbb{L}$  has almost all of the desired properties, in particular a symmetric monoidal structure given by a smash product  $\wedge$  (see [24, Section I.5]). The only defect is that the sphere spectrum is not a unit object. This can be fixed by restricting to the full subcategory  $\mathcal{M}_S \subset \mathcal{S}[\mathbb{L}]$ , on which  $S$  acts like a unit, and the objects of  $\mathcal{M}_S$  are called  $S$ -modules [24, Def. II.1.1].

A (commutative)  $S$ -algebra is a (commutative) monoid with respect to  $\wedge$  in  $\mathcal{M}_S$ . We will write  $\text{Com}_S$  for the category of commutative  $S$ -algebras [24, Section II.3]. Both,  $\mathcal{M}_S$  and  $\text{Com}_S$ , are model categories [24, Section VII.4]; the weak equivalences are the morphisms that induce an isomorphism on  $\pi_*$ .

A commutative  $S$ -algebra representing connective  $K$ -theory can be constructed from the bipermutative category of finite-dimensional complex inner product spaces and unitary isomorphisms as follows: Let  $\mathcal{U}$  be the topological category with objects  $\mathbb{N}_0$ , i.e. the natural numbers including 0, where we think of  $n \in \mathbb{N}_0$  as  $\mathbb{C}^n$ . The morphism spaces are given by

$$\text{hom}(m, n) = \begin{cases} \emptyset & \text{if } m \neq n, \\ U(n) & \text{if } m = n, \end{cases}$$

where  $U(0)$  is the trivial group. The sum and product operations on  $\mathbb{N}_0$  extend to the morphisms of  $\mathcal{U}$  via the block sum  $\oplus$  and the Kronecker product  $\otimes$  of unitary matrices. This gives  $(\mathcal{U}, \oplus, \otimes)$  the structure of a bipermutative category.

Combining [45, Section 3, discussion on p. 24] and [24, Cor. 3.6] shows that there is a commutative  $S$ -algebra  $ku$  associated to  $\mathcal{U}$  that represents connective complex topological  $K$ -theory as a multiplicative cohomology theory by [44, VIII.2.1]. Its periodic counterpart, the commutative  $S$ -algebra  $KU$ , can be constructed by inverting the Bott element  $\beta \in \pi_2(ku)$ . (In fact,  $KU$  is even a commutative  $ku$ -algebra by [24, Thm. VIII.4.3].)

The 0th space of an  $\Omega$ -spectrum is an infinite loop space. Extending this observation to the category  $\mathcal{S}[\mathbb{L}]$  we obtain a functor  $\Omega^\infty: \mathcal{S}[\mathbb{L}] \rightarrow \mathcal{T}[\mathbb{L}]$  with codomain given by the  $\mathbb{L}$ -algebras in the category of based topological spaces  $\mathcal{T}$ . Let  $R \in \text{Com}_S \subset \mathcal{S}[\mathbb{L}]$  be a commutative  $S$ -algebra. The abelian group  $\pi_0(\Omega^\infty R)$  turns out to be a ring with respect to the multiplication inherited from  $R$ . The space  $GL_1(R)$  of units of  $R$  is defined by the pullback diagram

$$(23) \quad \begin{array}{ccc} GL_1(R) & \longrightarrow & \Omega^\infty R \\ \downarrow & & \downarrow \\ GL_1(\pi_0(\Omega^\infty R)) & \longrightarrow & \pi_0(\Omega^\infty R) \end{array}$$

Thus,  $GL_1(R)$  consists of those components in  $\Omega^\infty R$  that are invertible with respect to the multiplication of  $R$ . The space  $GL_1(R)$  turns out to be an  $E_\infty$ -ring space. As such it gives rise to a connective spectrum  $gl_1(R)$  such that  $\Omega^\infty gl_1(R) = GL_1(R)$ . This construction can be made functorial and takes values in the category  $\mathcal{S}^\Omega$  of *weak  $\Omega$ -spectra*.

In contrast to highly structured spectra a weak  $\Omega$ -spectrum just consists of a sequence of based spaces  $(X_n)_{n \in \mathbb{N}_0}$  together with structure maps  $X_n \rightarrow \Omega X_{n+1}$  that are weak homotopy equivalences. A map of weak  $\Omega$ -spectra is a sequence of maps  $(f_n)_{n \in \mathbb{N}_0}$  with  $f_n: X_n \rightarrow Y_n$  compatible with the structure maps. An equivalence of weak  $\Omega$ -spectra  $X, Y$  is a chain of such maps

$$X \longleftarrow Z^{(1)} \longrightarrow Z^{(2)} \longleftarrow Z^{(3)} \longrightarrow \dots \longleftarrow Z^{(k)} \longrightarrow Y$$

that are levelwise weak equivalences [47, p. 209]. For details about the functor

$$gl_1: \text{Com}_S \rightarrow \mathcal{S}^\Omega$$

from commutative  $S$ -algebras to the category  $\mathcal{S}^\Omega$  we refer the reader to Section 4 and 5 in [5] which is based on the treatment of units in [44].

If we replace the full group of invertible elements in  $\pi_0(\Omega^\infty R)$  in the pullback diagram defining  $GL_1(R)$  by the trivial subgroup (consisting only of the unit element of  $\pi_0(\Omega^\infty R)$ ), then we obtain the space of special units  $SL_1(R)$  and a corresponding connective spectrum  $sl_1(R)$  (see for example [46, Def. 7.6]). As a spectrum  $sl_1(R) \rightarrow gl_1(R)$  is the 0-connected cover.

**4.1. Localizations, connective  $K$ -theory,  $bsu_\oplus$  and  $bsu_\otimes$ .** Recall that we denote by  $KU$  and  $ku$  the commutative  $S$ -algebras representing periodic complex  $K$ -theory and its connective counterpart, respectively. The associated unit spectra are  $gl_1(KU)$  and  $gl_1(ku)$ . The localisation map  $ku \rightarrow KU$  that inverts the Bott element is a morphism of commutative  $S$ -algebras which induces an equivalence of the underlying infinite loop spaces and therefore also an equivalence

$$gl_1(ku) \rightarrow gl_1(KU) .$$

We give some background and explain below how  $gl_1(KU)$  is related to the spectrum  $bu_\otimes$  appearing in [44] (and, using slightly different machinery, in [54]). We denote by  $U$  the infinite unitary group, i.e. the colimit over the groups  $U(n)$ , and by  $SU$  the infinite special unitary group, i.e. the colimit over the groups  $SU(n)$ . As pointed out in the last section the operations of direct sum and tensor product on these groups are known to induce  $H$ -space structures on the corresponding classifying spaces. We indicate these structure by using the notation  $BU_\oplus$ ,  $BSU_\oplus$  and respectively  $BU_\otimes$ ,  $BSU_\otimes$ .

Bott periodicity shows that  $BU_\oplus$  and  $BSU_\oplus$  are infinite loop spaces. The corresponding spectra are denoted by  $bu_\oplus$  and  $bsu_\oplus$ . Note that  $GL_1(\pi_0(\Omega^\infty KU)) \cong GL_1(\pi_0(BU \times \mathbb{Z})) \cong \mathbb{Z}/2\mathbb{Z}$ .

Hence, we obtain

$$\begin{aligned} SL_1(KU) &\simeq BU_{\otimes} , \\ GL_1(KU) &\simeq \mathbb{Z}/2\mathbb{Z} \times BU_{\otimes} . \end{aligned}$$

Note that the first equivalence can be used to equip  $BU_{\otimes}$  with an infinite loop space structure. We will therefore denote the spectrum  $sl_1(KU)$  also by  $bu_{\otimes}$ . Let  $bsu_{\otimes}$  be the 2-connected cover of the spectrum  $bu_{\otimes}$  and note that  $\Omega^{\infty}bsu_{\otimes} \simeq BSU_{\otimes}$ . Denote by  $BGL_1(KU)$  the first delooping of  $GL_1(KU)$ , i.e. the first space in the sequence forming the spectrum  $gl_1(KU)$ . Likewise, let  $BBSU_{\otimes}$  be the first delooping of  $BSU_{\otimes}$  with respect to the spectrum  $bsu_{\otimes}$ . By [44, Lem. V.3.1] (see also [41, p.406]) we have a splitting of infinite loop spaces

$$(24) \quad BU_{\otimes} \simeq K(\mathbb{Z}, 2) \times BSU_{\otimes} .$$

While it is true that the spaces  $GL_1(KU)$  and  $BGL_1(KU)$  also decompose as products

$$\begin{aligned} GL_1(KU) &\simeq \mathbb{Z}/2 \times BU_{\otimes} \simeq \mathbb{Z}/2 \times K(\mathbb{Z}, 2) \times BSU_{\otimes} , \\ BGL_1(KU) &\simeq K(\mathbb{Z}/2, 1) \times K(\mathbb{Z}, 3) \times BBSU_{\otimes} , \end{aligned}$$

these decompositions do not respect the infinite loop space structure as noted implicitly in [6]. Indeed, as we verify in the paper

$$[X, BGL_1(KU)] \cong (H^1(X, \mathbb{Z}/2) \times_{tw} H^3(X, \mathbb{Z})) \oplus bsu_{\otimes}^1(X)$$

which explains the twisting of the multiplication in Theorem A.

By a classic result of Adams and Priddy [4] (see also [44, Thm. V.4.2], or [46, Cor. 10.3] for a more general statement),  $BSU_{\oplus}$  and  $BSU_{\otimes}$  become equivalent as infinite loop spaces on localization at any prime  $p$ .

**Theorem 4.1** ([4]). *There is an equivalence of infinite loop spaces*

$$(BSU_{\oplus})_{(p)} \simeq (BSU_{\otimes})_{(p)} .$$

Thus, the corresponding spectra  $bsu_{\oplus}$  and  $bsu_{\otimes}$  become equivalent as spectra on localisation at any prime  $p$ . This also turns out to be true for the completions at any prime  $p$ , but we will not need that statement.

If we are only interested in computing the groups  $bsu_{\otimes}^*(X)$  and neglect naturality, then we can use the following observation: Let  $X$  be a space with the homotopy type of a finite CW-complex. By Theorem 4.1 there is a natural isomorphism  $bsu_{\oplus}^*(X, \mathbb{Z}_{(p)}) \cong bsu_{\otimes}^*(X, \mathbb{Z}_{(p)})$ . Since  $\mathbb{Z}_{(p)}$  is flat, by the universal coefficient theorem for generalized cohomology theories [2],  $bsu_{\oplus}^*(X, \mathbb{Z}_{(p)}) \cong bsu_{\oplus}^*(X) \otimes \mathbb{Z}_{(p)}$  and  $bsu_{\otimes}^*(X, \mathbb{Z}_{(p)}) \cong bsu_{\otimes}^*(X) \otimes \mathbb{Z}_{(p)}$ . Two finitely generated abelian groups which are isomorphic after localization at each prime are necessarily isomorphic as is apparent from the structure theorem of such groups. Therefore, for every finite CW-complex there is a (not natural) isomorphism

$$(25) \quad bsu_{\otimes}^*(X) \cong bsu_{\oplus}^*(X).$$

## 5. UNIQUENESS OF $gl_1$ FOR $K$ -THEORY SPECTRA

We have seen two ways of defining the unit spectrum of topological  $K$ -theory: the first one starting from the  $S$ -algebra  $KU$  and the second one from the commutative symmetric ring spectrum  $KU^{\mathbb{C}}$ . The output of both constructions is a weak  $\Omega$ -spectrum. Our goal in this section is to compare them. By [44, VIII.2.1] the spectrum  $ku$  represents connective topological  $K$ -theory, so after inverting the Bott element to obtain  $KU$ , this spectrum represents periodic topological  $K$ -theory as defined by Atiyah and Hirzebruch. Throughout this section we will use

both notations,  $KU^*$  and  $K^*$ , interchangeably to denote the cohomology theory represented by  $KU$  and similarly,  $K_*$  and  $KU_*$  for the corresponding homology theory. We will prove:

**Theorem 5.1.** *Let  $F$  be a commutative symmetric ring spectrum representing complex topological  $K$ -theory as a multiplicative cohomology theory on finite CW-complexes in the sense that there exists a natural multiplicative isomorphism  $F^*(X) \cong K^*(X)$ . Suppose also that  $F$  is a positive  $\Omega$ -spectrum. Then there is an equivalence of weak  $\Omega$ -spectra between  $gl_1(F)$  and  $gl_1(KU)$ .*

**Remark 5.2.** By Lemma 2.1 the commutative symmetric ring spectra  $KU^{\mathbb{C}}$ ,  $KU^{\mathbb{Z}}$  and  $KU^{\mathcal{O}_\infty}$  satisfy the hypotheses of the theorem.

**Corollary 5.3.** *There is a natural isomorphism of cohomology theories on finite CW complexes*

$$gl_1(KU^{\mathbb{C}})^*(X) \cong gl_1(KU)^*(X) .$$

**5.1. Change of categories.** In the rest of this section we will work with commutative  $S$ -algebras ([24]) rather than commutative symmetric ring spectra, because we need to use facts from [24, Chapters V and VIII] whose analogues for symmetric spectra are not written down in the literature. In subsections 5.2–5.6 we will prove the following:

**Theorem 5.4.** *Let  $G$  be a commutative  $S$ -algebra that represents complex topological  $K$ -theory as a multiplicative cohomology theory on finite CW-complexes. Then there is a chain of weak equivalences of commutative  $S$ -algebras between  $G$  and  $KU$ .*

In this subsection we show that Theorem 5.4 implies Theorem 5.1.

We will use a fact about the relevant model categories. First recall that if  $\mathcal{C}$  is a model category then the *homotopy category*  $\text{Ho}\mathcal{C}$  is obtained by inverting the weak equivalences ([31, Section 1.2]), so two objects in the model category become isomorphic in the homotopy category if they are connected by a chain of weak equivalences.

Recall that we write  $\mathcal{S}p^\Sigma$  for the category of symmetric spectra; we will write  $\text{Com}^\Sigma$  for the category of commutative symmetric ring spectra. These categories have model category structures given in [43, Sections 9 and 15]; the weak equivalences for these model category structures are the stable equivalences.

The category of  $S$ -modules is denoted  $\mathcal{M}_S$ . We will write  $\text{Com}_S$  for the category of commutative  $S$ -algebras ([24, Section II.3]). These are model categories ([24, Section VII.4]); the weak equivalences are the morphisms that induce an isomorphism of  $\pi_*$ .

The fact we need is

**Proposition 5.5.** *There is an equivalence of categories*

$$\Upsilon : \text{HoCom}^\Sigma \rightarrow \text{HoCom}_S$$

*If  $F \in \text{Com}^\Sigma$  then  $F$  and  $\Upsilon(F)$  represent the same cohomology theory on finite CW-complexes.*

*Proof of Proposition 5.5.* Let  $\text{Com}^{\mathcal{I}}$  be the category of commutative orthogonal ring spectra ([43, Example 4.4]). By [43, Theorem 0.7] there is an equivalence of categories

$$\Upsilon_1 : \text{HoCom}^\Sigma \rightarrow \text{HoCom}^{\mathcal{I}}$$

with the property that  $F$  and  $\Upsilon_1(F)$  represent the same cohomology theory on finite CW-complexes. By [42, Theorem 1.5] there is an equivalence of categories

$$\Upsilon_2 : \text{HoCom}^{\mathcal{I}} \rightarrow \text{HoCom}_S,$$

and by [42, Theorem 7.13]  $F$  and  $\Upsilon_2(F)$  represent the same cohomology theory on finite CW-complexes.  $\square$

Before we continue we explain how Theorem 5.4 implies Theorem 5.1. Let  $F$  be as in Theorem 5.1 and note that  $\Upsilon(F)$  is a commutative  $S$ -algebra representing  $K$ -theory as a multiplicative cohomology theory. The functor  $\Upsilon$  is the composition of the functors  $\mathbf{L}\mathbb{P}$  and  $\mathbf{L}\mathbb{N}$  in [40, Prop. 13.9] and [40, Prop. 14.1], respectively. Thus, combining these two propositions we obtain an equivalence of weak  $\Omega$ -spectra

$$gl_1(\Upsilon(F_c)) \simeq gl_1(F)$$

where  $F_c$  is a cofibrant replacement of  $F$  as a commutative symmetric ring spectrum (which still represents  $K$ -theory as a multiplicative cohomology theory). By Theorem 5.4 there is a chain of weak equivalences of commutative  $S$ -algebras between  $\Upsilon(F_c)$  and  $KU$ . Since  $gl_1$  preserves weak equivalences, it gives a chain of equivalences of weak  $\Omega$ -spectra

$$gl_1(F) \simeq gl_1(\Upsilon(F_c)) \simeq gl_1(KU) .$$

**5.2. Obstruction theory.** In subsections 5.3–5.6 we will prove:

**Lemma 5.6.** *Let  $G$  be a commutative  $S$ -algebra representing complex topological  $K$ -theory as a multiplicative cohomology theory on finite CW-complexes. Then there is an isomorphism from  $G$  to  $KU$  in the homotopy category  $\mathrm{HoM}_S$  which makes the diagrams*

$$\begin{array}{ccc} G \wedge G & \longrightarrow & KU \wedge KU \\ \downarrow & & \downarrow \\ G & \longrightarrow & KU \end{array}$$

and

$$\begin{array}{ccc} & S^0 & \\ \swarrow & & \searrow \\ G & \longrightarrow & KU \end{array}$$

commute in  $\mathrm{HoM}_S$ .

In this subsection we use Lemma 5.6 to prove Theorem 5.4.

We use the obstruction theory of Goerss and Hopkins [27], specifically Corollary 5.9 of [27] with  $E = KU$  and  $A = KU_*KU$ .<sup>1</sup>

First we consider the obstruction groups in that Corollary:

$$D_{E_*\mathcal{T}/E_*E}^{n+1}(A, \Omega^n A).$$

According to [8, Theorem 2.6], these groups are isomorphic to the Gamma cohomology groups

$$H\Gamma^{n+1}(A|E_*, \Omega^n A),$$

and according to [7, Theorem 6.2] these groups (with our choice of  $A$  and  $E$ ) are 0.

Next we observe that  $KU_*G$  is isomorphic to  $KU_*KU$  as a commutative algebra in the category of  $KU_*KU$  comodules; this is immediate from Lemma 5.6 and the definition of the comodule structures ([3, page 281]).

Now [27, Corollary 5.9] and the definition of  $\mathcal{TM}(A)$  on page 183 of [27] give a diagram of commutative  $S$ -algebras

$$(26) \quad G \rightarrow G_1 \leftarrow G_2 \rightarrow \cdots \leftarrow KU$$

in which each map is a  $KU_*$  isomorphism.

<sup>1</sup>Section 1 of [27] explains that this obstruction theory applies to the category of commutative  $S$ -algebras.



Next we apply  $KU$ -localization (see [24, Sections VIII.1 and VIII.2]). Theorem VIII.2.2 of [24] says that if  $H$  is a cell commutative  $S$ -algebra (see [24, Definition VII.4.11]) then there is a  $KU$ -localization  $\lambda: H \rightarrow \overline{H}$  with the property that  $\overline{H}$  is a commutative  $S$ -algebra and  $\lambda$  is a map of commutative  $S$ -algebras. In order to apply this theorem we need to know that for every commutative  $S$ -algebra  $J$  there is a cell commutative  $S$ -algebra  $CJ$  and a weak equivalence of commutative  $S$ -algebras  $\kappa: CJ \rightarrow J$ , and moreover this construction gives a functor  $C: \text{Com}_S \rightarrow \text{Com}_S$  and a natural transformation  $\kappa$  from  $C$  to the identity functor (see [43, Lemma 5.8] and [24, Lemma VII.5.8]).

Now consider the diagram

$$(27) \quad \begin{array}{ccccccc} G & \longrightarrow & G_1 & \longleftarrow & G_2 & \longrightarrow & \cdots & \longleftarrow & KU \\ \kappa \uparrow & & \kappa \uparrow & & \kappa \uparrow & & & & \kappa \uparrow \\ CG & \longrightarrow & CG_1 & \longleftarrow & CG_2 & \longrightarrow & \cdots & \longleftarrow & CKU \\ \lambda \downarrow & & \lambda \downarrow & & \lambda \downarrow & & & & \lambda \downarrow \\ \overline{CG} & \longrightarrow & \overline{CG_1} & \longleftarrow & \overline{CG_2} & \longrightarrow & \cdots & \longleftarrow & \overline{CKU} \end{array}$$

The maps in the third row are given by Theorem VIII.2.2 of [24], and the lower half of the diagram homotopy commutes. Because the  $\kappa$  are weak equivalences, all the maps in the second row are  $KU_*$ -isomorphisms, and because the  $\lambda$  are  $KU_*$ -isomorphisms, all the maps in the third row are  $KU_*$ -isomorphisms. Then the maps in the third row are weak equivalences, because a  $KU_*$ -isomorphism between  $KU$ -local spectra is a weak equivalence. The map  $\lambda: CKU \rightarrow \overline{CKU}$  is also a weak equivalence, because  $KU$  is  $KU$ -local, and Lemma 5.6 implies that  $G$  is weakly equivalent (as an  $S$ -module) to  $KU$ , and therefore  $CG$  is  $KU$ -local and the map  $\lambda: CG \rightarrow \overline{CG}$  is a weak equivalence. Now the diagram gives the chain of weak equivalences promised by Theorem 5.4.

**5.3. The  $S$ -module  $\Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}]$ .** Let  $\xi$  be the canonical complex line bundle over  $\mathbb{C}P^\infty$ . The classifying map for  $\xi$  is a homotopy equivalence from  $\mathbb{C}P^\infty$  to the classifying space  $BS^1$ . The multiplication of  $S^1$  induces an associative and commutative multiplication on  $BS^1$  and this gives a homotopy associative and homotopy commutative multiplication on  $\mathbb{C}P^\infty$ . This in turn gives a homotopy associative and homotopy commutative multiplication on the  $S$ -module  $\Sigma^\infty \mathbb{C}P_+^\infty$  (where  $+$  denotes a disjoint basepoint).

Next recall that if  $X$  is a based CW complex there is a natural isomorphism in the homotopy category  $\text{Ho}\mathcal{M}_S$

$$\nu: \Sigma^\infty X_+ \rightarrow \Sigma^\infty S^0 \vee \Sigma^\infty X$$

for which the composite

$$\Sigma^\infty X_+ \xrightarrow{\nu} \Sigma^\infty X \vee \Sigma^\infty S^0 \rightarrow \Sigma^\infty S^0$$

is induced by the based map

$$p: X_+ \rightarrow S^0$$

which takes  $X$  to the non-basepoint, and the composite

$$\Sigma^\infty X_+ \xrightarrow{\nu} \Sigma^\infty S^0 \vee \Sigma^\infty X \rightarrow \Sigma^\infty X$$

is induced by the based map

$$q: X_+ \rightarrow X$$

which is the identity on  $X$ . The inverse to  $\nu$  is the map

$$\Sigma^\infty S^0 \vee \Sigma^\infty X \xrightarrow{\Sigma^\infty p' \vee Q} \Sigma^\infty X_+$$

where  $p'$  takes the non-basepoint of  $S^0$  to the basepoint of  $X$  and  $Q \circ \Sigma^\infty q$  is the map  $1 - \Sigma^\infty(p' \circ p)$ .

Let

$$a: S^2 \rightarrow \mathbb{C}P^\infty$$

be the inclusion of the 2-skeleton and let

$$b: \Sigma^\infty S^2 \rightarrow \Sigma^\infty \mathbb{C}P_+^\infty$$

be the composite

$$\Sigma^\infty S^2 \xrightarrow{\Sigma^\infty a} \Sigma^\infty \mathbb{C}P^\infty \xrightarrow{Q} \Sigma^\infty \mathbb{C}P_+^\infty$$

Let us define the  $S$ -module  $\Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}]$  to be the telescope

$$\text{Tel } \Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty$$

where the map

$$\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty \rightarrow \Sigma^{-2n-2} \Sigma^\infty \mathbb{C}P_+^\infty$$

is left multiplication by  $\Sigma^{-2}b: S^0 \rightarrow \Sigma^{-2} \Sigma^\infty \mathbb{C}P_+^\infty$  (cf. [24, page 94]).

**Lemma 5.7.**  $\pi_*(\Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}])$  is 0 in odd degrees and in even degree  $2n$  is a copy of  $\mathbb{Z}$  generated by  $b^n$ .

*Proof.*  $\pi_*(\Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}])$  is isomorphic to  $(\pi_* \Sigma^\infty \mathbb{C}P_+^\infty)[b^{-1}]$  (cf. [24, bottom of page 94]), so the lemma follows from a theorem of Snaith [56, Theorem 2.12 and equation 2.4].  $\square$

**5.4. Equivalences from  $\Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}]$  to  $KU$  and  $G$ .** First we construct a weak equivalence

$$\eta: \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] \rightarrow KU$$

The canonical bundle  $\xi$  over  $\mathbb{C}P^\infty$  restricts to a bundle  $\xi_n$  over  $\mathbb{C}P^n$ , which gives a sequence of elements  $x_n \in K^0(\mathbb{C}P^n)$ , and this sequence gives an element  $x \in K^0(\mathbb{C}P^\infty)$  by [3, Proposition III.8.1 and Exercise (ii) on page 222].  $x$  gives a map

$$\bar{x}: \Sigma^\infty \mathbb{C}P_+^\infty \rightarrow KU$$

**Lemma 5.8.** *The composite*

$$\Sigma^\infty S^2 \xrightarrow{b} \Sigma^\infty \mathbb{C}P_+^\infty \xrightarrow{\bar{x}} KU$$

*is  $\pm\beta$ , where  $\beta$  is the Bott element.*

*Proof.* It suffices to show that the element of  $\tilde{K}^0(S^2)$  represented by this composite is a generator. With the notation of Subsection 5.3, it suffices to show that  $Q^*$  takes the class  $x_2$  to a generator of  $\tilde{K}^0(S^2)$ . In the direct sum diagram

$$\tilde{K}^0(S^2) \begin{array}{c} \xleftarrow{q^*} \\ \xrightarrow{Q^*} \end{array} \tilde{K}^0(S_+^2) \begin{array}{c} \xleftarrow{p^*} \\ \xrightarrow{(p')^*} \end{array} \tilde{K}^0(S^0)$$

we have

$$q^*Q^*(x_2) = x_2 - p^*(p')^*(x_2) = x_2 - 1$$

Now  $q^*$  maps isomorphically to the image of  $\text{id} - p^*(p')^*$  and  $x_2 - 1$  generates the image of  $\text{id} - p^*(p')^*$ , so  $Q^*(x_2)$  is a generator as required.  $\square$

Using this lemma, there is a map

$$\text{Tel } \Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty \rightarrow KU$$

whose restriction to  $\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty$  is the composite

$$\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty \xrightarrow{\Sigma^{-2n} \bar{x}} \Sigma^{-2n} KU \xrightarrow{\beta^{-n}} KU$$

for each  $n$ , and moreover this map is unique up to homotopy by [3, Proposition III.8.1 and Exercise (ii) on page 222]. Define

$$\eta: \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] \rightarrow KU$$

to be this map.

Next, using the fact that  $G$  represents  $K$ -theory on finite CW-complexes, the argument that constructed  $\eta$  gives a map

$$\bar{y}: \Sigma^\infty \mathbb{C}P_+^\infty \rightarrow G$$

and a weak equivalence

$$\theta: \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] \rightarrow G$$

**5.5. A product map for  $\mathbb{C}P_+^\infty[b^{-1}]$ .** In this subsection we construct a map

$$\mu: \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] \wedge \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] \rightarrow \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}]$$

in  $\text{Ho}\mathcal{M}_S$ .

First observe that there is a weak equivalence

$$\text{Tel } ((\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty) \wedge (\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty)) \xrightarrow{\simeq} (\text{Tel } \Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty) \wedge (\text{Tel } \Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty)$$

induced by the diagonal map of the unit interval. Next observe that, because the multiplication of  $\Sigma^\infty \mathbb{C}P_+^\infty$  is homotopy associative and homotopy commutative, there is a map

$$m: \text{Tel } ((\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty) \wedge (\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty)) \rightarrow \text{Tel } \Sigma^{-4n} \Sigma^\infty \mathbb{C}P_+^\infty$$

whose restriction to  $(\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty) \wedge (\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty)$  is the composite

$$(\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty) \wedge (\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty) \rightarrow \Sigma^{-4n} (\mathbb{C}P_+^\infty \wedge \mathbb{C}P_+^\infty) \rightarrow \Sigma^{-4n} \Sigma^\infty \mathbb{C}P_+^\infty$$

for each  $n$ . Moreover, up to homotopy there is only one map  $m$  with this property, by [3, Proposition III.8.1 and Exercise (ii) on page 222] (using the fact that multiplication by  $\beta$  is a weak equivalence on the target of  $m$ ). Now let  $\mu$  be the composite

$$\begin{aligned} \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] \wedge \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] &= (\text{Tel } \Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty) \wedge (\text{Tel } \Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty) \simeq \\ &\text{Tel } ((\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty) \wedge (\Sigma^{-2n} \Sigma^\infty \mathbb{C}P_+^\infty)) \xrightarrow{m} \text{Tel } \Sigma^{-4n} \Sigma^\infty \mathbb{C}P_+^\infty \simeq \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] \end{aligned}$$

**5.6. Proof of Lemma 5.6.** To prove Lemma 5.6 it suffices to show that the diagrams

$$(28) \quad \begin{array}{ccc} & \Sigma^\infty S^0 & \\ & \swarrow & \searrow \\ \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] & \xrightarrow[\simeq]{\eta} & KU \end{array}$$

$$(29) \quad \begin{array}{ccc} \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] \wedge \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] & \xrightarrow[\simeq]{\eta \wedge \eta} & KU \wedge KU \\ \downarrow \mu & & \downarrow \\ \Sigma^\infty \mathbb{C}P_+^\infty & \xrightarrow[\simeq]{\eta} & KU \end{array}$$

$$(30) \quad \begin{array}{ccc} & \Sigma^\infty S^0 & \\ & \swarrow & \searrow \\ \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] & \xrightarrow[\simeq]{\theta} & G \end{array}$$

and

$$(31) \quad \begin{array}{ccc} \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] \wedge \Sigma^\infty \mathbb{C}P_+^\infty[b^{-1}] & \xrightarrow[\simeq]{\theta \wedge \theta} & G \wedge G \\ \downarrow \mu & & \downarrow \\ \Sigma^\infty \mathbb{C}P_+^\infty & \xrightarrow[\simeq]{\theta} & G \end{array}$$

commute in  $\text{HoM}_S$ .

We will give the proof for diagrams (28) and (29); the proof for the other two diagrams is the same.

**Lemma 5.9.** *The diagrams*

$$(32) \quad \begin{array}{ccc} & \Sigma^\infty S^0 & \\ & \swarrow & \searrow \\ \Sigma^\infty \mathbb{C}P_+^\infty & \xrightarrow{\bar{x}} & KU \end{array}$$

and

$$(33) \quad \begin{array}{ccc} \Sigma^\infty \mathbb{C}P_+^\infty \wedge \Sigma^\infty \mathbb{C}P_+^\infty & \xrightarrow{\bar{x} \wedge \bar{x}} & KU \wedge KU \\ \downarrow & & \downarrow \\ \Sigma^\infty \mathbb{C}P_+^\infty & \xrightarrow{\bar{x}} & KU \end{array}$$

commute in  $\text{HoM}_S$ .

*Proof.* Diagram (32) commutes because the restriction of  $x$  to  $\mathbb{C}P^0$  is the standard generator of  $K(\mathbb{C}P^0)$ . For diagram (33) we need to show that the pullback of  $x$  along the map

$$\Sigma^\infty \mathbb{C}P_+^\infty \wedge \Sigma^\infty \mathbb{C}P_+^\infty \rightarrow \Sigma^\infty \mathbb{C}P_+^\infty$$

is the exterior product  $x \times x$ . For this it suffices, by [3, Proposition III.8.1 and Exercise (ii) on page 222], to show that the pullback of  $x$  along the map

$$\Sigma^\infty \mathbb{C}P_+^n \wedge \Sigma^\infty \mathbb{C}P_+^n \rightarrow \Sigma^\infty \mathbb{C}P_+^\infty$$

is  $x_n \times x_n$  for each  $n$ . The map

$$\mathbb{C}P^n \times \mathbb{C}P^n \rightarrow \mathbb{C}P^\infty \times \mathbb{C}P^\infty \rightarrow \mathbb{C}P^\infty$$

factors up to homotopy through  $\mathbb{C}P^{2n}$  by cellular approximation, so we have a homotopy commutative diagram

$$(34) \quad \begin{array}{ccc} \mathbb{C}P^n \times \mathbb{C}P^n & \longrightarrow & \mathbb{C}P^\infty \times \mathbb{C}P^\infty \\ \downarrow & & \downarrow \\ \mathbb{C}P^{2n} & \longrightarrow & \mathbb{C}P^\infty \end{array}$$

In this diagram, the pullback of the bundle  $\xi$  along the clockwise composite is isomorphic to the pullback of the bundle  $\xi$  along the counterclockwise composite, so the pullback of  $\xi_{2n}$  along

the left vertical arrow is  $\xi_n \otimes \xi_n$ . Applying  $K$ -theory to diagram (34) gives the commutative diagram

$$\begin{array}{ccc} K^0(\mathbb{C}P^n \times \mathbb{C}P^n) & \longleftarrow & K^0(\mathbb{C}P^\infty \times \mathbb{C}P^\infty) \\ \uparrow & & \uparrow \\ K^0(\mathbb{C}P^{2n}) & \longleftarrow & K^0(\mathbb{C}P^\infty) \end{array}$$

We have just shown that the image of  $x_{2n}$  under that left vertical arrow is  $x_n \times x_n$ , so the image of  $x$  under the counterclockwise composite is  $x_n \times x_n$  as required.  $\square$

Now the commutativity of diagram (32) implies that of (28), so it only remains to show commutativity of diagram (29). By [3, Proposition III.8.1 and Exercise (ii) on page 222] it suffices to show that the two composites in diagram (29) have homotopic restrictions to  $(\Sigma^{-2n}\Sigma^\infty\mathbb{C}P_+^\infty) \wedge (\Sigma^{-2n}\Sigma^\infty\mathbb{C}P_+^\infty)$  for each  $n$ , and this follows from Lemma 5.9.

**Acknowledgements** We would like to thank Mike Mandell for pointing out that the isomorphism (25) follows from the Adams-Priddy theorem. The third author would like to thank John Lind for helpful discussions about the two incarnations of unit spectra used in this paper.

## REFERENCES

- [1] J. Aastrup, R. Nest, and E. Schrohe. A continuous field of  $C^*$ -algebras and the tangent groupoid for manifolds with boundary. *J. Funct. Anal.*, 237(2):482–506, 2006. 2
- [2] J. F. Adams. Lectures on generalised cohomology. In *Category Theory, Homology Theory and their Applications, III (Battelle Institute Conference, Seattle, Wash., 1968, Vol. Three)*, pages 1–138. Springer, Berlin, 1969. 18
- [3] J. F. Adams. *Stable homotopy and generalised homology*. University of Chicago Press, Chicago, Ill.-London, 1974. Chicago Lectures in Mathematics. 20, 22, 23, 24, 25
- [4] J. F. Adams and S. B. Priddy. Uniqueness of  $BSO$ . *Math. Proc. Cambridge Philos. Soc.*, 80(3):475–509, 1976. 3, 8, 18
- [5] M. Ando, A. J. Blumberg, D. Gepner, M. J. Hopkins, and C. Rezk. Units of ring spectra, orientations and Thom spectra via rigid infinite loop space theory. *J. Topol.*, 7(4):1077–1117, 2014. 17
- [6] M. Atiyah and G. Segal. Twisted  $K$ -theory. *Ukr. Mat. Visn.*, 1(3):287–330, 2004. 18
- [7] A. Baker and B. Richter. On the  $\Gamma$ -cohomology of rings of numerical polynomials and  $E_\infty$  structures on  $K$ -theory. *Comment. Math. Helv.*, 80(4):691–723, 2005. 20
- [8] M. Basterra and B. Richter. (Co-)homology theories for commutative ( $S$ -)algebras. In *Structured ring spectra*, volume 315 of *London Math. Soc. Lecture Note Ser.*, pages 115–131. Cambridge Univ. Press, Cambridge, 2004. 20
- [9] É. Blanchard and E. Kirchberg. Global Glimm halving for  $C^*$ -bundles. *J. Oper. Theory*, 52:385–420, 2004. 1
- [10] K. S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994. Corrected reprint of the 1982 original. 11
- [11] A. Connes. *Noncommutative Geometry*. Academic Press, San Diego, CA, 1994. 2
- [12] A. Connes and N. Higson. Déformations, morphismes asymptotiques et  $K$ -théorie bivariante. *C. R. Acad. Sci. Paris Sér. I Math.*, 311(2):101–106, 1990. 1
- [13] J. Cuntz. Simple  $C^*$ -algebras generated by isometries. *Comm. Math. Phys.*, 57:173–185, 1977. 4
- [14] M. Dadarlat, I. Hirshberg, and N. C. Phillips. Simple nuclear  $C^*$ -algebras not equivariantly isomorphic to their opposites. arXiv:1602.04612, 2016. 2
- [15] M. Dadarlat and J. McClure. When are two commutative  $C^*$ -algebras stably homotopy equivalent? *Math. Z.*, 235(3):499–523, 2000. 9

- [16] M. Dadarlat and A. Némethi. Shape theory and (connective)  $K$ -theory. *J. Operator Theory*, 23(2):207–291, 1990. [9](#)
- [17] M. Dadarlat and U. Pennig. A Dixmier–Douady theory for strongly self-absorbing  $C^*$ -algebras II: the Brauer group. *J. Noncommut. Geom.*, 9(4):1137–1154, 2015. [11](#), [13](#)
- [18] M. Dadarlat and U. Pennig. Unit spectra of  $K$ -theory from strongly self-absorbing  $C^*$ -algebras. *Algebr. Geom. Topol.*, 15(1):137–168, 2015. [2](#), [3](#), [6](#), [9](#), [10](#), [11](#)
- [19] M. Dadarlat and U. Pennig. A Dixmier–Douady theory for strongly self-absorbing  $C^*$ -algebras. *J. Reine Angew. Math.*, 718:153–181, 2016. [2](#), [9](#), [11](#), [12](#)
- [20] M. Dadarlat and U. Pennig. Bundles of strongly self-absorbing  $C^*$ -algebras with a Clifford grading. preprint 2022. [9](#), [10](#), [11](#), [12](#), [13](#), [14](#)
- [21] M. Dadarlat and W. Winter. On the  $KK$ -theory of strongly self-absorbing  $C^*$ -algebras. *Math. Scand.*, 104(1):95–107, 2009. [2](#)
- [22] J. Dixmier.  *$C^*$ -algebras*. North Holland, Amsterdam, 1982. [1](#)
- [23] P. Donovan and M. Karoubi. Graded Brauer groups and  $K$ -theory with local coefficients. *Inst. Hautes Études Sci. Publ. Math.*, (38):5–25, 1970. [10](#)
- [24] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May. *Rings, modules, and algebras in stable homotopy theory*, volume 47 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997. With an appendix by M. Cole. [16](#), [19](#), [21](#), [22](#)
- [25] J. M. G. Fell. The structure of algebras of operator fields. *Acta Math.*, 106:233–280, 1961. [1](#)
- [26] J. Gabe and G. Szabó. The dynamical Kirchberg–Phillips theorem. preprint 2022, arXiv:2205.04933 [math.OA]. [2](#)
- [27] P. G. Goerss and M. J. Hopkins. Moduli spaces of commutative ring spectra. In *Structured ring spectra*, volume 315 of *London Math. Soc. Lecture Note Ser.*, pages 151–200. Cambridge Univ. Press, Cambridge, 2004. [20](#)
- [28] N. Higson. The Mackey analogy and  $K$ -theory. preprint 2007. [2](#)
- [29] N. Higson and E. Guentner. Group  $C^*$ -algebras and  $K$ -theory. In *Noncommutative geometry*, volume 1831 of *Lecture Notes in Math.*, pages 137–251. Springer, Berlin, 2004. [2](#)
- [30] N. Higson and G. Kasparov.  $E$ -theory and  $KK$ -theory for groups which act properly and isometrically on Hilbert space. *Invent. Math.*, 144(1):23–74, 2001. [2](#)
- [31] M. Hovey. *Model categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999. [19](#)
- [32] M. Hovey, B. Shipley, and J. Smith. Symmetric spectra. *J. Amer. Math. Soc.*, 13(1):149–208, 2000.
- [33] M. Izumi. Group actions on operator algebras. In *Proceedings of the International Congress of Mathematicians. Volume III*, pages 1528–1548. Hindustan Book Agency, New Delhi, 2010. [2](#)
- [34] M. Izumi and H. Matui. Poly- $\mathbb{Z}$  group actions on Kirchberg algebras II. *Invent. Math.*, 224(3):699–766, 2021. [2](#)
- [35] X. Jiang and H. Su. On a simple unital projectionless  $C^*$ -algebra. *American J. Math.*, 121(2):359–413, 1999. [4](#)
- [36] M. Joachim. Higher coherences for equivariant  $K$ -theory. In *Structured ring spectra*, volume 315 of *London Math. Soc. Lecture Note Ser.*, pages 87–114. Cambridge Univ. Press, Cambridge, 2004. [4](#), [5](#)
- [37] G. G. Kasparov. Equivariant  $KK$ -theory and the Novikov conjecture. *Invent. Math.*, 91(1):147–201, 1988. [2](#)
- [38] E. Kirchberg. The classification of purely infinite  $C^*$ -algebras using Kasparov’s theory. preprint, 1994. [4](#)
- [39] N. P. Landsman. *Mathematical topics between classical and quantum mechanics*. Springer Monographs in Mathematics. Springer-Verlag, New York, 1998. [2](#)
- [40] J. A. Lind. Diagram spaces, diagram spectra and spectra of units. *Algebr. Geom. Topol.*, 13(4):1857–1935, 2013. [7](#), [8](#), [20](#)
- [41] I. Madsen, V. Snaitch, and J. Tornehave. Infinite loop maps in geometric topology. *Math. Proc. Cambridge Philos. Soc.*, 81(3):399–430, 1977. [18](#)



- [42] M. A. Mandell and J. P. May. Equivariant orthogonal spectra and  $S$ -modules. *Mem. Amer. Math. Soc.*, 159(755):x+108, 2002. [19](#)
- [43] M. A. Mandell, J. P. May, S. Schwede, and B. Shipley. Model categories of diagram spectra. *Proc. London Math. Soc. (3)*, 82(2):441–512, 2001. [7](#), [8](#), [19](#), [21](#)
- [44] J. P. May.  *$E_\infty$  ring spaces and  $E_\infty$  ring spectra*. Lecture Notes in Mathematics, Vol. 577. Springer-Verlag, Berlin-New York, 1977. With contributions by Frank Quinn, Nigel Ray, and Jørgen Tornehave. [15](#), [16](#), [17](#), [18](#)
- [45] J. P. May. Multiplicative infinite loop space theory. *J. Pure Appl. Algebra*, 26(1):1–69, 1982. [16](#)
- [46] J. P. May. What are  $E_\infty$  ring spaces good for? In *New topological contexts for Galois theory and algebraic geometry (BIRS 2008)*, volume 16 of *Geom. Topol. Monogr.*, pages 331–365. Geom. Topol. Publ., Coventry, 2009. [17](#), [18](#)
- [47] J. P. May and R. Thomason. The uniqueness of infinite loop space machines. *Topology*, 17(3):205–224, 1978. [17](#)
- [48] R. Meyer. On the classification of group actions on  $C^*$ -algebras up to equivariant KK-equivalence. *Ann. K-Theory*, 6(2):157–238, 2021. [2](#)
- [49] E. M. Parker. The Brauer group of graded continuous trace  $C^*$ -algebras. *Trans. Amer. Math. Soc.*, 308(1):115–132, 1988. [10](#)
- [50] U. Pennig. A non-commutative model for higher twisted  $K$ -theory. *J. Topol.*, 9(1):27–50, 2016. [2](#)
- [51] N. C. Phillips. A classification theorem for nuclear purely infinite simple  $C^*$ -algebras. *Documenta Math.*, (5):49–114, 2000. [4](#)
- [52] M. A. Rieffel. Quantization and  $C^*$ -algebras. In  *$C^*$ -algebras: 1943–1993 (San Antonio, TX, 1993)*, volume 167 of *Contemp. Math.*, pages 66–97. Amer. Math. Soc., Providence, RI, 1994. [2](#)
- [53] C. Schlichtkrull. Units of ring spectra and their traces in algebraic  $K$ -theory. *Geom. Topol.*, 8:645–673 (electronic), 2004. [3](#), [7](#)
- [54] G. Segal. Categories and cohomology theories. *Topology*, 13:293–312, 1974. [15](#), [17](#)
- [55] G. Segal.  $K$ -homology theory and algebraic  $K$ -theory. pages 113–127. Lecture Notes in Math., Vol. 575, 1977. [9](#)
- [56] V. Snaith. Localized stable homotopy of some classifying spaces. *Math. Proc. Cambridge Philos. Soc.*, 89(2):325–330, 1981. [22](#)
- [57] A. S. Toms and W. Winter. Strongly self-absorbing  $C^*$ -algebras. *Trans. Amer. Math. Soc.*, 359(8):3999–4029, 2007. [2](#)
- [58] S. Toms, A and W. Winter. Strongly self-absorbing  $C^*$ -algebras. *Trans. Amer. Math. Soc.*, 359(8):3999–4029 (electronic), 2007. [4](#), [5](#)
- [59] J. Trout. On graded  $\mathbb{K}$ -theory, elliptic operators and the functional calculus. *Illinois Journal of Mathematics*, 44(2):294–309, 2000. [6](#)
- [60] J.-L. Tu. La conjecture de Baum-Connes pour les feuilletages moyennables. *K-Theory*, 17(3):215–264, 1999. [2](#)
- [61] W. Winter. Strongly self-absorbing  $C^*$ -algebras are  $\mathcal{Z}$ -stable. *J. Noncommut. Geom.*, 5(2):253–264, 2011. [2](#)
- [62] W. Winter. QDQ vs. UCT. In *Operator algebras and applications—the Abel Symposium 2015*, volume 12 of *Abel Symp.*, pages 327–348. Springer, [Cham], 2017. [2](#), [4](#)

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, USA

SCHOOL OF MATHEMATICS, CARDIFF UNIVERSITY, CARDIFF, CF24 4AG, WALES, UK