

Systems of differential equations

$$\frac{d}{dt} \begin{pmatrix} u_1(t) \\ u_2(t) \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} u_1(t) \\ u_2(t) \end{pmatrix}$$

They arise in many applications

① Dynamical systems for planet/satellite motion

(FYI only;
NOT Required)

$$\begin{cases} x''(t) = \frac{-GM}{[x(t)^2 + y(t)^2]^{3/2}} x(t) \\ y''(t) = \frac{-GM}{[x(t)^2 + y(t)^2]^{3/2}} y(t) \end{cases}$$

This is a nonlinear ODE,

for which we need to understand linear ODE first.

② Partial Differential Equations (FYI only;
NOT Required)

Example: The heat equation describes how heat diffuses on a given a region.

Consider a long thin metal bar



Let $u(x, t)$ be the temperature of the bar at location x and time t .

The heat equation is given as

$$\frac{\partial}{\partial t} u(x, t) = c \frac{\partial^2}{\partial x^2} u(x, t)$$

$$\text{or } u_t = c u_{xx}$$


where c is a constant for thermal diffusivity of the metal bar.

An initial value problem is $\begin{cases} u_t = c u_{xx} \\ u(x, 0) = f(x) \end{cases}$

for a given initial temperature $f(x)$.

To approximate the solution, we consider grid points x_i with uniform spacing Δx

$$x_i = i\Delta x$$



Let $u_i(t) = u(x_i, t)$, denoted by u_i .

The second order derivative can be approximated by

$$u_{xx}(x_i, t) \approx \frac{u_{i+1} - 2u_i + u_{i-1}}{\Delta x^2}$$

because

$$u(x_{i+1}) = u(x_i + \Delta x) \approx u(x_i) + \Delta x u_x(x_i) + \frac{\Delta x^2}{2} u_{xx}(x_i) + O(\Delta x^3)$$


$$u(x_{i-1}) = u(x_i - \Delta x) \approx u(x_i) - \Delta x u_x(x_i) + \frac{\Delta x^2}{2} u_{xx}(x_i) + O(\Delta x^3)$$

$$\Rightarrow \frac{u_{i+1} - 2u_i + u_{i-1}}{\Delta x^2} \approx u_{xx}(x_i) + O(\Delta x)$$

Suppose we consider a finite interval $x \in [0, 1]$

and impose boundary condition $u(0, t) = u(1, t) = 0$

which means the bar is cold on two ends.



$$u_0(t) = 0$$

$$u_t = c u_{xx}$$

$$u_4(t) = 0$$

$$\frac{d}{dt} u_1(t) = c \frac{u_0 - 2u_1 + u_2}{\Delta x^2} = c \frac{-2u_1 + u_2}{\Delta x^2}$$

$$\frac{d}{dt} u_2(t) = c \frac{u_1 - 2u_2 + u_3}{\Delta x^2}$$

$$\frac{d}{dt} u_3(t) = c \frac{u_2 - 2u_3 + u_4}{\Delta x^2} = c \frac{u_2 - 2u_3}{\Delta x^2}$$

$$\Rightarrow \frac{d}{dt} \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix} = \frac{c}{\Delta x^2} \begin{pmatrix} -2 & 1 & 0 \\ 1 & -2 & 1 \\ 0 & 1 & -2 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix}$$

More points \Rightarrow $\left\{ \begin{array}{l} \cdot \text{ Better approximation} \\ \cdot \text{ More equations} \\ \cdot \text{ Larger system/matrix} \end{array} \right.$
 Smaller Δx

$$\begin{array}{l} \textcircled{1} \quad y'''(t) - 2y''(t) - 3y'(t) + y(t) = 0, \quad \vec{u} = \begin{pmatrix} y \\ y' \\ y'' \end{pmatrix} \\ \textcircled{2} \quad \frac{d}{dt} \vec{u}(t) = A \vec{u}(t), \quad \vec{u}(t) \in \mathbb{R}^3, \quad A \in \mathbb{R}^{3 \times 3} \end{array} \quad (*)$$

Consider all solutions to $(*)$, which form a subset

of $V = \left\{ \vec{u} = \begin{pmatrix} u_1(t) \\ u_2(t) \\ u_3(t) \end{pmatrix} : u_1(t), u_2(t), u_3(t) \text{ are differentiable} \right\}$

① V is an abstract vector space

1) $+$ and \cdot are defined

2) closed under two ops

② Let $\vec{v}(t), \vec{w}(t)$ be solutions to $(*)$

$$\frac{d}{dt} \vec{v}(t) = A \vec{v}(t)$$

$$\frac{d}{dt} \vec{w}(t) = A \vec{w}(t)$$

$$\Rightarrow \frac{d}{dt} (\vec{v} + \vec{w}) = A (\vec{v} + \vec{w})$$

$$\left\{ \begin{array}{l} \frac{d}{dt} [a \vec{v}(t)] = A [a \vec{v}(t)] \end{array} \right. , a \in \mathbb{R}$$

If S is the set of all solutions to $(*)$,

then S is closed under $+$ and \cdot .

So $S \subseteq V$ is a subspace.

③ The dimension of S is less than or equal to 3.

If $\vec{v}_1(t), \vec{v}_2(t), \vec{v}_3(t), \vec{v}_4(t) \in S$,

$$a \vec{v}_1(t) + b \vec{v}_2(t) + c \vec{v}_3(t) + d \vec{v}_4(t) = \vec{0}$$

$$\Rightarrow a \vec{v}_1(t_0) + b \vec{v}_2(t_0) + c \vec{v}_3(t_0) + d \vec{v}_4(t_0) = \vec{0}$$

for any fixed t_0

for example, $t_0 = 1$

$$a \vec{v}_1(t) + b \vec{v}_2(t) + c \vec{v}_3(t) + d \vec{v}_4(t) = \vec{0}$$

$$\vec{v}_1(t), \vec{v}_2(t), \vec{v}_3(t), \vec{v}_4(t) \in \mathbb{R}^3$$

\Rightarrow Nonzero $\begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix}$ exists \Rightarrow Dependence

④ If \vec{v} is an eigenvector of A with eigenvalue λ ,

then $\vec{u}(t) = e^{\lambda t} \vec{v}$ is a sol to (*)

$$\text{LHS} = \frac{d}{dt} \vec{u}(t) = \lambda e^{\lambda t} \vec{v}$$

$$\text{RHS} = A \vec{u}(t) = A e^{\lambda t} \vec{v} = e^{\lambda t} A \vec{v} = e^{\lambda t} \lambda \vec{v}$$

⑤

Theorem

Then $A = V D V^{-1}$ if and only if

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are linearly independent eigenvectors of A , with eigenvalues d_1, \dots, d_n .

- A being diagonalizable means dimensions of all eigenspaces sum to n .

Next lecture, we will see that

real symmetric ($A = A^T$) and real skew-symmetric ($A = -A^T$)

are always diagonalizable.

$$\begin{pmatrix} -2 & 1 & 0 & 0 \\ 1 & -2 & 1 & 0 \\ 0 & 1 & -2 & 1 \\ 0 & 0 & 1 & -2 \end{pmatrix}$$

$$\begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}$$

Assume $A \in \mathbb{R}^{3 \times 3}$ is diagonalizable, then

we have
$$\begin{cases} A \vec{v}_1 = \lambda_1 \vec{v}_1 \\ A \vec{v}_2 = \lambda_2 \vec{v}_2 \\ A \vec{v}_3 = \lambda_3 \vec{v}_3 \end{cases}$$

and $\begin{cases} \vec{v}_1, \vec{v}_2, \vec{v}_3 \text{ are independent,} \\ \lambda_1, \lambda_2, \lambda_3 \text{ may or may not be the same.} \end{cases}$

We get 3 sols to (*):

$$\begin{cases} \vec{u}_1(t) = e^{\lambda_1 t} \vec{v}_1 \\ \vec{u}_2(t) = e^{\lambda_2 t} \vec{v}_2 \\ \vec{u}_3(t) = e^{\lambda_3 t} \vec{v}_3 \end{cases}$$

Check independence:

$$a \vec{u}_1(t) + b \vec{u}_2(t) + c \vec{u}_3(t) = \vec{0}$$

$$\Rightarrow a \vec{u}_1(0) + b \vec{u}_2(0) + c \vec{u}_3(0) = \vec{0}$$

$$\Rightarrow a \vec{v}_1 + b \vec{v}_2 + c \vec{v}_3 = \vec{0}$$

$$\Rightarrow a = b = c = 0$$

\Rightarrow 3 independent sols to (*)

Recall the solution set S is at most 3-dim

So $\{\vec{u}_1(t), \vec{u}_2(t), \vec{u}_3(t)\}$ is a basis of S .

For any $\vec{w}(t) \in S$, we have

$$\vec{w} = a \vec{u}_1 + b \vec{u}_2 + c \vec{u}_3$$

$$a \vec{u}_1 + b \vec{u}_2 + c \vec{u}_3 = \vec{w}$$

$$[\vec{u}_1 \ \vec{u}_2 \ \vec{u}_3] \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \vec{w}$$

$$[e^{\lambda_1 t} \vec{v}_1 \quad e^{\lambda_2 t} \vec{v}_2 \quad e^{\lambda_3 t} \vec{v}_3] \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} w_1(t) \\ w_2(t) \\ w_3(t) \end{bmatrix}$$

Plug in $t=0$

$$\Rightarrow [\vec{v}_1 \ \vec{v}_2 \ \vec{v}_3] \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} w_1(0) \\ w_2(0) \\ w_3(0) \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} a \\ b \\ c \end{bmatrix} = [\vec{v}_1 \ \vec{v}_2 \ \vec{v}_3]^{-1} \begin{bmatrix} w_1(0) \\ w_2(0) \\ w_3(0) \end{bmatrix} \\ = V^{-1} \vec{w}(0)$$

$$\Rightarrow \vec{w} = a \vec{u}_1 + b \vec{u}_2 + c \vec{u}_3$$

$$= [\vec{u}_1 \ \vec{u}_2 \ \vec{u}_3] \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

$$= [e^{\lambda_1 t} \vec{v}_1 \quad e^{\lambda_2 t} \vec{v}_2 \quad e^{\lambda_3 t} \vec{v}_3] \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

$$= [\vec{v}_1 \ \vec{v}_2 \ \vec{v}_3] \begin{bmatrix} e^{\lambda_1 t} & 0 & 0 \\ 0 & e^{\lambda_2 t} & 0 \\ 0 & 0 & e^{\lambda_3 t} \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

$$= \begin{bmatrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \\ & V & \end{bmatrix} \begin{bmatrix} e^{\lambda_1 t} & 0 & 0 \\ 0 & e^{\lambda_2 t} & 0 \\ 0 & 0 & e^{\lambda_3 t} \end{bmatrix} \begin{bmatrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \\ & V^{-1} & \end{bmatrix} \begin{bmatrix} w_1(t_0) \\ w_2(t_0) \\ w_3(t_0) \end{bmatrix}$$

$$= e^{At} \vec{w}(t_0)$$

Example: Initial Value Problem (IVP)

$$\begin{cases} y''(t) - y'(t) - 2y(t) = 0 \\ y(0) = 0, \quad y'(0) = 1 \end{cases}$$

$$\frac{d}{dt} \begin{pmatrix} y \\ y' \end{pmatrix} = \begin{pmatrix} y' \\ y'' \end{pmatrix} = \begin{pmatrix} y' \\ y' + 2y \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} y \\ y' \end{pmatrix}$$

$$|A - \lambda I| = \begin{vmatrix} -\lambda & 1 \\ 2 & 1-\lambda \end{vmatrix} = -\lambda + \lambda^2 - 2 = (\lambda - 2)(\lambda + 1)$$

$$\Rightarrow \lambda = -1, 2$$

$$\vec{v}_1 = \begin{pmatrix} -1 \\ 1 \end{pmatrix} \quad \vec{v}_2 = \begin{pmatrix} \frac{1}{2} \\ 1 \end{pmatrix}$$

$$e^{-t} \begin{pmatrix} -1 \\ 1 \end{pmatrix} \quad e^{2t} \begin{pmatrix} \frac{1}{2} \\ 1 \end{pmatrix}$$

$$V = \begin{pmatrix} -1 & \frac{1}{2} \\ 1 & 1 \end{pmatrix}$$

$$V^{-1} = \frac{1}{-1 - \frac{1}{2}} \begin{pmatrix} 1 & -\frac{1}{2} \\ -1 & -1 \end{pmatrix} = \frac{2}{3} \begin{pmatrix} -1 & \frac{1}{2} \\ 1 & 1 \end{pmatrix}$$

$$\begin{pmatrix} y(t) \\ y'(t) \end{pmatrix} = V \begin{pmatrix} e^{-t} & 0 \\ 0 & e^{2t} \end{pmatrix} V^{-1} \begin{pmatrix} y(0) \\ y'(0) \end{pmatrix}$$

$$\begin{aligned}
 \begin{pmatrix} y(t) \\ y'(t) \end{pmatrix} &= \begin{pmatrix} -1 & \frac{1}{2} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} e^{-t} & 0 \\ 0 & e^{2t} \end{pmatrix} \frac{2}{3} \begin{pmatrix} -1 & \frac{1}{2} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\
 &= \begin{pmatrix} -1 & \frac{1}{2} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} e^{-t} & 0 \\ 0 & e^{2t} \end{pmatrix} \begin{pmatrix} \frac{1}{3} \\ \frac{2}{3} \end{pmatrix} \\
 &= \begin{pmatrix} -1 & \frac{1}{2} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{3}e^{-t} \\ \frac{2}{3}e^{2t} \end{pmatrix}
 \end{aligned}$$

$$\Rightarrow \begin{cases} y(t) = -\frac{1}{3}e^{-t} + \frac{1}{3}e^{2t} \\ y'(t) = \frac{1}{3}e^{-t} + \frac{2}{3}e^{2t} \end{cases}$$

$$\text{IVP: } \begin{cases} y''(t) - y'(t) - 2y(t) = 0 \\ y(0) = 0, \quad y'(0) = 1 \end{cases}$$

$$\begin{aligned}
 y'' &= -\frac{1}{3}e^{-t} + \frac{4}{3}e^{2t} \\
 -y' &= -\frac{1}{3}e^{-t} - \frac{2}{3}e^{2t} \\
 -2y &= +\frac{2}{3}e^{-t} - \frac{2}{3}e^{2t}
 \end{aligned}$$

$$\begin{array}{cc}
 \underbrace{y_1(t)} & \underbrace{y_2(t)} & \begin{pmatrix} y_1 \\ y_1' \end{pmatrix} & \begin{pmatrix} y_2 \\ y_2' \end{pmatrix} \\
 \left| \begin{array}{cc} y_1(t) & y_2(t) \\ y_1'(t) & y_2'(t) \end{array} \right| & = & \underbrace{y_1(t)y_2'(t)} - \underbrace{y_2(t)y_1'(t)} \neq 0
 \end{array}$$