

Stochastic Optimization

Consider $\min_{x \in \mathbb{R}^n} \frac{1}{N} \sum_{i=1}^N f_i(x)$

Example: Linear Regression for given data $(x_i, y_i)_{i=1}^N$
 $\phi_j(x), j=1, 2, \dots, n$ are some model basis
 Want to solve the eqn in least square sense.

$$\underbrace{c_1 \phi_1(x_i) + c_2 \phi_2(x_i) + \dots + c_n \phi_n(x_i)}_{g(x_i, c)} = y_i, \quad i=1, \dots, N$$

$$\min_{c \in \mathbb{R}^n} \frac{1}{N} \sum_{i=1}^N \|g(x_i, c) - y_i\|^2$$

(Full batch) Gradient Descent:

$$x_{k+1} = x_k - \eta \left[\frac{1}{N} \sum_{i=1}^N \nabla f_i(x) \right]$$

Stochastic Gradient Descent:

$$x_{k+1} = x_k - \eta_k \nabla f_{i(k)}(x_k)$$

$i(k) \in \{1, \dots, N\}$ are i.i.d. random variables
 with uniform distribution

Now consider $\begin{cases} x_{k+1} = x_k - \eta_k A(x_k) \\ x_{k+1} = x_k - \eta_k A_{i(k)}(x_k) \end{cases}$

Example: $A_i = \nabla f_i$ and $A = \frac{1}{N} \sum_{i=1}^N \nabla f_i$
 SGD GD

Def Monotone Operator

$$\langle Tx - Ty, x - y \rangle \geq 0$$

Example: $f(x)$ is convex, ∂f is monotone

$$\begin{cases} f(y) \geq f(x) + \langle \partial f(x), y - x \rangle \\ f(x) \geq f(y) + \langle \partial f(y), x - y \rangle \end{cases}$$
$$\Rightarrow \langle \partial f(x) - \partial f(y), x - y \rangle \geq 0$$

Def Graph of T is $\{(x, Tx) : \forall x\}$

Def T is maximal monotone if its graph is not a strict subset of graph of a monotone operator S .

If $f(x)$ is well-defined & continuous, ∂f is maximal

Assumptions:

① $A_i(x)$ are maximal monotone

② There are constants C_1, C_2 s.t.

$$\frac{1}{N} \sum_{i=1}^N \|A_i(x)\|^2 \leq \frac{C_1}{2} \|x\|^2 + C_2$$

Assume ∂f_i are L-const.

$$\|\partial f_i(x) - \partial f_i(0)\| \leq L \|x - 0\|$$

$$\Rightarrow \|\partial f_i(x)\| \leq L \|x\| + \|\partial f_i(0)\|$$

③ $\tau(0), \tau(1), \dots, \tau(k), \dots$ are i.i.d. R.V.

with uniform distribution in $\{1, \dots, N\}$

$$\textcircled{3} \quad \sum_{k=0}^{\infty} \eta_k = \infty, \quad \sum_{k=0}^{\infty} \eta_k^2 < \infty$$

$$\eta_k = \frac{C}{(k+1)^p}, \quad k=0, 1, 2, \dots$$

$$\frac{1}{2} < p \leq 1$$

\textcircled{4} Assume A is demipositive

For any x_* satisfying $0 \in A(x_*)$

Example: $\langle Ax, x - x_* \rangle > 0, \forall x$ and Ax does not include 0

2.2.3 Convergence for convex functions

Theorem 2.8. Assume $\nabla f(\mathbf{x})$ is Lipschitz-continuous with Lipschitz constant L and $f(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex. Then for any \mathbf{x}, \mathbf{y} :

1. $f(\mathbf{y}) \geq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle + \frac{1}{2L} \|\nabla f(\mathbf{x}) - \nabla f(\mathbf{y})\|^2$
2. $\|\nabla f(\mathbf{x}) - \nabla f(\mathbf{y})\|^2 \leq L \langle \nabla f(\mathbf{x}) - \nabla f(\mathbf{y}), \mathbf{x} - \mathbf{y} \rangle$.

1) Assume $f = \frac{1}{N} \sum_{i=1}^N f_i(\mathbf{x})$ is convex and $A = \nabla f$ is L-cont.

$$\begin{aligned} \langle \nabla f(\mathbf{x}), \mathbf{x} - x_* \rangle &\geq f(\mathbf{x}) - f(x_*) + \frac{1}{2L} \|\nabla f(\mathbf{x})\|^2 \\ &\geq \frac{1}{2L} \|\nabla f(\mathbf{x})\|^2 > 0 \end{aligned}$$

2) Assume $f(\mathbf{x})$ is convex, then $A = \nabla f$ is demipositive

$$\begin{aligned} f(\mathbf{y}) &\geq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), \mathbf{y} - \mathbf{x} \rangle \\ \Rightarrow f(x_*) &\geq f(\mathbf{x}) + \langle \nabla f(\mathbf{x}), x_* - \mathbf{x} \rangle \\ \Rightarrow \langle \nabla f(\mathbf{x}), \mathbf{x} - x_* \rangle &\geq f(\mathbf{x}) - f(x_*) > 0 \end{aligned}$$

Theorem 1

Under assumptions ①, ②, ③ and demipositivity
 for $x_{k+1} = x_k - \eta_k A_{ik_k}(x_k)$,
 and any x_0 ,

$$x_k \rightarrow x_* \text{ with prob. 1}$$

Polyak-Ruppert Averaging:

For $x_{k+1} = x_k - \eta_k A_{ik_k}(x_k)$

define averaged iterate

$$\bar{x}_k = \frac{\sum_{j=0}^k \eta_j x_k}{\sum_{j=0}^k \eta_j}$$

Theorem 2 (Convergence with averaging)

Under assumptions ①, ②, ③, without demipositivity
 for $x_{k+1} = x_k - \eta_k A_{ik_k}(x_k)$ and any x_0 ,

$$\bar{x}_k \rightarrow x_* \text{ with prob. 1}$$

$$\text{where } 0 \in A(x_*)$$

Proof of Theorem 1 :

We make this general by also including $A = \emptyset$ which is a set

$$\begin{aligned}x_{k+1} &= x_k - \eta_k A_{i(k)}(x_k) \\ \Rightarrow \|x_{k+1} - x_*\|^2 &= \|x_k - x_*\|^2 + \eta_k^2 \|A_{i(k)}(x_k)\|^2 \\ &\quad - 2\eta_k \langle A_{i(k)}(x_k), x_k - x_* \rangle\end{aligned}$$

$$\boxed{\text{Assumption } \frac{1}{N} \sum_{i=1}^N \|A_i(x)\|^2 \leq \frac{C_1}{2} \|x\|^2 + C_2}$$

$$\begin{aligned}\Rightarrow \|A_{i(k)}(x_k)\|^2 &= \|A_{i(k)}(x_k) - A(x_*)\|^2 \\ &\leq 2\|A_{i(k)}(x_k)\|^2 + 2\|A(x_*)\|^2 \\ &\leq 2(\|x_k\|^2 + \|x_*\|^2) + 4C_2\end{aligned}$$

$$\Rightarrow \|x_{k+1} - x_*\|^2 \leq \|x_k - x_*\|^2 + \eta_k^2 \left[(\|x_k\|^2 + \|x_*\|^2) + 4c_2 \right] - 2\eta_k \langle A(i_k)(x_k), x_k - x_* \rangle$$

Take conditional expectation E_k w.r.t. $i(k)$

for both sides

$$\Rightarrow E_k \|x_k - x_*\|^2 \leq \|x_k - x_*\|^2$$

$$+ \eta_k^2 \left[(\|x_k\|^2 + \|x_*\|^2) + 4c_2 \right] + E_k \left[-2\eta_k \langle A(i_k)(x_k), x_k - x_* \rangle \right]$$

||

$$-2\eta_k \langle E_k (A(i_k)(x_k)), x_k - x_* \rangle$$

||

$$-2\eta_k \langle \frac{1}{N} \sum_{i=1}^k A_i(x_k), x_k - x_* \rangle$$

||

$$-2\eta_k \langle A(x_k), x_k - x_* \rangle$$

$$\Rightarrow E_k \|x_k - x_*\|^2 \leq \|x_k - x_*\|^2$$

$$+ \eta_k^2 \left[(\|x_k\|^2 + \|x_*\|^2) + 4c_2 \right]$$

$$-2\eta_k \langle A(x_k), x_k - x_* \rangle$$

$$\|x_k\|^2 + \|x_*\|^2 \leq 2\|x_k - x_*\|^2 + 3\|x_*\|^2$$

$$E_k \|x_k - x_*\|^2 \leq (1 + 2\eta_k^2) \|x_k - x_*\|^2 + 3\eta_k^2 [\|x_*\|^2 + 4Q] \\ - 2\eta_k < \underbrace{A(x_k), x_k - x_*}_{\geq 0 \text{ due to demipositivity}}$$

Theorem [Robbins & Siegmund Quasi-Martingale Convergence Theorem]

If

① $V_k \geq 0, S_k \geq 0, U_k \geq 0$ are R. V.

② $\sum_{k=0}^{\infty} \beta_k < \infty, \beta_k \geq 0$ are numbers

③ $E_k[V_{k+1}] \leq (1 + \beta_k)V_k - S_k + U_k$

④ $\sum_{i=1}^{\infty} U_i < +\infty$

Then 1) $V_k \rightarrow V_\infty$ with prob. 1
 2) $\sum_{k=0}^{\infty} S_k < +\infty$

To apply this Quasi-Martingale Thm,

we need $\sum_{k=0}^{\infty} \eta_k^2 < +\infty$

Everything is under prob. 1

$$\Rightarrow \textcircled{1} \sum_{k=0}^{\infty} \eta_k < A(x_k), x_k - x_* > < \infty$$

$$\sum_{k=0}^{\infty} \eta_k = +\infty \quad \& \quad < A(x_k), x_k - x_* > \geq 0$$

$$\text{implies} \quad \liminf_k < A(x_k), x_k - x_* > = 0$$

If $\liminf_k < A(x_k), x_k - x_* > = c > 0$,

$$\text{then } \sum \eta_k < A(x_k), x_k - x_* >$$

is essentially at least $\sum_{k=M}^{\infty} \eta_k c = +\infty$

\textcircled{2} $\|x_k - x_*\|$ has a limit

$\Rightarrow \{x_k\}$ is bounded (thus $A(x_k)$ bounded)

\Rightarrow There is a subsequence x_{k_j} s.t.

$$x_{k_j} \rightarrow x_\infty$$

$$A(x_{k_j}) \rightarrow a_\infty$$

$$< A(x_{k_j}), x_{k_j} - x_* > \rightarrow 0$$

1) $\liminf_k < A(x_k), x_k - x_* > = 0$

\Rightarrow There is a subsequence x_{k_n} s.t.

$$\langle A(x_{k_n}), x_{k_n} - x_* \rangle \rightarrow 0$$

2) Pick another subsequence X_{kn_i} from X_{k_n}

S.t. $A(X_{kn_i}) \rightarrow a_\infty$

3) Pick a subsequence $X_{kn_{ij}}$ from X_{kn_i}

S.t. $X_{kn_{ij}} \rightarrow x_\infty$

So A is maximal monotone $\Rightarrow a_\infty \in A(x_\infty)$

$$\langle a_\infty, x_\infty - x_* \rangle = 0$$

$$\Rightarrow 0 \in \langle A(x_\infty), x_\infty - x_* \rangle$$

demi-positivity $\Rightarrow x_\infty$ is a zero point of A .