

NAME Solution

STUDENT ID # _____

RECITATION INSTRUCTOR _____


RECITATION TIME _____

DIRECTIONS

- 1) Fill in the above information. Also write your name at the top of each page of the exam.
- 2) The exam has 6 pages, including this one.
- 3) Problems 1 through 6 are multiple choice; circle the correct answer. No partial credit for these problems.
- 4) Problems 7 through 9 are problems to be worked out. Partial credit for correct work is possible. Write your answer in the box provided. **YOU MUST SHOW SUFFICIENT WORK TO JUSTIFY YOUR ANSWERS. CORRECT ANSWERS WITH INCONSISTENT WORK MAY NOT RECEIVE CREDIT.**
- 5) Points for each problem are given in parenthesis in the left margin.
- 6) No books, notes, or calculators may be used on this test.

Page 2	/16
Page 3	/16
Page 4	/16
Page 5	/22
Page 6	/30
TOTAL	/100

- (8) 1. Find the volume of the solid region in the first octant bounded above by the plane $x + z = 3$, on the sides by the planes $x + y = 1$, $x = 0$, and $y = 0$ and below by the plane $z = 0$.




$$V = \int_0^1 \int_0^{1-x} \int_0^{3-x} dz dy dx$$

- A. $\frac{5}{3}$
 B. $\frac{2}{3}$
 C. $\frac{1}{2}$
 D. $\frac{4}{3}$
 E. 2

$$= \int_0^1 (1-x)(3-x) dx$$

$$= \int_0^1 (3 - 4x + x^2) dx = \left[3x - 2x^2 + \frac{1}{3}x^3 \right]_0^1 = \frac{4}{3}$$

- (8) 2. Find the surface area of the part of the parabolic cylinder $z = y^2$ that lies over the triangle with vertices $(0, 0)$, $(0, 1)$, $(1, 1)$ in the xy -plane.



$$0 \leq x \leq y$$

$$0 \leq y \leq 1$$

$$F(x, y, z) = y^2 - z$$

$$\frac{\partial F}{\partial x} = 0, \quad \frac{\partial F}{\partial y} = 2y$$

$$\frac{\partial F}{\partial z} = -1$$

A. $\frac{1}{12}(5\sqrt{5} - 1)$

B. $\frac{5}{12}\sqrt{5}$

C. $\frac{2}{3}$

D. $\frac{1}{4}$

E. $\frac{1}{12}$

$$ds = \sqrt{\left(\frac{\partial F}{\partial x}\right)^2 + \left(\frac{\partial F}{\partial y}\right)^2 + \left(\frac{\partial F}{\partial z}\right)^2} dx dy$$

$$= \sqrt{4y^2 + 1} dx dy$$

$$S = \int_0^1 \int_0^y \sqrt{4y^2 + 1} dx dy$$

$$= \int_0^1 y \sqrt{4y^2 + 1} dy = \frac{1}{8} \int_0^1 8y (4y^2 + 1)^{1/2} dy$$

$$= \frac{1}{8} \cdot \frac{2}{3} (4y^2 + 1)^{3/2} \Big|_0^1 = \frac{1}{12} (5^{3/2} - 1)$$

- (8) 3. The iterated triple integral

$$\int_{-2}^2 \int_0^{\sqrt{4-x^2}} \int_x^{3+x^2+y^2} 10y \, dz \, dy \, dx$$

in cylindrical coordinates is:

$$y = r \sin \theta$$

$$dA = r \, dr \, d\theta$$

$$x = r \cos \theta$$

$$3 + x^2 + y^2 = 3 + r^2$$

$$y = \sqrt{4-x^2} \text{ is the semicircle } r = 2, \quad 0 \leq \theta \leq \pi$$

$$-2 \leq x \leq 2$$

$$0 \leq \theta \leq \pi$$

A. $\int_0^\pi \int_0^2 \int_{\cos \theta}^{3+r^2} 10r \sin \theta \, dz \, dr \, d\theta$

B. $\int_0^{\frac{\pi}{2}} \int_0^2 \int_{r \cos \theta}^{3+r^2} 10r \sin \theta \, dz \, dr \, d\theta$

C. $\int_0^\pi \int_{-2}^2 \int_{r \cos \theta}^{3+r^2} 10r^2 \sin \theta \, dz \, dr \, d\theta$

D. $\int_0^\pi \int_0^2 \int_{r \cos \theta}^{3+r^2} 10r^2 \sin \theta \, dz \, dr \, d\theta$

E. $\int_0^{\frac{\pi}{2}} \int_{-2}^2 \int_{r \cos \theta}^{3+r^2} 10r^2 \sin \theta \, dz \, dr \, d\theta$

- (8) 4. If
- $\vec{F}(x, y, z) = xy\vec{i} + z^2\vec{j} + e^y\vec{k}$
- then
- $\vec{F} \cdot \text{curl } \vec{F} =$

$$\text{curl } \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xy & z^2 & e^y \end{vmatrix}$$

$$= \left(\frac{\partial e^y}{\partial y} - \frac{\partial z^2}{\partial z} \right) \vec{i} + \left(\frac{\partial xy}{\partial z} - \frac{\partial e^y}{\partial x} \right) \vec{j}$$

$$+ \left(\frac{\partial z^2}{\partial x} - \frac{\partial xy}{\partial y} \right) \vec{k}$$

$$= (e^y - 2z) \vec{i} + 0 \vec{j} - x \vec{k} = (e^y - 2z) \vec{i} - x \vec{k}$$

$$\vec{F} \cdot \text{curl } \vec{F} = xy(e^y - 2z) + z \cdot 0 - xe^y$$

$$= xy(e^y - 2z) - xe^y$$

A. 0

B. xy^2

C. $xy(e^y - 2z) - xe^y$

D. $e^y(xy + 2z - x) - yz^2$

E. $e^y(xy + 2z - x)$

(8) 5. Compute $\int_C 6x \, ds$ where C is the graph of $y = x^2$ for $0 \leq x \leq 1$.

$y = x^2$ can be parametrized as
 $x(t) = t, \quad y(t) = t^2, \quad z(t) = 0, \quad 0 \leq t \leq 1$

$$ds = \frac{dr}{dt} dt = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt$$

$$= \sqrt{1 + 4t^2} dt$$

$$\therefore \int_C 6x \, ds = \int_0^1 6t \sqrt{1+4t^2} dt$$

$$= \frac{3}{4} \int_0^1 8t (1+4t^2)^{1/2} dt$$

$$= \frac{3}{4} \cdot \frac{2}{3} (1+4t^2)^{3/2} \Big|_0^1 = \frac{1}{2} (5^{3/2} - 1)$$

A. $5\sqrt{5} - 1$

B. $\frac{1}{2} (5\sqrt{5} - 1)$

C. 3

D. 2

E. $\frac{3}{2}$

(8) 6. Compute $\int_C e^x dx + 3xy \, dy + xyz \, dz$ where C is the curve parametrized by
 $\vec{r}(t) = t\vec{i} + t\vec{j} + 2t\vec{k}$ for $0 \leq t \leq 1$.

$$I = \int_0^1 (e^t \cdot 1 + 3t^2 \cdot 1 + 2t^3 \cdot 2) dt$$

$$= \int_0^1 (e^t + 3t^2 + 4t^3) dt$$

$$= \left[e^t + t^3 + t^4 \right]_0^1$$

$$= (e + 1 + 1) - e^0$$

$$= e + 1$$

A. e

B. $e + \frac{1}{3}$

C. $e + \frac{1}{2}$

D. $e + 1$

E. $e + \frac{3}{2}$

- (11) 7. Find a function
- $f(x, y)$
- whose gradient is:

$$\text{grad } f(x, y) = (3x^2 e^{2y} - y)\vec{i} + (2x^3 e^{2y} - x + 2y)\vec{j} \quad \text{and } f(1, 0) = 3.$$

$$\frac{\partial f}{\partial x} = 3x^2 e^{2y} - y.$$

$$f(x, y) = x^3 e^{2y} - xy + h(y)$$

$$\frac{\partial f}{\partial y} = 2x^3 e^{2y} - x + 2y, \quad \frac{\partial f}{\partial y} = 2x^3 e^{2y} - x + h'(y)$$

$$h'(y) = 2y, \quad h(y) = y^2 + C.$$

$$3 = f(1, 0) = 1 + C \quad \therefore C = 2$$

$$f(x, y) = x^3 e^{2y} - xy + y^2 + 2.$$

- (11) 8. Use Green's Theorem to evaluate
- $\int_C (y^3 + 2y)dx + 3xy^2 dy$
- , where
- C
- is the circle
- $x^2 + y^2 = 16$
- oriented counterclockwise.

$$\frac{\partial N}{\partial x} = 3y^2, \quad \frac{\partial M}{\partial y} = 3y^2 + 2.$$

$$\begin{aligned} \int_C (y^3 + 2y)dx + 3xy^2 dy &= \iint_R -2 dA \\ &= \int_0^{2\pi} \int_0^4 -2r dr d\theta \end{aligned}$$

$$\text{or } = -2(\text{area of circle})$$

$$-32\pi$$

- (30) 9. Let D be the solid region above the upper nappe of the cone $z^2 = x^2 + y^2$ and below the sphere $x^2 + y^2 + z^2 = 18$. If $\vec{F}(x, y, z) = 3x^2\vec{i} + \vec{j} + \frac{1}{2}z^2\vec{k}$, express the triple

$\iiint_D \operatorname{div} \vec{F} dV$ as an iterated triple integral in (a) rectangular coordinates, (b) cylindrical coordinates, and (c) spherical coordinates. (Include the limits of integration.)

(a) Rectangular coordinates

$$\operatorname{div} \vec{F} = 6x + z$$

or

$$\int_{-3}^3 \int_{-\sqrt{9-y^2}}^{\sqrt{9-y^2}} \int_{\sqrt{x^2+y^2}}^{\sqrt{18-x^2-y^2}} (6x+z) dz dx dy$$

$$\int_{-3}^3 \int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} \int_{\sqrt{x^2+y^2}}^{\sqrt{18-x^2-y^2}} (6x+z) dz dy dx$$

(b) Cylindrical coordinates

$$\operatorname{div} \vec{F} = 6r \cos \theta + z$$

$$\int_0^{2\pi} \int_0^3 \int_r^{\sqrt{18-r^2}} (6r \cos \theta + z) r dz dr d\theta$$

(c) Spherical coordinates

$$\operatorname{div} \vec{F} = 6\rho \sin \phi \cos \theta + \rho \cos \phi$$

$$\int_0^{2\pi} \int_0^{\pi/4} \int_0^{\sqrt{18}} (\operatorname{div} \vec{F}) \rho^2 \sin \phi d\rho d\phi d\theta$$