

# SOLUTION

MA 387

Exam I

February 26, 1975

Name \_\_\_\_\_

1. Prove that for all sets  $A$  and  $B$ ,  $A \subseteq B$  if and only if  $A \cup (B - A) = B$ .

proof:

→ Suppose  $A \subseteq B$  then

$$A \cup (B - A) = A \cup (B \cap \bar{A})$$

$$= (A \cup B) \cap (A \cup \bar{A})$$

$$= (A \cup B) \cap U$$

$$= A \cup B$$

$$= B \quad \text{because } A \subseteq B$$

← Suppose  $A \cup (B - A) = B$  and  $x \in A$

then since  $x \in A \cup (B - A)$ , it follows

that  $x \in B$ . Therefore,  $A \subseteq B$

2. Show that if  $F : A \rightarrow B$ ,  $G : B \rightarrow A$  and  $F \circ G = i_B$   
then  $F$  maps  $A$  onto  $B$  and  $G$  is 1-1.

(a)  $F$  maps  $A$  onto  $B$ .

Suppose  $y \in B$ . Then since  $F \circ G = i_B$ ,  
 $F(G(y)) = y$ . Since  $G : B \rightarrow A$ ,  $G(y) \in A$ .

Therefore  $x = G(y)$  is the element in  
 $A$  which is mapped onto  $y$  by the  
function  $F$ .

(b)  $G$  is 1-1.

Suppose  $x, y \in B$  and  $G(x) = G(y)$ .

Then  $F(G(x)) = F(G(y))$ . But, by hypothesis,

$F \circ G = i_B$ , so  $F(G(x)) = x$  and

$F(G(y)) = y$ . Consequently,  $x = y$  so

$G$  is 1-1

3. Calculate each of the following:

$$(a) \bigcap_{n \in \mathbb{Z}^+} [0, 1/n] = \{0\}$$

proof:

Clearly  $0 \in [0, 1/n]$  for every  $n > 0$ , so  $\{0\} \subseteq \bigcap_{n \in \mathbb{Z}^+} [0, 1/n]$

Conversely if  $x \in \bigcap_{n \in \mathbb{Z}^+} [0, \frac{1}{n}]$  then  $x \in [0, \frac{1}{n}]$

for every  $n \in \mathbb{Z}^+$ . This implies  $x \geq 0$ . If

$x > 0$  choose  $n \in \mathbb{Z}^+$  so that  $0 < \frac{1}{n} < x$ .

Then  $x \notin [0, \frac{1}{n}]$ . Therefore  $x \notin \bigcap_{n \in \mathbb{Z}^+} [0, \frac{1}{n}]$ .

$\therefore$  if  $x \in \bigcap_{n \in \mathbb{Z}^+} [0, \frac{1}{n}]$  then  $x = 0$ .

$$(b) \bigcup \{(a, b) \mid (a, b) \subseteq [1, 2]\} = (1, 2)$$

proof:

Suppose  $x \in (1, 2)$ . Choose  $a$  and  $b$  so that  $1 < a < x < b < 2$ . Then  $x \in (a, b)$ . Therefore,

$$x \in \bigcup \{(a, b) \mid (a, b) \subseteq [1, 2]\}$$

Conversely, suppose  $x \in$  Left side. Then there exist  $a$  &  $b$  such that  $1 < a < b < 2$  such that  $x \in (a, b)$ .  $\therefore 1 < x < 2$  so  $x \in (1, 2)$ .

4. (a) Prove that if  $R$  is a linear ordering relation then so is  $R^{-1}$ .

proof Suppose  $R$  is a linear ordering.

(a) Show  $R^{-1}$  is reflexive. Let  $x \in D_{R^{-1}} = D_R$ .

Then  $\langle x, x \rangle \in R$  because  $R$  is reflexive. Consequently,  $\langle x, x \rangle \in R^{-1}$  so  $R^{-1}$  is reflexive.

(b) Show  $R^{-1}$  is anti-symmetric. Suppose  $\langle x, y \rangle, \langle y, x \rangle \in R^{-1}$ . Then  $\langle y, x \rangle, \langle x, y \rangle \in R$ . Since  $R$  is anti-symmetric,  $x = y$ .

(c) Show  $R^{-1}$  is transitive. Suppose  $\langle x, y \rangle, \langle y, z \rangle \in R^{-1}$ . Then  $\langle y, x \rangle, \langle z, y \rangle \in R$ . Then since  $R$  is transitive,  $\langle z, x \rangle \in R$  which implies  $\langle x, z \rangle \in R^{-1}$ .

(d) Show  $R^{-1}$  is connected. Suppose  $x, y \in D_{R^{-1}} = D_R$ . Then since  $R$  is connected, either  $\langle x, y \rangle \in R$  or  $\langle y, x \rangle \in R$  or  $x = y$ . This implies, either  $\langle y, x \rangle \in R^{-1}$  or  $\langle x, y \rangle \in R^{-1}$ , so  $R^{-1}$  is connected.

- (b) Give an example of a relation  $R$  which is a well-ordering relation but  $R^{-1}$  is not.

Let  $R = \leq$  on  $\mathbb{Z}^+$  then  $R^{-1} = \geq$   
 $\langle \mathbb{Z}^+, \leq \rangle$  is well-ordered but  
 $\langle \mathbb{Z}^+, \geq \rangle$  is not

5. Find two sets  $X$  and  $Y$  such that  $X \approx Y$  but  $(X - Y) \neq (Y - X)$ .

let  $X = \mathbb{Z}^+ \cup \{0\}, Y = \mathbb{Z}^+$

Then  $X \approx Y$  by the mapping  $F$

where  $F(m) = m + 1$ . But  $X - Y = \{0\}$

and  $Y - X = \emptyset$  so

$$X - Y \neq Y - X.$$