

MATH 162 – SPRING 2010 – THIRD EXAM – APRIL 13, 2010  
VERSION 01  
MARK TEST NUMBER 01 ON YOUR SCANTRON

STUDENT NAME \_\_\_\_\_

STUDENT ID \_\_\_\_\_

RECITATION INSTRUCTOR \_\_\_\_\_

INSTRUCTOR \_\_\_\_\_

RECITATION TIME \_\_\_\_\_

INSTRUCTIONS

1. Fill in all the information requested above and the version number of the test on your scantron sheet.
2. This booklet contains 13 problems. Problem 1 is worth 4 points. The others are worth 8 points each. The maximum score is 100 points.
3. For each problem mark your answer on the scantron sheet and also circle it in this booklet.
4. Work only on the pages of this booklet.
5. Books, notes and calculators are not allowed.
6. At the end turn in your exam and scantron sheet to your recitation instructor.

Useful Formulas

$$\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}, \quad \sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n.$$

1)(4 points) For the series  $\sum_{n=1}^{\infty} (-1)^n n^2$ , the partial sum  $s_4$  equals

A) 2.

B) 10.

C) -10.

D) -2.

E) 30.

2)(8 points) Which of the following statements are true?

(I) If  $\lim_{n \rightarrow \infty} a_n = 0$ , then  $\sum_{n=1}^{\infty} a_n$  converges.

(II) If  $\sum_{n=1}^{\infty} |a_n|$  converges, then  $\sum_{n=1}^{\infty} a_n$  converges.

(III) If  $\sum_{n=1}^{\infty} \left| \frac{a_{n+1}}{a_n} \right|$  converges, then  $\sum_{n=1}^{\infty} a_n$  converges.

(IV) If  $0 \leq a_n \leq b_n$  and  $\sum_{n=1}^{\infty} b_n$  diverges, then  $\sum_{n=1}^{\infty} a_n$  diverges.

(V) If  $\lim_{n \rightarrow \infty} 5^n a_n = 2$ , then  $\sum_{n=1}^{\infty} a_n$  converges.

A)(I), (II) and (III) only.

B)(I), (II) and (IV) only.

C)(II), (IV) and (V) only.

D)(II), (III) and (V) only.

E)(II), (III) and (IV) only.

3)(8 points) Which of the following alternatives is true about the series  $\sum_{n=2}^{\infty} \frac{1}{n(\log n)^2}$ ?

- A) It converges by the comparison test with  $\sum_{n=1}^{\infty} \frac{1}{n}$ .
- B) It diverges by the comparison test with  $\sum_{n=1}^{\infty} \frac{1}{n}$ .
- C) It converges by the comparison test with  $\sum_{n=1}^{\infty} \frac{1}{n^2}$ .
- D) It diverges by the comparison test with  $\sum_{n=1}^{\infty} \frac{1}{n^2}$ .
- E) It converges by the integral test.

4)(8 points) Which of the following series diverge?

- (I)  $\sum_{n=1}^{\infty} \frac{n+1}{n^2}$
- (II)  $\sum_{n=1}^{\infty} \frac{n^2+n}{n^2-n}$
- (III)  $\sum_{n=2}^{\infty} \frac{1}{\ln n}$

A)(I) only.

B)(II) only.

C)(I) and (II) only.

D)(II) and (III) only.

E)All of them.

5)(8 points) Which statement is true about the following series?

$$(I) \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$$

$$(II) \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$$

$$(III) \sum_{n=1}^{\infty} (-1)^n \sqrt{n}$$

A) All are conditionally convergent.

B) All are divergent.

C) (I) is conditionally convergent; (II) is absolutely convergent.

D) (I) is absolutely convergent; (II) is conditionally convergent.

E) (I) and (II) are conditionally convergent; (III) is divergent.

6)(8 points) Let  $S = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(n+1)(n-1)}$ . Find the smallest integer  $N$  such that we

can be sure that  $|S_N - S| < \frac{1}{100}$ , where  $S_N = \sum_{n=1}^N \frac{(-1)^{n+1}}{(n+1)(n-1)}$

A) 8.

B) 9.

C) 10.

D) 11.

E) 12.

7)(8 points) The radius and interval of convergence of the power series  $\sum_{n=1}^{\infty} \frac{(-1)^n(x-2)^n}{(n+1)}$  satisfy

- A) The radius is equal to 1 and the interval is  $(-1, 1)$ .
- B) The radius is equal to 2 and the interval is  $(0, 4)$ .
- C) The radius is equal to 1 and the interval is  $(1, 3)$ .
- D) The radius is equal to 1 and the interval is  $(1, 3]$ .
- E) The radius is equal to 1 and the interval is  $[1, 3]$ .

8)(8 points) Which of the following is a power series representation of the function

$$f(x) = \frac{x-2}{x^2-4x+5}?$$

- A)  $\sum_{n=0}^{\infty} \frac{1}{n!}(x-2)^n.$
- B)  $\sum_{n=0}^{\infty} \frac{(-1)^n}{n}(x-2)^n.$
- C)  $\sum_{n=0}^{\infty} (x-2)^{n+1}.$
- D)  $\sum_{n=0}^{\infty} (-1)^n(x-2)^{n+1}.$
- E)  $\sum_{n=0}^{\infty} \frac{(-1)^n}{(n+1)}(x-2)^{n+1}.$

**9)(8 points)** The Maclaurin series of the function  $f(x) = \frac{1}{(4-x)^3}$  is

(Hint: Start with the power series of  $(4-x)^{-1}$  and differentiate it enough times.)

- A)  $\sum_{n=2}^{\infty} \frac{n(n-1)}{2(4^{n+1})} x^{n-2}$ .
- B)  $\sum_{n=2}^{\infty} \frac{n^2}{4^n} x^{n-2}$ .
- C)  $\sum_{n=2}^{\infty} \frac{(-1)^n n(n-1)}{4^n} x^{n-2}$ .
- D)  $\sum_{n=2}^{\infty} \frac{(-1)^n n(n-1)}{4^{n+2}} x^{n-2}$ .
- E)  $\sum_{n=2}^{\infty} \frac{(-1)^n n^2(n-1)}{2(4^n)} x^{n-2}$ .

**10)(8 points)** The Maclaurin series of  $f(x) = (\cos x)^2$  is equal to

(Hint: Use that  $(\cos x)^2 = \frac{1}{2}(1 + \cos 2x)$ .)

- A)  $\frac{1}{2} + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} x^n$ .
- B)  $\frac{1}{2} + \sum_{n=0}^{\infty} \frac{(-1)^n 4^n}{2(2n)!} x^{2n}$ .
- C)  $\frac{1}{2} + \sum_{n=0}^{\infty} \frac{(-1)^n 8^n}{4(n!)} x^{2n}$ .
- D)  $\frac{1}{2} + \sum_{n=0}^{\infty} \frac{(-1)^n 4^n}{2(2n)!} x^n$ .
- E)  $\frac{1}{2} + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} x^{4n}$ .

11)(8 points) Let  $f(x) = \sum_{n=0}^{\infty} \frac{2^n}{n!} (x-2)^n$ . We can say that the fifth derivative of  $f$  at the point 2 is equal to

A)  $f^{(5)}(2) = 10$ .

B)  $f^{(5)}(2) = 64$ .

C)  $f^{(5)}(2) = 32$ .

D)  $f^{(5)}(2) = 21$ .

E)  $f^{(5)}(2) = 100$ .

12)(8 points) If we use that  $\frac{1}{\sqrt{1-x}} = 1 + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2^n n!} x^n$ , and that

$\frac{d}{dx} \arcsin x = \frac{1}{\sqrt{1-x^2}}$ , we conclude that the Maclaurin series of  $\arcsin x$  is equal to

A)  $x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2^n (2n+1) n!} x^{2n+1}$ .

B)  $x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2^{n+3} (2n+1)!} x^{2n+1}$ .

C)  $x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2^n (2n+1) n!} x^n$ .

D)  $1 + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2^n n!} x^{2n+1}$ .

E)  $x + \sum_{n=1}^{\infty} \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2^n n!} x^{2n+3}$ .

**13)(8 points)** Let  $f(x)$  be a function defined on  $[1, \infty)$  such that  $f(x) > 1$  for all  $x$  and  $\lim_{x \rightarrow \infty} \frac{f(x)}{x} = 1$ . What can we say about the convergence of the series

$$S_1 = \sum_{n=1}^{\infty} \sin\left(\frac{1}{f(n)}\right) \text{ and } S_2 = \sum_{n=1}^{\infty} \sin\left(\frac{1}{f(n)^3}\right)?$$

- A)  $S_1$  and  $S_2$  diverge.
- B)  $S_1$  converges and  $S_2$  diverges.
- C)  $S_1$  diverges and  $S_2$  converges.
- D)  $S_1$  and  $S_2$  converge.
- E) Nothing can be said about the convergence of the series.