## Math 530

## Homework 3

1. Suppose that f(z) and g(z) are given by convergent power series  $\sum_{n=0}^{\infty} a_n z^n$  and  $\sum_{n=0}^{\infty} b_n z^n$ , respectively, where  $R_f > 0$  and  $R_g > 0$ . Prove that, if  $g(0) \neq 0$ , then f/g is analytic in a neighborhood of the origin and the power series for f(z)/g(z) is  $\sum_{n=0}^{\infty} c_n z^n$  where the  $c_n$ 's can be determined recursively via the formula,

$$a_n = \sum_{k=0}^n b_k c_{n-k}.$$
 (\*)

Is the radius of convergence of this series at least as big as the minimum of the radii of convergence for the series for f and g? Can the radius be larger than this?

- 2. Use formula (\*) of problem 1 and the complex power series for sine and cosine to find the first three nonzero terms in the power series expansion for  $\tan z = \sin z / \cos z$  about z = 0. What is the radius of convergence of the Taylor series for  $\tan z$  about z = 0?
- 3. (Recall that a *domain* is an open connected set). It can be shown that an analytic function f(z) on a domain Ω must be constant if A) f is *real valued* on Ω, or B) |f| is constant on Ω, or C) arg f is constant on Ω. Instead of giving separate proofs for A-C, do the following single problem that implies them all. Suppose a curve Γ in the complex plane is described as the level set of a function ρ:

$$\Gamma = \{ z = x + iy \in \mathbb{C} : \rho(x, y) = 0 \},\$$

where  $\rho$  is a real valued twice continuously differentiable function on  $\mathbb{R}^2$  and  $\nabla \rho$  is non-vanishing on  $\Gamma$ . Prove that if f(z) is an analytic function on a domain  $\Omega$  such that  $f(\Omega) \subset \Gamma$ , then f must be constant on  $\Omega$ .

- 4. We know that  $\exp(x+iy) = e^x(\cos y+i\sin y)$ . Find a similar formula for  $\sin(x+iy)$ .
- 5. Show that  $\int_{-\infty}^{\infty} e^{-t^2} \cos 2bt \, dt = \sqrt{\pi} e^{-b^2}$  by integrating  $e^{-z^2}$  around the rectangle with corners at  $\pm a$  and  $\pm a + ib$ . Let  $a \to \infty$ .