

Comb functions

Alexandre Eremenko* and Peter Yuditskii†

October 15, 2011

Dedicated to the memory of Franz Peherstorfer

Abstract

We discuss a class of regions and conformal mappings which are useful in several problems of approximation theory, harmonic analysis and spectral theory.

MSC: 30C20, 41A10, 47B36, 41A50. Keywords: conformal map, Green function, Martin function, uniform approximation, Jacobi matrices, Riesz bases, spectral theory.

1 Introduction

We begin with two classical problems which serve as motivation. Then in sections 2–5 we describe some classes of regions, corresponding conformal maps and entire and subharmonic functions. In sections 6–7 we discuss various problems where these classes appear.

1. Polynomials of least deviation from zero. Let $E \subset \mathbf{R}$ be a compact set on the real line, and P_n a polynomial with minimal sup-norm $L_n = \|P_n\|_E$ among all monic polynomials of degree n .

If $n < \text{card } E$, then P_n is unique and can be characterized by the following properties:

- (i) P_n is real, and all its zeros are real and simple,

*Supported by NSF grant DMS-1067886.

†Supported by the Austrian Science Fund FWF, project no: P22025-N18.

(ii) For every pair of adjacent zeros $x_1 < x_2$ there is a point $y \in (x_1, x_2) \cap E$ such that $|P_n(y)| = L_n$.

(iii) At the points $a_1 = \inf E$ and $a_2 = \sup E$, we have $|P_n(a_j)| = L_n$.

For a simple variational argument which proves (i)–(iii) see [3, 47]. These polynomials P_n can be represented in terms of special conformal maps.

Let m, k be integers, $k - m = n$, and let D be a region obtained from the half-strip

$$\{z = x + iy : \pi m < x < \pi k, y > 0\}$$

by removing vertical intervals $\{\pi j + it : 0 \leq t \leq h'_j\}$, $m < j < k$, where $h'_j \geq 0$, see Fig. 1 (right).




Fig. 1. Comb regions of V (left) and MO (right) types.

Let θ be a conformal map from the upper half-plane \mathbf{H} to D , such that $\theta(\infty) = \infty$, $\theta(a_1) = \pi m$, $\theta(a_2) = \pi k$. Then $z \mapsto \cos \theta(z)$ is an analytic function in the upper half-plane, which is real on the real line. So it extends to an entire function, and the behavior at ∞ shows that this entire function is a polynomial of degree n . Choose L so that the polynomial $P = L \cos \theta$ is monic. It is easy to check that our extremal polynomials satisfying (i)–(iii) are of this form, with an appropriate choice of parameters $h'_j \geq 0$.

The set E is contained in $E' := \theta^{-1}([\pi m, \pi k])$. This set E' is the maximal extension of E , for which the extremal polynomial is the same as the one for E . Critical points of P_n are preimages of the tips of the slits under θ , and critical values are $\pm \cosh h'_j$. The θ -preimages of the points πj , $m < j < k$ are solutions of $P_n(z) = \pm L$, and all these solutions are real.

For example, if $E = [-1, 1]$ we take all $h'_j = 0$, and P_n is the n -th Chebyshev polynomial. If E consists of two intervals symmetric with respect to 0, and n is even, we take all $h'_k = 0$, except one, $h'_{(m+k)/2} > 0$.

On polynomials of least deviation from 0 on several intervals we refer to [2, 3], [4, vol. 1] and the survey [47], where the representation $P_n = L \cos \theta$ is used systematically.

2. Spectra of periodic Jacobi matrices. Consider a doubly infinite, periodic Jacobi matrix

$$J = \begin{pmatrix} \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ p_{-1} & q_{-1} & p_0 & 0 & 0 & 0 \\ 0 & p_0 & q_0 & p_1 & 0 & 0 \\ 0 & 0 & p_1 & q_1 & p_2 & 0 \\ 0 & 0 & 0 & p_2 & q_2 & p_3 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \end{pmatrix}$$

which is constructed of two periodic sequences of period n , where q_j are real, and $p_j > 0$. This matrix defines a bounded self-adjoint operator on ℓ_2 , and we wish to describe its spectrum [18, 33, 45].

To do this we consider a generalized eigenvector $u \in \ell_\infty$ which satisfies

$$Ju = zu, \quad z \in \mathbf{C}.$$

For fixed z , this can be rewritten as a recurrent relation on the coordinates of u :

$$p_{j+1}u_j + (q_{j+1} - z)u_{j+1} + p_{j+2}u_{j+2} = 0,$$

which we rewrite in the matrix form as

$$\begin{pmatrix} u_{j+1} \\ p_{j+2}u_{j+2} \end{pmatrix} = \begin{pmatrix} 0 & 1/p_{j+1} \\ -p_{j+1} & (z - q_{j+1})/p_{j+1} \end{pmatrix} \begin{pmatrix} u_j \\ p_{j+1}u_{j+1} \end{pmatrix}.$$

Thus

$$\begin{pmatrix} u_n \\ p_{n+1}u_{n+1} \end{pmatrix} = T_n(z) \begin{pmatrix} u_0 \\ p_1u_1 \end{pmatrix},$$

where $T_n(z)$ is a polynomial matrix with determinant 1, which is called the transfer-matrix. To have a bounded generalized eigenvector u , both eigenvalues of T_n must have absolute value 1. This happens if and only if

$$|P_n(z)| := |\operatorname{tr} T_n(z)|/2 \leq 1.$$

As P_n is a real polynomial, the spectrum is the preimage of the interval $[-1, 1]$. As our matrix J is symmetric, the spectrum must be real, this is the

same as the condition that all solutions of the equations $P_n(z) = \pm 1$ are real, so we obtain a polynomial of the same kind as in Example 1. For every real polynomial with this property, there exists a periodic Jacobi matrix whose spectrum is $P_n^{-1}([-1, 1])$, and all matrices J with a given spectrum can be explicitly described [34, 45]. Our polynomial has a representation $P_n = \cos \theta$, where θ is a conformal map of the upper half-plane onto a comb region D as in Example 1. We obtain the result that the spectrum of a periodic Jacobi matrix consists of the intervals – preimage of the real line under a conformal map θ .

We can prescribe an arbitrary sequence h'_j , $1 \leq j \leq n - 1$, construct a conformal map $\theta : \mathbf{H} \rightarrow D$, where D is the region shown in Fig. 1 (right), and the polynomial $P = \cos \theta$ will have critical values $(-1)^j \cosh h'_j$ and all solutions of $P(z) = \pm 1$ will be real. Such polynomial P is defined by its critical values of alternating sign up to a change of the independent variable $z \mapsto az + b$, $a > 0, b \in \mathbf{R}$. Later we will show that any real polynomial with arbitrary real critical points is defined by its critical values up to a change of the independent variable $z \mapsto az + b$, $a > 0, b \in \mathbf{R}$.

We conclude this introduction with several historical remarks. After the works of Sergei Bernstein, in the 1950s and 1960s the development of the Chebyshev approximation was clearly dominated by the study of *entire functions* of least deviation from zero, see, for example [1, p. 320-363]. In particular, Boas and Schaeffer [9] (see also [36]) proved under very general assumptions that the extremal function can be expressed in terms of the hyperelliptic integral

$$f(z) = L \sin \left\{ \int \frac{p(z)}{\sqrt{q(z)}} dz \right\}.$$

The new wave of interest in the comb polynomials is most likely related to the theory of integrable systems, see e.g. [7], and the application of the iteration theory to the spectral theory of almost periodic operators with Cantor type spectrum, see e.g. [20]. Franz Peherstorfer explicitly formulated both properties of the comb polynomials (as polynomials of the least deviation from zero and in connection with periodic Jacobi matrices) in [40], see also [41, 42, 47].

An attractive aspect of this circle of questions is the numerous connections with diverse classical problems in analysis: investigations of Abel on

expressing elliptic and hyperelliptic integrals in elementary functions; continued fractions and Pell's equations; factorization of functions on Riemann surfaces; subharmonic majorants; and so on. For modern surveys of these questions we refer to [47, 31]. For other applications of comb functions we mention [10, 11].

In parallel with our paper Injo Hur and Christian Remling presented a special paper [21] on applications of comb functions in spectral theory of ergodic Jacobi matrices and we refer the reader to it in particular, for further references related to this important direction.

2 Comb representation of LP entire functions

In both examples in the Introduction, the class of real polynomials P such that all solutions of $P(z) = \pm 1$ are real appears. Evidently, all zeros of such polynomials must be real and simple. Here we discuss a representation of polynomials with real zeros, not necessarily simple, using conformal mappings and generalization of this representation to a class of entire functions.

Let P be a non-constant real polynomial of degree n with all zeros real. Let $\varphi = \log P$ be a branch of the logarithm in the upper half-plane \mathbf{H} . Then

$$-\varphi' = -\frac{P'}{P} = -\sum_{n=j}^n \frac{1}{z - z_j}$$

is an analytic function in \mathbf{H} with positive imaginary part.

Lemma 1. *An analytic function ψ in \mathbf{H} whose derivative has positive imaginary part is univalent.*

Proof. Suppose that $\psi(z_1) = \psi(z_2)$, $z_j \in \mathbf{H}$, $z_1 \neq z_2$. Then

$$0 = \frac{\psi(z_1) - \psi(z_2)}{z_1 - z_2} = \int_0^1 \psi'(z_2 + t(z_1 - z_2)) dt,$$

but the last integral has positive imaginary part and thus cannot be 0.

It is easy to describe the image $\varphi(\mathbf{H})$. By Rolle's theorem, all zeros of P' are real and we arrange them in a sequence $x_1 \leq \dots \leq x_{n-1}$ where each zero is repeated according to its multiplicity. Let $c_j = P(x_j)$ be the *critical sequence*

of P . Then the region $D = \varphi(\mathbf{H})$ is obtained from a strip by removing $n - 1$ rays:

$$D = \{x + iy : \pi m < y < \pi k\} \setminus \bigcup_{m < j < k} \{x + i\pi j : -\infty < x \leq h_j\}. \quad (1)$$

Here $m - k = n$, and $h_j = \log |c_{k-j}| \geq -\infty$. Thus

$$P = \exp \varphi, \quad (2)$$

where φ is a conformal map of the upper half-plane onto a region D we just described (see Fig. 1, left). Such regions will be called polynomial V -combs. Letter V in this notation is used because this representation was introduced by E. B. Vinberg in [49].

Now suppose that an arbitrary finite sequence $h_j \in [-\infty, \infty)$, $m < j < k$ is given. Consider a V -comb D corresponding to this sequence, and a conformal map $\varphi : \mathbf{H} \rightarrow D$. Using the same argument with reflection as in Example 1 in the Introduction, it is easy to see that $\exp \varphi$ is a real polynomial of degree n with critical values $(-1)^j e^{h_j}$.

We obtain

Theorem 1. *For every finite sequence c_1, \dots, c_{n-1} with the property*

$$c_{j+1}c_j \leq 0, \quad (3)$$

there exists a real polynomial with real zeros for which this sequence is the critical sequence. Such polynomial is defined by its critical sequence up to a real affine change $z \mapsto az + b$, $a > 0$ of the independent variable.

Now we extend this result to entire functions. Recall that an entire function belongs to the class LP (Laguerre-Pólya) if it is a limit of real polynomials with all zeros real. For more information on the LP -class and its applications we refer to [25]. Consider the following class of regions. Begin with $\{x + iy : \pi m < y < \pi k\}$, where m, k are integers or $\pm\infty$, $-\infty \leq m < k \leq \infty$ and remove from this region the rays of the form

$$\{x + i\pi j : x \leq h_j\}, \quad m < j < k, \quad (4)$$

where $h_j \in [-\infty, \infty)$. A region of this form is called a V -comb corresponding to a sequence (h_j) , $h_j \in [-\infty, \infty)$. The sequence can be finite, or infinite in one direction, or infinite in both directions.

Theorem 2. *The following statements are equivalent:*

- (i) $f \in LP$,
- (ii) $f = \exp \varphi$, where $\varphi : \mathbf{H} \rightarrow D$ is a conformal map onto a V -comb,
- (iii) $f(z) = z^q e^{-az^2+bz} \prod_{j=1}^{\infty} (1 - z/z_j) e^{z/z_j}$, where $z_j, a, b \in \mathbf{R}$;

$$\sum_j |z_j|^{-2} < \infty,$$

$a \geq 0$ and $q \geq 0$ is an integer.

It follows that there exists a function $f \in LP$ with a prescribed sequence of critical values c_j satisfying (3), and prescribed limits $\lim_{x \rightarrow \pm\infty} f(x) \in \{0, \infty\}$ (asymptotic values). Such function is defined by its critical sequence and asymptotic values up to an increasing real affine change of the independent variable.

Here are some examples of comb representations (ii).

- $f(z) = z + b, b \in \mathbf{R}$. There are no critical values, asymptotic values are $\pm\infty$. D is a strip $\{0 < |\operatorname{Im} z| < \pi\}$.
- $f(z) = \cos z$. The critical sequence is $(-1)^j$, infinite in both directions, there are no asymptotic values. D is the plane, cut along the rays $\{x + i\pi j : -\infty < x \leq 0\}$.
- $f(z) = \exp(-z^2)$. D is the plane cut along the negative ray.
- $f(z) = 1/\Gamma(z)$. D is the plane cut along the rays $\{x + i\pi j : -\infty < x \leq \log |c_j|\}$, $j < 0$, where c_j are the critical values of the Γ -function, there is an asymptotic value $0 = \lim_{x \rightarrow +\infty} 1/\Gamma(x)$.
- f a polynomial of degree n . D is obtained from the strip $\{x + iy : 0 < y < \pi n\}$ by removing the rays (4), where $h_j = \log |c_j|$, and c_j are the critical values of f .

An important subclass of LP is defined by the condition that $h_j \geq 0$, $m < j < k$, and whenever the sequence of critical points is bounded from below (or above, or from both sides), then the corresponding asymptotic value is ∞ . This subclass of LP will be called the MO -class. It was used for the first time in spectral theory in [35]. Functions of MO -class have another

representation in terms of conformal mappings. Consider a region D of the form

$$\{x + iy : y > 0, \pi m < x < \pi k\} \setminus \bigcup_{j=m+1}^{k-1} \{\pi k + iy : 0 \leq y \leq h'_j\}, \quad (5)$$

where $-\infty \leq m < k \leq \infty$, and $h'_j \geq 0$, see Fig. 1 (right). Such regions will be called *MO-combs*. Let $\theta : \mathbf{H} \rightarrow D$ be a conformal map $\theta(\infty) = \infty$. Then

$$f = \cos \theta \quad (6)$$

is a function of the class *MO* with critical values $(-1)^j \cosh h'_j$. Every function of *MO* class can be represented in this way, and the function is defined by its critical sequence up to a real affine change of the independent variable. We have the following important characterization of the *MO* class [35]:

Theorem 3. *For a real entire function f , the equation $f^2(z) - 1$ has only real roots if and only if $f \in MO$.*

Such functions occur in the situation similar to the Examples 2 and 1 in the Introduction: they describe the spectra of periodic canonical systems [23, 12] and entire functions of smallest deviation from zero on closed subsets of the real axis [47].

3 MacLane's theorem

In this section we give a geometric characterization of integrals of *LP* functions. Roughly speaking, we will show that critical values of these integrals can be arbitrarily prescribed, subject to the evident restriction (7). Notice that differentiation maps *LP* into itself, so the class of integrals of *LP*-functions contains *LP*.

We follow the exposition in [49] with some corrections and simplifications, see also [14] on related questions. Let f be a real entire function with all critical points real.

Consider the preimage $f^{-1}(\mathbf{R})$. It contains the real line, and it is a smooth curve in a neighborhood of any point which is not a critical point. At a critical point of order n it looks like the preimage of the real line under z^{n+1} .

MacLane's class consists of real entire functions for which the preimage of the real line looks like one of the pictures in Fig. 2,




Fig. 2. Fish-bones.

up to an orientation preserving homeomorphism of the plane, commuting with the complex conjugation. We call this picture a fish-bone.

There are several cases. In the simplest case, the sequence of critical points $\dots \leq x_j \leq x_{j+1} \leq \dots$ is unbounded from below and from above. Each critical point is repeated in this sequence according to its multiplicity. Preimage of the real line consists of the real line itself, crossed by infinitely many simple curves, each curve is symmetric with respect to the real line. The crossing points are mapped onto the critical values $c_j = f(x_j)$. Several “vertical” lines cross the real line at a multiple critical point. The complement to the union of curves in Fig. 2, consists of simply connected regions, each of them is mapped conformally onto the upper or lower half-plane.

The sequence of critical points can be bounded from above or from below or both. Suppose that it is bounded from below, and enumerate the sequence as $x_1 \leq x_2 \leq \dots$. Then the left end of the fish-bone can be of two types. For the first type, shown in Fig 2 (left), the full preimage of the real line is connected. We have two large complementary regions adjacent along a negative ray, each of them is mapped by f homeomorphically onto the upper or lower half-plane. It is easy to see that in this case we have $f(x) \rightarrow \infty$ as $x \rightarrow -\infty$. We set $c_0 = \infty$ and extend our critical sequence (c_j) by adding this term to it.

The second type of the end is shown in Fig. 2 (right). In this case, the preimage of the real has infinitely many components. In addition to one component of $f^{-1}(\mathbf{R})$, as above, there are infinitely many simple curves tending to infinity at both ends. Strip-like regions between these curves are mapped homeomorphically onto the upper or lower half-plane. In this case $c_0 = \lim_{x \rightarrow -\infty} f(x) \neq \infty$, and we extend our critical sequence by c_0 .

Similar situations may occur on the right end when the sequence of critical points is bounded from above.

In all cases, the fish-bone is completely determined by the *augmented critical sequence* (c_j) . We use the following notation: if the sequence of critical points is unbounded from above and below, then $-\infty < j < +\infty$. In all other cases, the critical values are c_j ; $m < j < k$, where $-\infty \leq m < k \leq +\infty$, and if m or k or both are finite, we add to our sequence the term c_m or c_k or both, which are the limits of $f(x)$ as $x \rightarrow -\infty$ or $x \rightarrow +\infty$.

The augmented critical sequence satisfies the following condition

$$(c_{j+1} - c_j)(c_j - c_{j-1}) \leq 0. \tag{7}$$

All c_j are real, except possibly the first and/or the last term which can be $\pm\infty$.

We call such sequences “up-down sequences”. If the sequence of critical points is unbounded from below and from above, then the sequences x_j and c_j are defined for a given f up to a shift of the subscript.

MacLane’s theorem [30] *For every up-down sequence, finite or infinite in one or both directions, there exists a function $f \in M$ for which this sequence is the critical sequence. Any two functions corresponding to the same sequence are related by $f_1(z) = f_2(az + b)$ with $a > 0, b \in \mathbf{R}$.*

In other words, one can prescribe a piecewise-monotone graph on the real line, and after a strictly increasing continuous change of the independent variable, this will be the graph of an entire function of MacLane’s class, which is essentially unique.

Uniqueness statement in MacLane’s theorem is easy. Suppose that f_1 and f_2 are two functions of MacLane class with the same augmented critical sequence. Then it is easy to construct a homeomorphism ϕ of the plane such that $f_1 = f_2 \circ \phi$. Then ϕ must be conformal and commute with complex conjugation, so ϕ must be a real affine map.

Class LP is contained in the MacLane class. It corresponds to the case when the critical sequence satisfies the condition (3) and in addition, the first and last terms of the sequence c_j , if present, are 0 or ∞ . It is clear that (3) is stronger than (7).

We proved this special case of MacLane’s theorem in the previous section. Now we give the proof of MacLane’s theorem in full generality.

First we recover the fish-bone from the given sequence (c_j) as explained

above. Then we construct a continuous map $F : \mathbf{C}_z \rightarrow \mathbf{C}_w$ as follows. We map each interval $[x_j, x_{j+1}] \in \mathbf{R}$ linearly onto the interval $[c_j, c_{j+1}]$. Then we map each infinite ray of the fish-bone onto a corresponding ray of the real line, linearly with respect to length. The curves on the left of Fig. 2 (right) are mapped on the rays $[c_0, \infty)$. Then we extend our map to the components of the complement of the fish-bone, so that each component is mapped on the upper or lower half-plane homeomorphically.

The resulting continuous map F is a local homeomorphism everywhere except the points x_j where it is ramified. There is unique conformal structure ρ in the plane \mathbf{C}_z which makes this map holomorphic. By the uniformization theorem, the simply connected Riemann surface (\mathbf{C}, ρ) is conformally equivalent to a disc $|z| < R$, where $R \leq \infty$. This means that there exists a homeomorphism $\phi : \{z : |z| < R\} \rightarrow \mathbf{C}$ such that $F \circ \phi$ is a holomorphic function. As all our construction can be performed symmetrically with respect to the real line, F is a real function. It remains to prove that $R = \infty$.

If the sequence (x_j) is finite, and both asymptotic values are ∞ , our map extends to a continuous map of the Riemann sphere $\overline{\mathbf{C}}_z \rightarrow \overline{\mathbf{C}}_w$ by putting $F(\infty) = \infty$. So the Riemann surface $(\overline{\mathbf{C}}, \rho)$ must be conformally equivalent to the sphere, and we obtain that $R = \infty$. In this case $f = F \circ \phi$ is evidently a polynomial.

If the sequence (x_j) is infinite in both directions, we consider truncated sequences $(c_j)_{j=-n}^n$, augmented by asymptotic values ∞ on both sides, and the corresponding fish-bones and maps F_n as above. By the previous argument we have homeomorphisms ϕ_n and polynomials $f_n = F_n \circ \phi_n$.

We can always arrange that $x_1 < x_2$, $0 \in (x_1, x_2)$, and $F_n(0) = a \in (c_1, c_2)$, where a is independent of n . Then we choose ϕ_n so that $\phi_n(0) = 0$, and

$$f'_n(0) = 1. \tag{8}$$

Then f_n maps univalently some disc $\{z : |z| < r\}$ onto a region G_n which contains a disc $\{w : |w| < \epsilon\}$ and is contained in a disc $\{w : |w| < \delta\}$ with some $r > 0, \epsilon > 0, \delta > 0$ which are independent of n . This follows from the Schwarz lemma applied to f_n and f_n^{-1} in a neighborhood of 0. We conclude that (f_n) is a normal family in $\{z : |z| < r\}$ and the limit functions are non-constant.

Now we use the following lemma [25].

Lemma 2. *Let g_n be a sequence of real polynomials whose all zeros are real, and suppose that $g_n \rightarrow g \not\equiv 0$ uniformly in some neighborhood of 0. Then g*

is entire, and $g_n \rightarrow g$ uniformly on compact subsets of \mathbf{C} .

Proof. By a shift of the independent variable we may assume that $g(0) \neq 0$. Then $g_n(0) \neq 0$ for large n . We have

$$-\left(\frac{g'_n}{g_n}\right)'(0) = \sum_k \frac{1}{z_{n,k}^2},$$

where $z_{n,k}$ are zeros of g_n . The left hand side is bounded by a constant independent of n , while all summands in the right hand side are positive. So for every interval I on the real line there exists a constant $c(I)$ independent of n such that the g_n have at most $c(I)$ roots on I . Thus from every sequence of g_n one can choose a subsequence such that the zero-sets of polynomials of this subsequence tend to a limit set which has no accumulation points in \mathbf{C} . So our subsequence converges to an entire function. Evidently this entire function is an analytic continuation of g , and the statement of the lemma follows.

We apply this lemma to the sequence (f'_n) and conclude that f is entire, that is $R = \infty$, as advertised.

Now we describe the necessary modifications of this proof for the case that the sequence of critical points is bounded from below (the case of semi-infinite sequence bounded from above is treated similarly). If the asymptotic value $c_0 = \infty$, no modification is needed. If $c_0 \neq \infty$, we may assume without loss of generality that $c_0 = 0$, by adding a real constant to all functions f, F, f_n . Then we approximate our critical sequence $c_0, c_1 \dots$ by the finite sequences $c_0, c_0, \dots, c_0, c_1 \dots, c_n$, where $c_0 = 0$ is repeated n times. The corresponding fish-bone is shown in Fig. 3, where β is the additional zero of multiplicity n . As $n \rightarrow \infty$, $\beta \rightarrow -\infty$. The rest of the argument goes without change.




Fig. 3. Approximation of a fish-bone by polynomial ones.

Finally we consider the case when there are finitely many critical points and two different asymptotic values. In this case, no approximation argument is needed, and $f(z) = \int_{-\infty}^z P(\zeta) \exp(-a\zeta^2 + b\zeta) d\zeta$, where P is a real polynomial with all zeros real and $a \geq 0$ and $b \in \mathbf{R}$.

4 Representation of Green's and Martin's functions

Here we discuss the relation between comb regions and Green and Martin functions of complements of closed sets on the real line.

Let $E \subset \mathbf{R}$ be a compact set of positive capacity. Then there exists the Green function G of $\Omega = \overline{\mathbf{C}} \setminus E$ with pole at ∞ . We have

$$G(z) = \int_E \log |z - t| d\mu(t) + \gamma(E), \quad (9)$$

where μ is a probability measure on E which is called the equilibrium measure, and γ the Robin constant of E . Function G is positive and harmonic in $\mathbf{C} \setminus E$, and has boundary values 0 a. e. with respect to μ . We have

$$G(z) = \log |z| + \gamma + o(1), \quad z \rightarrow \infty. \quad (10)$$

These properties characterize G and μ [24].

There exists an analytic function $\phi : \mathbf{H} \rightarrow \mathbf{H}$, such that $G = \text{Im } \phi$. It is called the complex Green function. Since the derivative

$$\phi' = \frac{d}{dz} \left(i \int_E \log(z - t) d\mu(t) \right) = i \int_E \frac{d\mu(t)}{z - t}$$

has positive real part in \mathbf{H} , we conclude from Lemma 1 that ϕ is univalent. Let $D = \phi(\mathbf{H})$. This region D has the following characteristic properties:

- (i) D is contained in a vertical half-strip $\{x + iy : a < x < b, y > 0\}$ with $b - a = \pi$, and contains a half-strip $\{x + iy : a < x < b, y > K\}$ with some $K > 0$.
- (ii) For every $z \in D$, the vertical ray $\ell_z = \{z + it : t \geq 0\}$ is contained in D .

(iii) For almost every $x \in (a, b)$, the ray $\{x + iy : y > 0\}$ is contained in D .

These properties can be restated shortly as follows:

$$D = \{x + iy : a < x < b, y > h(x)\}, \quad (11)$$

where h is as non-negative upper semi-continuous function bounded from above and equal to 0 a. e.

We sketch a proof of (i)–(iii). Function G given by (9) is upper semi-continuous, so it must be continuous at every point where $G(z) = 0$. If $h(x) = 0$ for some $x \in (a, b)$, then for the similar reason, h is continuous at x , so ∂D is locally connected at x . It follows that $x = \phi(x')$ for some $x' \in \mathbf{R}$, and ϕ is continuous at x' . In other words, existence of a radial limit $\phi(x')$, such that $\text{Im } \phi(x') = 0$ implies continuity of ϕ and G at the point x'

$$\text{Re } \phi(z) = - \int_E \arg(z - t) d\mu(t) = y \int_E \frac{\mu(t) dt}{(x - t)^2 + y^2}, \quad (12)$$

where $z = x + iy$ and $\mu(t) = \mu((-\infty, t])$ is the distribution function. As μ has no atoms, $t \mapsto \mu(t)$ is continuous. So $\text{Re } \phi$ is continuous in $\overline{\mathbf{H}}$. The first statement of (i) follows because $0 \leq \mu(t) \leq 1$, and the second because $G(z) = \text{Im } \phi$ is bounded on any compact set in \mathbf{C} .

To prove (ii), let α be a tangent vector to the ray ℓ_z , so $\alpha = i$. Then $\beta = (\phi^{-1})'\alpha$ will be the tangent vector to the ϕ -preimage of this ray, and we have seen that $\arg(\phi^{-1})' \in (-\pi/2, \pi/2)$. So β is in the upper half-plane thus the preimage of ℓ_z can never hit the real line, and an analytic continuation of ϕ^{-1} is possible along the whole ray ℓ_z .

To prove (iii), we use (12) again. As $\mu(t)$ is continuous, $\text{Re } \phi$ is continuous in $\overline{\mathbf{H}}$. Moreover,

$$\text{Re } \phi(\beta) - \text{Re } \phi(\alpha) = \mu(\beta) - \mu(\alpha), \quad \alpha < \beta.$$

This means that measure μ on E corresponds to the Lebesgue measure on base of the comb (a, b) . Furthermore, if for some $x \in E$ we have $G(x) = 0$ then $h(\text{Re } \phi(x)) = 0$. Thus $h = 0$ almost everywhere with respect to the Lebesgue measure on (a, b) . This proves (iii),

Now we show that for every D satisfying (i)–(iii), the conformal map $\phi : \mathbf{H} \rightarrow D$ is related with the Green function G of some closed set E by the formula $G = \text{Im } \phi$.

Imaginary part $v = \text{Im } \phi$ is a positive harmonic function in the upper half-plane. We extend it to the lower half-plane by symmetry, $v(\bar{z}) = v(z)$, and to the real line by upper semicontinuity: $v(x) = \limsup_{z \rightarrow x} v(z)$. In view of (i), ∂D has a rectilinear part near infinity, the extended function v is harmonic in a punctured neighborhood of ∞ and has asymptotics of the form

$$v(z) = \log |z| + \text{const} + o(1), \quad z \rightarrow \infty.$$

Let us prove that v is subharmonic in the whole plane, and has a representation (9) with some probability measure μ with compact support on the real line.

Let $\{h_k\}$ be a dense set on ∂D . Let D_n be the region obtained from the half-strip $\{x + iy : a < x < b, y > 0\}$ by removing the vertical segments $\{\text{Re } h_k + iy, 0 < y \leq \text{Im } h_k\}$. Then $D_1 \supset D_2 \supset \dots \rightarrow D$. Let ϕ_n be conformal maps of \mathbf{H} onto D_n , normalized by $\phi_n(0) = a$, $\phi_n(1) = b$, $\phi_n(\infty) = \infty$. Then it is easy to check that $\text{Im } \phi_n$ is the Green function of some set $E_n \subset [0, 1]$ consisting of finitely many closed intervals. So

$$\text{Im } \phi_n(z) = \int \log |z - t| d\mu_n(t) + \gamma_n,$$

with some probability measures μ_n on $[0, 1]$ and some constants γ_n . We can choose a subsequence such that $\mu_n \rightarrow \mu$ weakly, where μ is a probability measure on $[0, 1]$, and it is easy to check that (9) holds with some γ . Thus v is subharmonic in the plane. Since $v \geq 0$, the measure μ has no atoms. It remains to prove that $v(x) = 0$ a. e. with respect to μ . This follows from the property (iii) of the region D . Indeed, let $x \in (a, b)$ be a point such that the vertical ray ℓ_x is in D , except the endpoint x . By a well-known argument, the curve $\phi(\ell_x)$ has an endpoint at some $x' \in (0, 1)$, and the angular limit of $v = \text{Im } \phi$ is zero at this point x' . By the remark above, $v(x') = 0$. We define E as the closed support of μ . Then $v = \text{Im } \phi$ is positive and harmonic outside E and $v(x) = 0$ μ -almost everywhere, so v is the Green function of E .

Our construction of D from E defines D up to a shift by a real number. The inverse construction defines E up to a real affine transformation, and changing E by a set of zero capacity.

Now we give a similar representation of Martin functions. Let $E \subset \mathbf{R}$ be an unbounded closed set of positive capacity. Let U be the cone of positive harmonic functions in $\mathbf{C} \setminus E$, and $U_s \subset U$ the cone of symmetric positive harmonic functions, $v(z) = v(\bar{z})$. Martin's functions are minimal elements of

U , that is functions $v \in U$ with the property

$$u \in U, \quad u \leq v \quad \text{implies} \quad u = cv,$$

where $c > 0$ is a constant. Similarly we define *symmetric Martin functions* using U_s instead of U . Martin functions always exist and form a convex cone. If v is a Martin function, then $u(z) + u(\bar{z})$ is a symmetric Martin function, so symmetric Martin functions also exist and form a convex cone.

Let v be a symmetric Martin function, and let ϕ be an analytic function in \mathbf{H} so that $v(z) = \text{Im } \phi(z)$, $z \in \mathbf{H}$. Then $\phi : \mathbf{H} \rightarrow D$ is a conformal map onto a region $D \subset \mathbf{H}$. This is proved in the same way as for Green's functions. Regions D arising from symmetric Martin functions are characterized by the properties (ii), (iii) above and the *negation* of the property (i): either $a = -\infty$ or $b = +\infty$, or h is unbounded in (11). Levin introduced the following classification of regions D :

Class A: $a = -\infty$ and $b = \infty$.

Class B: one of the numbers a, b is finite, another infinite.

Class C: both a and b are finite.

Kesarev [19] gave a geometric criterion in terms of the set E which distinguish the cases A,B and C.

Notice that function ϕ maps h into \mathbf{H} , so the angular derivative of ϕ at infinity exists, that is

$$\phi(z) = cz + o(z), \quad z \rightarrow \infty \quad \text{in any Stolz angle,}$$

where $c \geq 0$. One can derive from this that every Martin function satisfies

$$B(r, v) := \max_{|z|=r} v(z) = O(r), \quad r \rightarrow \infty.$$

This implies that the cone of Martin functions has dimension at most 2, [22, 29, 12], and the cone of symmetric Martin functions is always one-dimensional.

Dimension of the cone of Martin's functions is an important characteristic of the set E , see [8, 29]. One can show that the cone of Martin functions is two-dimensional if and only if

$$\limsup_{r \rightarrow \infty} B(r, v)/r > 0.$$

A geometric criterion which tells in terms of the set E the dimension of the cone of Martin's functions is given by Andrievskii [6].

Now we impose various conditions on E and find their exact counterparts in terms of E and μ .

The first important condition is that the set E is regular in the sense of potential theory [24]. In this case Green's and Martin's functions are continuous in \mathbf{C} . For the region D this is equivalent to the local connectedness of ∂D in the case of Green's function, and local connectedness of the part $\partial D \setminus X$, where X is the union of the vertical rays on ∂D , if these rays are present. In terms of function h in (11), local connectedness is equivalent in the case of Green's function to the condition that the set $X = \{x : h(x) > 0\}$ is at most countable, and the sets $X_\epsilon = \{x : h(x) > \epsilon\}$ are finite for every $\epsilon > 0$, that is D is obtained from \mathbf{H} by making countably many cuts, and the length of a cut tends to 0.

In the case of Martin's function, local connectedness of D means that the sets X_ϵ can only accumulate to a or b .

Next we discuss the condition on D which corresponds to absolute continuity of μ . We thank Misha Sodin who passed to us the contents of his conversation with Ch. Pommerenke on this subject.

To state the result we first recall McMillan's sector theorem [32], [43, Thm. 6.24]. Let f be a conformal map from \mathbf{H} to a region G . Let $\text{sect}(f)$ be the set of points $x \in \mathbf{R}$ such that the non tangential limit $f(x)$ exists and $f(x)$ is the vertex of an angular sector in G .

McMillan Sector Theorem [43, Theorem 6.24, p.146]. *Assume that $A \subset \text{sect}(f)$. Then*

$$|A| = 0 \quad \text{if and only if} \quad |f(A)| = 0. \quad (13)$$

We say that the sector condition holds in the comb region D if the function

$$H(x) = \sup_{y \in (a,b)} \frac{h(y)}{|y-x|} \quad (14)$$

is finite for almost all $x \in (a, b)$. Geometrically it means that for almost all x in the base of the comb there exists a Stolz angle with the vertex in x .

Theorem 4. *Region D satisfies the sector condition if and only if μ is absolutely continuous with respect to the Lebesgue measure on \mathbf{R} .*

Proof. Recall that the Lebesgue measure on the base of the comb corresponds to the harmonic measure μ on E .

Assume that the sector condition holds. This means that a Borel support of the harmonic measure μ is contained in $\text{sect}(\phi)$. Let A be a Borel support of the singular component of μ . By the definition $|A| = 0$. Thus, by McMillan's theorem $\mu(A) = 0$, thus μ is absolutely continuous.

Conversely, assume that the harmonic measure is absolutely continuous. Recall that ϕ' has positive imaginary part, and therefore possesses non-tangential limits for almost all x with respect to the Lebesgue measure. Therefore the limit exists for almost all x with respect to the harmonic measure as well.

Example 1. There exist irregular regions with absolutely continuous measures μ . Indeed, let C be the standard Cantor set in $[a, b]$. Let $h(x)$ be the characteristic function of C . Then the region generated by this comb is irregular, on other hand $H(x)$ is finite for all $x \in [a, b] \setminus C$.

Example 2. We give an example of a comb such that the conditions of the previous theorem do not hold, moreover $H(x) = \infty$ for almost all $x \in [a, b]$. This comb is related to the Julia set of a polynomial $T(z) = z^2 - \lambda$ [46]. For $\lambda > 2$ there exists $h_0 > 0$ such that the Julia set of T is the preimage of the base of the comb given in Fig. 4.




Fig. 4. Comb related to the Julia set of $T(z) = z^2 - \lambda$.

Recall that almost every number x contains arbitrarily long strings of zeros in its dyadic representation, that is, for almost every x , and every non-negative integer N , there exists a string y_m of 0's and 1's, ending with 1, such that

$x = (y_m, \overbrace{0, \dots, 0}^N, \dots)$. Then $h(y_m) = 2^{-m}h_0$, and $|x - y_m| = 2^{-(m+N)}$. That is $H(x) \geq 2^N h_0$. In fact the Lebesgue measure of the Julia set is 0, i.e. the harmonic measure is singular continuous. Note that since the boundary is locally connected the region $\overline{\mathbb{C}} \setminus E$ is regular.

Even stronger condition is that

$$\sum_x h(x) < \infty, \tag{15}$$

in other words, the total length of slits is finite. This is the so-called Widom condition. It appears in his studies of asymptotics for extremal polynomials associated with a system of curves in the complex plane. Let $\pi_1(\Omega)$ be the fundamental group of the given region $\Omega = \overline{\mathbf{C}} \setminus E$. To a fixed character $\alpha \in \pi_1^*(\Omega)$ one associates the set of multi-valued (character-automorphic) analytic functions

$$H^\infty(\alpha) = \{f : f \circ \gamma = \alpha(\gamma)f, \forall \gamma \in \pi_1(\Omega), \sup_{z \in \Omega} |f(z)| < \infty\}.$$

The region Ω is of Widom type if the space $H^\infty(\alpha)$ is non-trivial (contains a non-constant function) for every $\alpha \in \pi_1^*(\Omega)$. A regular region $\Omega = \overline{\mathbf{C}} \setminus E$ is of Widom type if and only if (15) holds. For the role of this condition in the spectral theory of almost periodic Jacobi matrices see [48]. A well-known fact that the derivative of a conformal mapping on a region bounded by a rectifiable curve belongs to H^1 implies that the corresponding equilibrium measure μ is absolutely continuous.

5 More general combs

In this section we consider more general comb regions: those which satisfy property (ii) of the previous section. These regions D can be described as

$$D = \{x + iy : a < x < b, y > h(x)\}, \tag{16}$$

where $-\infty \leq a < b \leq \infty$ and h is an upper semi-continuous function on (a, b) . Let **Combs** be the set of such regions modulo horizontal shift, or equivalently, the set of all triples (a, b, h) modulo the equivalence relation $(a, b, h) \sim (a + c, b + c, h(x - c))$, $c \in \mathbf{R}$.

Let **Conf** be the set of univalent functions in \mathbf{H} such that $\phi(\mathbf{H}) \in \mathbf{Combs}$, normalized by $\phi(\infty) = \infty$, modulo the equivalence relation $\phi(z) \sim \phi(z - c)$, $c \in \mathbf{R}$.

Let R be the set of all analytic functions with positive real part in \mathbf{H} .

Let Sub be the set of all subharmonic functions v in the plane of the form

$$v(z) = \text{Re} \left(-az^2 + bz + \int_{\mathbf{R}} \left(\log \left(1 - \frac{z}{t} \right) + \frac{zt}{1+t^2} \right) d\mu(t) \right),$$

where $a > 0$, $b \in \mathbf{R}$ and μ is an increasing right-continuous function, such that

$$\int_0^\infty \frac{\mu(t) - \mu(-t)}{1+t^2} dt < \infty.$$

Two such functions are considered equivalent if their difference is constant.

Theorem 5. *There are the following canonical bijections between the sets Conf , R , Sub :*

$$\phi \mapsto \phi' : \text{Conf} \rightarrow R, \quad \phi \mapsto \text{Im } \phi : \text{Conf} \rightarrow \text{Sub}.$$

Moreover, $\text{Re } \phi = \mu + \text{const}$, $(2\pi)^{-1} \Delta v = d\mu$.

6 Uniform approximation and extremal problems

Here we consider several extremal problems whose solutions are expressed in terms of comb functions.

Applications of comb functions to extremal problems begins with the work of Akhiezer and Levin [5] on extension of Bernstein's inequality. Further applications are contained in [26, 27, 28, 29]. A survey of polynomials and entire functions of least deviation from zero on closed sets on the real line is given in [47].

Here we mention only few results.

1. Let f be an entire function of exponential type 1 satisfying $|f(x)| \leq 1$, $x < 0$ and $|f(x)| \leq B$, $x > 0$, where $B \geq 1$. One looks for maximal values of $|f(x)|$ for given x and of $|f'(0)|$, [13]. The extremal function is expressed in terms of the MO -comb with $h'_j = 0$, $j < 0$ and $h'_j = \cosh^{-1} B$, $j \geq 0$. Let $\theta : \mathbf{H} \rightarrow D$ be the conformal map onto the region (5), such that $\theta(z) \sim z$, as $z \rightarrow \infty$ non-tangentially, $\theta(0) = 0_-$. Set $x_1 = \theta^{-1}(ih_0)$. Then the function

$$f_0(x) = \begin{cases} B, & x > x_1, \\ \cos \theta(x), & 0 \leq x \leq x_1, \\ 1, & x < 0 \end{cases}$$

gives the solution of the first extremal problem: $|f(x)| \leq f_0(x)$ for f in the class described above, and $f'_0(0)$ is the maximal value of $|f'(0)|$.

2. Best uniform approximation of $\text{sgn}(x)$ on two rays/intervals. The simplest problem of this kind is to find the best uniform approximation of $\text{sgn}(x)$ on the set $X = (-\infty, -a] \cup [a, \infty)$ by entire functions of exponential type at most 1. The extremal entire function belongs to the MacLane class and its critical sequence is of the form

$$c_j = \begin{cases} -1 + (-1)^j L, & j \leq 0, \\ 1 + (-1)^j L, & j > 0, \end{cases} \quad (17)$$

where L is the error of the best approximation. Unfortunately, MacLane's functions do not have simple representations in terms of conformal mappings like (2) or (6), however in certain cases representation in terms of conformal maps of the kind described in section 5 can be obtained [15, 17].

3. Let us consider a uniform counterpart of the classical orthogonal Jacobi polynomials. Let $\alpha, \beta \geq 0$ and let $\mathcal{J}_n(x; \alpha, \beta) = x^n + \dots$ denote the monic polynomial of least deviation from zero on $[0, 1]$ with respect to the weight function $x^\alpha(1-x)^\beta$.

Lemma 3. *For non-negative α, β and an integer n*

$$x^\alpha(1-x)^\beta \mathcal{J}_n(x) = Le^\phi,$$

where ϕ is the conformal map on the V -comb region

$$D = \left\{ z = x + iy : -\beta < \frac{y}{\pi} < \alpha + n \right\} \setminus \bigcup_{j=0}^n \left\{ z = x + iy : \frac{y}{\pi} = j, x \leq 0 \right\}.$$

Such polynomials turn out to be useful in the description of multidimensional polynomials of least deviation from zero [37]. As an example we formulate the following theorem. Note that in multidimensional situation an extremal polynomial is not necessarily unique.

Theorem 6. [38] *A best polynomial approximation $P(z_1, \dots, z_d, \overline{z_1}, \dots, \overline{z_d})$ to the monomial $z_1^{k_1} \dots z_d^{k_d} \overline{z_1}^{l_1}$, $k_1 \geq l_1$, by polynomials of the total degree less than $k_1 + \dots + k_d + l_1$ in the ball $|z_1|^2 + \dots + |z_d|^2 \leq 1$ can be given in the form*

$$\begin{aligned} & z_1^{k_1} \dots z_d^{k_d} \overline{z_1}^{l_1} + P(z_1, \dots, z_d, \overline{z_1}, \dots, \overline{z_d}) \\ &= z_1^{k_1 - l_1} z_2^{k_2} \dots z_d^{k_d} \mathcal{J}_{l_1} \left(|z_1|^2; \frac{k_1 - l_1}{2}, \frac{k_2 + \dots + k_d}{2} \right). \end{aligned}$$

4. We finish this section with an extremal problem for entire functions of exponential type which arises in harmonic analysis [16].

For a fixed $\sigma > 0$, consider the class A_σ of entire functions which can be represented in the form

$$f(z) = \frac{1}{2\pi} \int_\gamma F(\zeta) e^{-i\zeta z} d\zeta,$$

where F is analytic in $\mathbf{C} \setminus [-\sigma, \sigma]$, $F(\infty) = 0$ and γ is a closed contour surrounding once the segment $[-\sigma, \sigma]$. We are interested in the upper estimate of the *upper density* of zeros of f ,

$$\bar{d}(f) = \limsup_{r \rightarrow \infty} \frac{n(r)}{r},$$

where $n(r)$ is the number of zeros, counting multiplicity in the disc $\{z : |z| \leq r\}$.

Theorem 7. *For a function $f \in A$, we have $\bar{d}(f) \leq c\sigma$, where $c \approx 1.508879$ is the unique solution of the equation*

$$\log(\sqrt{c^2 + 1} + c) = \sqrt{1 + c^{-2}} \quad \text{on } (0, +\infty). \quad (18)$$

This theorem is deduced from the solution of the following extremal problem for comb regions. *Among all univalent functions ϕ mapping H onto regions of the form $D = \{x + iy : y > h(|x|)\}$ with the properties $h(0) = 0, h(x) \leq 0, \phi(0) = 0, \phi(iy) \sim iy, y \rightarrow +\infty$, find the function with the largest $\operatorname{Re} \phi(1)$.*

The extremal region is described by

$$h_0(x) = \begin{cases} -\infty, & 0 < x < \pi c/2, \\ 0, & \text{otherwise.} \end{cases}$$

It is interesting that the same constant c as in (18) appears in the solution of another extremal problem [3, Appendix, 84] which has no apparent relation to Theorem 7.

7 Spectral theory and harmonic analysis

1. We say that an unbounded closed set E is homogeneous if there exists $\eta > 0$ such that for all $x \in E$ and all $\delta > 0$,

$$|(x - \delta, x + \delta) \cap E| \geq \eta\delta.$$

Theorem 8. [50] Let θ be a conformal map from the upper half-plane \mathbf{H} onto an MO-comb region D (Fig. 1, right). Assume that $E = \theta^{-1}(\mathbf{R})$ is homogeneous. Then E is the spectrum of a periodic canonical system, i.e., there exists an integrable on $[0, 1]$ non-negative 2×2 matrix function $H(t)$ of period 1, $H(t+1) = H(t)$, such that for an entire (transfer) matrix function $T(1, z)$ defined by the differential system

$$J\dot{T}(t, z) = zH(t)T(t, z), \quad T(0, z) = I, \quad J = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad (19)$$

the following relation holds

$$e^{i\theta} = \Delta - \sqrt{\Delta^2 - 1}, \quad \Delta(z) := (1/2) \operatorname{tr} T(1, z). \quad (20)$$

Moreover the parameter t in (19) corresponds to the “exponential type” of the matrix $T(t, z)$ with respect to the Martin function θ , that is,

$$t = \lim_{y \rightarrow +\infty} \frac{\log \|T(t, iy)\|}{\operatorname{Im} \theta(iy)}. \quad (21)$$

The whole collection of such matrices $H(t)$ for the given E can be parametrized by the characters of the fundamental group of the region $\Omega = \mathbf{C} \setminus E$.

The condition of homogeneity of E implies that $\Omega = \mathbf{C} \setminus E$ is of Widom type, and thus the region D_G , which corresponds to the Green function of Ω , satisfies Widom’s condition (15). Moreover, the so called Direct Cauchy Theorem holds in Ω . This fact plays a crucial role in the proof of Theorem 8. A very interesting and natural question: is it possible to characterize a Widom domain by means of geometric properties of the region D related to the Martin function of Ω ?

Example. A region D is defined by a system of slits forming a geometric progression

$$h_{jk} = \kappa j^k, \quad \kappa > 0, \quad h_0 = \infty,$$

otherwise $h_j = 0$. The corresponding set E is homogeneous.

2. Riesz bases. A sequence of vectors (e_n) in a Hilbert space H is called a *Riesz basis* if it is complete and there exist positive constants c, C such that

$$c \sum |a_n|^2 \leq \left\| \sum a_n e_n \right\|^2 \leq C \sum |a_n|^2$$

for every finite sequence (a_n) . A long-standing problem is how to find out whether for a given sequence of real exponents (λ_n) the sequence $e^{i\lambda_n x}$ is a Riesz basis in $L^2(-\pi, \pi)$. A recent result of Günter Semmler gives a parametric description of such Riesz bases.

We say that a sequence (d_n) , $d_n \geq 0$ satisfies the *discrete Muckenhoupt condition* if

$$\sum_{n \in I} d_n \sum_{n \in I} d_n^{-1} \leq C(\text{card } I)^2,$$

for every interval I of integers, and some $C > 0$.

Theorem 9. [44] *The sequence $(e^{i\lambda_n x})$ is a Riesz basis in $L^2(-\pi, \pi)$ if and only if it is the sequence of zeros of the entire function $f = \exp \phi$ of exponential type, where ϕ is a conformal map onto a V -comb with tips of the cuts h_n , and $\exp(2h_n)$ satisfies the discrete Muckenhoupt condition.*

For a given sequence (h_n) such that $(\exp(2h_n))$ satisfies the discrete Muckenhoupt condition, the conformal map ϕ can be always normalized so that $f = \exp \phi$ is of exponential type.

This theorem parametrizes all Riesz bases consisting of functions $e^{i\lambda_n x}$ in terms of sequences h_k .

We thank Misha Sodin for many illuminating discussions on the subject in the period 1980–2011.

References

- [1] N. I. Achieser, Vorlesungen der Approximationstheorie, Akademie-Verlag, Berlin 1967 xiii+412 pp.
- [2] N. Akhiezer, Elements of the theory of elliptic functions, AMS, Providence, RI, 1990.
- [3] N. Akhiezer, Theory of approximation, Frederick Ungar Publishing Co., New York 1956, Reprint by Dover Publications, Inc., New York, 1992.
- [4] N. Akhiezer, Selected works in 2 volumes, (Russian) Acta, Kharkiv, 2001.

- [5] N. Akhiezer and B. Levin, Generalization of S. N. Bernstein's inequality for derivatives of entire functions. (Russian) *Issledovaniya po sovremennym problemam teorii funktsii kompleksnogo peremennogo* pp. 111–165. Gosudarstv. Izdat. Fiz.-Mat. Lit., Moscow, 1960 (See also [4, vol. 2].)
- [6] V. Andrievskii, Positive harmonic functions on Denjoy domains in the complex plane, *J. d'Analyse*, 104 (2008) 83–124.
- [7] A.I. Aptekarev, Asymptotic properties of polynomials orthogonal on a system of contours, and periodic motions of Toda Lattices, *Mat. Sb.* 125 (1984) (Russian) 231–258; translation: *Math. USSR-Sbornik*, 53 (1986), 233–260.
- [8] M. Benedicks, Positive harmonic functions vanishing on the boundary of certain domains in R^n , *Ark. Mat.* 18 (1980), no. 1, 53–72.
- [9] R. P. Boas and A. C. Schaeffer, Variational methods in entire functions, *Amer. Journ. Math.*, 79, 1957, 857–884.
- [10] A. B. Bogatyrev, A combinatorial description of a moduli space of curves and of extremal polynomials, (Russian) *Mat. Sb.* 194 (2003), 27–48; translation: *Sb. Math.* 194 (2003), 1451–1473.
- [11] A. B. Bogatyrev, *Extremal polynomials and Riemann surfaces* (Russian), Moscow, MCNMO, 2005.
- [12] L. de Branges, *Hilbert spaces of entire functions*, Prentice-Hall, Inc., Englewood Cliffs, N.J. 1968.
- [13] A. Eremenko, Entire functions bounded on the real line, *Entire functions bounded on the real axis*, *Soviet Math. Dokl.* 37 (1988), no. 3, 693–695.
- [14] A. Eremenko, Geometric theory of meromorphic functions, in the book: *In the tradition of Ahlfors and Bers, III*, 221–230, *Contemp. Math.*, 355, Amer. Math. Soc., Providence, RI, 2004.
- [15] A. Eremenko and P. Yuditskii, Uniform approximation of $\operatorname{sgn}x$ by polynomials and entire functions, *J. Anal. Math.* 101 (2007), 313–324.
- [16] A. Eremenko and P. Yuditskii, An extremal problem for a class of entire functions, *C. R. Math. Acad. Sci. Paris* 346 (2008), no. 15-16, 825–828.

- [17] A. Eremenko and P. Yuditskii, Polynomials of the best uniform approximation to $\operatorname{sgn} x$ on two intervals, *J. d'Analyse*, 114 (2011) 285–315.
- [18] R. Gantmakher and M. Krein, Oscillation matrices and kernels and small vibrations of mechanical systems, AMS, Providence, RI, 2002.
- [19] A. Kesarev, Characterization of conformal maps of the upper half-plane on comb type domains, *J. Math. Phys. Anal. Geom.* 3 (1996) 290–307.
- [20] J. Geronimo, W. Van Assche, Orthogonal polynomials on several intervals via a polynomial mapping, *Trans. Amer. Math. Soc.*, 308 (1988) 559–581.
- [21] I. Hur, Ch. Remling, Ergodic Jacobi matrices and conformal maps, arXiv:1108.5370.
- [22] B. Kjellberg, On the growth of minimal positive harmonic functions in a plane region, *Ark. mat.* 1 (1950) 347–351.
- [23] M. Krein, On inverse problems of the theory of filters and λ -zones of stability (Russian) *Doklady Akad. Nauk SSSR (N.S.)* 93 (1953) 767–770. (English transl, see MR0073775).
- [24] N. Landkof, Foundations of modern potential theory, Springer, NY 1972.
- [25] B. Levin, Distribution of zeros of entire functions, AMS, Providence, R.I., 1980.
- [26] B. Levin, Majorants in classes of subharmonic functions, *Function theory, Func. Anal. and Appl.*, 51 (1989) 3–17.
- [27] B. Levin, Connection of the majorant with a conformal map, part 2, *Function theory, Func. Anal. and Appl.*, 52 (1989) 3–21.
- [28] B. Levin, Classification of closed sets on R and representation of a majorant, part 3, *Function theory, Func. Anal. and Appl.*, 52 (1989) 21–33.
- [29] B. Levin, Completeness of system of functions, quasi-analyticity and subharmonic majorants, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* 170 (1989), *Issled. Linein. Oper. Teor. Funktsii.* 17, 102–156, 322–323; translation in *J. Soviet Math.* 63 (1993), no. 2, 171–201.

- [30] G. MacLane, Concerning the uniformization of certain Riemann surfaces allied to the inverse-cosine and inverse-gamma surfaces, *Trans. Amer. Math. Soc.* 62 (1947) 99–113.
- [31] V. A. Malyshev, Abel’s equation, (Russian) *Algebra i Analiz* 13 (2001), 1–55; translation in *St. Petersburg Math. J.* 13 (2002), no. 6, 893–938.
- [32] McMillan, Boundary behavior of a conformal mapping, *Acta Math.*, 123 (1969) 43–67.
- [33] V. Marchenko, *Sturm–Liouville operators and applications*, AMS Chelsea Publishing, Providence, RI, 2011.
- [34] V. Marchenko, *Introduction to the theory of inverse problems of spectral analysis*, Acta, Kharkiv, 2005 (Russian).
- [35] V. Marchenko and I. Ostrovskii, A characterization of the spectrum of the Hill operator, *Math. USSR Sbornik* 26, 4 (1975) 493–554.
- [36] N. N. Meiman, Solution of the fundamental problems of the theory of polynomials and entire functions which deviate least from zero, (Russian) *Trudy Moskov. Mat. Obshch.* 9 1960, 507–535.
- [37] I. Moale and P. Yuditskii, On Complex (Non-Analytic) Chebyshev Polynomials in \mathbf{C}^2 , *Comp. Methods and Function Theory*, 11 (2011) 13–24.
- [38] I. Moale, *Explicit min-max polynomials in several variables*, PhD thesis, 2010, JKU, Linz.
- [39] F. Nazarov, F. Peherstorfer, A. Volberg and P. Yuditskii, Asymptotics of the best polynomial approximation of $|x|^p$ and of the best Laurent polynomial approximation of $\text{sgn}(x)$ on two symmetric intervals. *Constr. Approx.* 29 (2009), no. 1, 23–39.
- [40] F. Peherstorfer, On Bernstein–Szegő orthogonal polynomials on several intervals II, *Orthogonal polynomials with periodic recurrence coefficients*, *J. Approx. Theory*, 64 (1991) 123–161.
- [41] F. Peherstorfer, Orthogonal and extremal polynomials on several intervals, *J. Com. ppl. Math.*, 48 (1993) 187–205.

- [42] F. Peherstorfer, Minimal polynomials on several intervals with respect to the maximum norm – a survey. Complex methods in approximation theory (Almeria, 1995), 137–159, Univ. Almeria, Almeria, 1997.
- [43] Ch. Pommerenke, Boundary behaviour of conformal maps, Springer-Verlag, Berlin, 1992.
- [44] G. Semmler, Complete interpolating sequences, the discrete Muckenhoupt condition, and conformal mapping, Ann. Acad. Sci. Fenn. Math. 35 (2010), no. 1, 23–46.
- [45] B. Simon, Szegő’s theorem and its descendants, Princeton UP, 2011.
- [46] M. Sodin and P. Yuditskii, The limit-periodic finite-difference operator on $l^2(Z)$ associated with iterations of quadratic polynomials, J. Stat. Phys. 60 (1990), no. 5-6, 863–873.
- [47] M. Sodin and P. Yuditskii, Functions that deviate least from zero on closed subsets of the real axis (Russian) Algebra i Analiz 4 (1992), no. 2, 1–61; translation in St. Petersburg Math. J. 4 (1993), no. 2, 201–249.
- [48] M. Sodin and P. Yuditskii, Almost periodic Jacobi matrices with homogeneous spectrum, infinite-dimensional Jacobi inversion, and Hardy spaces of character-automorphic functions, J. Geom. Anal. 7 (1997), no. 3, 387–435.
- [49] E. Vinberg, Real entire functions with prescribed critical values, in the book: Problems of group theory and homological algebra, Yaroslavl’ Gos. Un-t., Yaroslavl, 1989, 127–138 (Russian).
- [50] P. Yuditskii, A special case of de Branges’ theorem on the inverse monodromy problem, Integral Equations Operator Theory 39 (2001), no. 2, 229–252.

*Department of Mathematics, Purdue University
West Lafayette, IN 47907 USA
eremenko@math.purdue.edu*

*Abteilung für Dynamische Systeme und Approximationstheorie,
Johannes Kepler Universität Linz,
A-4040 Linz, Austria
Petro.Yudytskiy@jku.at*