

Simultaneous diagonalization of two quadratic forms and a generalized eigenvalue problem

A. Eremenko

November 28, 2019

Theorem. *Let A, M be two real symmetric matrices of the same size, and let M be positive definite. Then there exists a non-singular matrix C such that*

$$C^T M C = I, \tag{1}$$

and

$$C^T A C = \Lambda, \tag{2}$$

where Λ is a real diagonal matrix.

Proof. We have

$$M = R^T R, \tag{3}$$

with some non-singular matrix R . Then the matrix

$$(R^{-1})^T A R^{-1}$$

is symmetric, so there exists an orthogonal matrix B such that

$$B^{-1}(R^{-1})^T A R^{-1} B = \Lambda. \tag{4}$$

Set

$$C = R^{-1} B. \tag{5}$$

Then $C^T = B^T (R^{-1})^T = B^{-1} (R^{-1})^T$, where we used that B is orthogonal. So (4) is the same as (2). To check (1) we use $B^T = B^{-1}$ and $(R^{-1})^T = (R^T)^{-1}$ and obtain

$$C^T M C = B^{-1} (R^T)^{-1} R^T R R^{-1} B = I.$$

This proves Theorem 1.

Next we will show that the entries λ_j of the diagonal matrix Λ in this theorem are *generalized eigenvalues* of A with respect to M :

$$Ax = \lambda Mx, \quad x \neq 0. \quad (6)$$

They can be determined from the characteristic equation

$$\det(A - \lambda M) = 0,$$

and the *generalized eigenvectors* are the columns of C from (1), (2).

We obtain from Theorem 1 and from its proof:

Corollary. *Let A, M be symmetric matrices of the same size, and let M be positive definite. Then all generalized eigenvalues (6) are real, and there is a basis of the whole space which consists of generalized eigenvectors.*

Proof. We refer to the proof of Theorem 1. Matrix $(R^{-1})^T AR^{-1}$ is symmetric, therefore all its eigenvalues are real and the eigenvectors form a basis. These eigenvectors are columns of B . If v_j is an eigenvector of $(R^{-1})^T AR^{-1}$ with eigenvalue λ_j then

$$(R^{-1})^T AR^{-1}v_j = \lambda_j v_j.$$

Multiplying on R^T and setting $v_j = Ru_j$ we obtain

$$Au_j = \lambda_j R^T Ru_j = \lambda_j R^T Ru_j = \lambda_j Mu_j.$$

As R is non-singular, the u_j form a basis of the space.

Since $B = RC$, this basis is nothing but the columns of C of Theorem 1. So the simultaneous diagonalization of two matrices is not more difficult than diagonalization of one matrix: solve the generalized characteristic equation and find generalized eigenvectors.

Notice that

$$u_i^T Mu_j = u_i R^T Ru_j = (Ru_i)^T (Ru_j) = v_i^T v_j = \begin{cases} 0, & i \neq j \\ 1, & i = j. \end{cases}$$

This means that u_j , the eigenvectors of (6) are orthonormal with respect to the dot product defined by

$$(x, y)_M = x^T Mx,$$

and our matrix R transforms this dot product to the standard dot product:

$$(x, y)_M = x^T M y = x^T R^T R y = (R x, R y).$$

Applications to mechanics.

Newton's form of equations of motion $ma = F$ is not always convenient, especially when one deals with curvilinear coordinates. A generalization was proposed by Lagrange. We consider a system of points whose position is completely determined by some *generalized coordinates* $\mathbf{q} = (q_1, \dots, q_n)$. For example, for one free point in space we have three coordinates $(q_1, q_2, q_3) = (x_1, x_2, x_3)$. Or \mathbf{q} may be cylindrical, or spherical coordinates. For m free points in space we need $n = 3m$ coordinates. For a pendulum oscillating in a vertical plane, we need one coordinate, for example the angle of deviation of this pendulum from the vertical is a convenient coordinate.

As the system moves, coordinates are functions of time $q_j(t)$. Their derivatives are called *generalized velocities*, $\dot{q} = dq/dt$. Derivatives with respect to time are usually denoted by dots over letters in mechanics, to distinguish them from other derivatives. To obtain the true velocity vector of a point $\mathbf{x}_k \in \mathbf{R}^3$, one has to write $\mathbf{x}_k = f_k(\mathbf{q})$, rectangular coordinates as functions of generalized coordinates, and differentiate:

$$\dot{\mathbf{x}}_k = \sum_{j=1}^n \frac{\partial f_k}{\partial q_j} \dot{q}_j,$$

and the kinetic energy is

$$T_k = m \|\dot{\mathbf{x}}_k\|^2 / 2 = \sum b_{k,i,j}(\mathbf{q}) \dot{q}_i \dot{q}_j, \quad (7)$$

where $b_{k,i,j}$ are some functions of \mathbf{q} . The total kinetic energy T of the system is the sum of such expressions over all points \mathbf{x}_k , $T = \sum T_k$. The important fact is that

Kinetic energy is a positive definite quadratic form of generalized velocities, with coefficients depending on the generalized coordinates.

It is positive definite because the LHS of (7) is non-negative and the sum of such expressions is positive, if at least one point actually moves.

Now we assume that *vector of forces is the gradient of some function $-U$ of generalized coordinates*:

$$F = -\text{grad } U = - \left(\frac{\partial U}{\partial q_1}, \dots, \frac{\partial U}{\partial q_n} \right). \quad (8)$$

This function U is called the potential energy or simply the *potential*.

Following Lagrange's recipe, we form the following function of generalized coordinates and velocities:

$$L = T - U,$$

the difference between kinetic and potential energy. This function is called the *Lagrangian* of the system. The equations of motion in the form of Lagrange are

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_j} \right) = \frac{\partial L}{\partial q_j}, \quad 1 \leq j \leq n. \quad (9)$$

The advantage of this formulation is that unlike for Newton's equations *arbitrary* curvilinear coordinate system can be used.

To see that these equations indeed generalize Newton's equations, consider a free point with coordinate $\mathbf{x} = (x_1, x_2, x_3)$ and mass m moving in the field of force with potential U . Then the kinetic energy is

$$T = \frac{m}{2} (\dot{x}_1^2 + \dot{x}_2^2 + \dot{x}_3^2),$$

and the Lagrangian is $L = T - U$. So equations (9) become

$$\frac{d}{dt}(m\dot{x}_j) = -\frac{\partial U}{\partial x_j} = F_j(x_1, x_2, x_3),$$

where F_j is the j -th component of the force.

Equations of motion are usually non-linear and cannot be solved.

One of the most common methods of dealing with them is *linearization*, that is approximation of non-linear equations by linear ones. The simplest case is the linearization near an equilibrium. An equilibrium is a point \mathbf{q}^0 such that the system in this state does not move. This means that equations (9) are satisfied by $\mathbf{q}(t) \equiv \mathbf{q}^0$.

Theorem 2. *A point \mathbf{q}^0 is an equilibrium if and only if it is a critical point of the potential energy U .*

Proof. Let us write (9) as

$$\frac{d}{dt} \frac{\partial T}{\partial \dot{q}_j} = \frac{\partial T}{\partial q_j} - \frac{\partial U}{\partial q_j}.$$

If $\mathbf{q}(t) \equiv \mathbf{q}^0$ is a solution, then $\dot{\mathbf{q}} = 0$ and thus $\partial T / \partial \dot{q}_j = 0$, and $\partial T / \partial q_j = 0$ for all j . So $\partial U / \partial q_j = 0$.

For the linearization we assume without loss of generality that $\mathbf{q}^0 = 0$, and that both T and U are *analytic* functions of \mathbf{q} and $\dot{\mathbf{q}}$. This means that they have convergent series expansions

$$T(q, \dot{q}) = T_0 + T_1(q, \dot{q}) + T_2(q, \dot{q}) + \dots,$$

where T_k are homogeneous polynomials of the variables q_j, \dot{q}_j . Similar expansion holds for U . When we differentiate a homogeneous polynomial, its degree decreases by 1, so to obtain *linear* equations in (9) both T and U have to be of degree 2. As T is of degree 2 in the variables \dot{q} , it must be independent of q . In U , the terms of the first degree vanish by Theorem 2, and the constant term disappears after differentiation.

Thus

The Lagrangian of the linearized systems at an equilibrium point 0 is obtained by setting $\mathbf{q} = 0$ in (7) and keeping only quadratic terms in U , in other words, this Lagrangian has the form

$$L = \sum_{i,j} m_{i,j} \dot{q}_i \dot{q}_j - \sum_{i,j} a_{i,j} q_i q_j, \quad (10)$$

the difference of two quadratic forms with matrices M and A such that M is positive definite.

To write the Lagrange equations with Lagrangian (10) we need the differentiation formula

$$\frac{d}{d\mathbf{x}}(\mathbf{x}^T A \mathbf{x}) = A \mathbf{x}.$$

Here $d/d\mathbf{x}$ is the column $(d/dx_1, \dots, d/dx_n)^T$. So the equation of motion with Lagrangian (10) is

$$\frac{d}{dt} M \dot{\mathbf{x}} = -A \mathbf{x}. \quad (11)$$

Now we can use the theorem on simultaneous diagonalization. We can apply it directly to (10) to conclude that there are new coordinates \mathbf{y} , $\mathbf{x} = C \mathbf{y}$, such that

$$L = \dot{\mathbf{y}}^T \dot{\mathbf{y}} - \mathbf{y}^T \Lambda \mathbf{y},$$

so the equations of motion *decouple* and become


$$\ddot{y}_j = -\lambda_j y_j. \quad (12)$$

Or alternatively we can apply the Corollary to the linear equation (11) and find a basis u_j of generalized eigenvectors. If y_j are coordinates with respect to this basis, we obtain (12) again. Stated in words this means that *the linearized equation of small oscillations always decouples and becomes (12) after a change of coordinates.*

Notice that if the matrix A is positive definite, then all $\lambda_j > 0$ and solutions have the form $y_j(t) = c_j e^{\pm i\omega_j t}$, where $\omega_j = \sqrt{\lambda_j}$ the system is stable and solutions oscillate with frequencies ω_j .

Example. Double pendulum.

The configuration is shown in the figure. Let us choose the angles between the two rods and the vertical direction as generalized coordinates q_1, q_2 . Angles are measured from the downward vertical direction, counterclockwise, as shown in the picture.



Then the kinetic energy of the mass m_1 is

$$T_1 = \frac{m_1}{2} \ell_1^2 \dot{q}_1^2,$$

and the kinetic energy of the second mass is

$$T_2 = \frac{m_2}{2} \left(\ell_1^2 \dot{q}_1^2 + \ell_2^2 \dot{q}_2^2 + 2\ell_1 \ell_2 \cos(q_2 - q_1) \dot{q}_1 \dot{q}_2 \right).$$

Potential energy of the system is

$$U = -(m_1 + m_2)g\ell_1 \cos q_1 - m_2 g \ell_2 \cos q_2.$$

Thus $T = T_1 + T_2$ and

$$\begin{aligned} L = T - U &= \frac{m_1 + m_2}{2} \ell_1^2 \dot{q}_1^2 + \frac{m_2}{2} \ell_2^2 \dot{q}_2^2 + m_2 \ell_1 \ell_2 \cos(q_2 - q_1) \dot{q}_1 \dot{q}_2 \\ &+ (m_1 + m_2)g\ell_1 \cos q_1 + m_2 g \ell_2 \cos q_2. \end{aligned}$$

The equations of motion are non-linear and difficult to solve, so we linearize them near the equilibrium $(q_1, q_2) = (0, 0)$. (There are four equilibria in our system). Linearization in this case means that we replace the cosine in the kinetic energy by 1 and the cosines in potential according to the formula $\cos x \approx 1 - x^2/2$, because we want to keep only second degree terms in the Lagrangian. The constant term in Potential energy can be omitted.

Thus the Lagrangian of the linearized system is

$$\begin{aligned} L^* &= \frac{m_1 + m_2}{2} \ell_1^2 \dot{q}_1^2 + \frac{m_2}{2} \ell_2^2 \dot{q}_2^2 + m_2 \ell_1 \ell_2 \dot{q}_1 \dot{q}_2 \\ &- \frac{m_1 + m_2}{2} g \ell_1 q_1^2 - \frac{m_2}{2} g \ell_2 q_2^2. \end{aligned}$$

and the linearized equation of motion is

$$\begin{pmatrix} (m_1 + m_2)\ell_1^2 & m_2\ell_1\ell_2 \\ m_2\ell_1\ell_2 & m_2\ell_2^2 \end{pmatrix} \begin{pmatrix} \ddot{q}_1 \\ \ddot{q}_2 \end{pmatrix} = - \begin{pmatrix} (m_1 + m_2)g\ell_1 & 0 \\ 0 & m_2g\ell_2 \end{pmatrix} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix},$$

which we write as

$$M\ddot{q} = -Aq.$$

It is easy to check directly that both M and A are positive definite. Notice that M is not diagonal in this example.