

Spectral Theorems for Hermitian and unitary matrices

A. Eremenko

October 26, 2017

1. An *Hermitian product* on a complex vector space V is an assignment of a complex number (x, y) to each pair of vectors x, y , which has the following properties for all vectors x, y, z and for all numbers α, β :

$$(x, y) = \overline{(y, x)},$$

$$(x, \alpha y + \beta z) = \alpha(x, y) + \beta(x, z),$$

$$(x, x) \geq 0,$$

with equality only for $x = 0$.

Example. $(x, y) = \overline{x_1}y_1 + \dots + \overline{x_n}y_n$. This example is called the *standard Hermitian product* on \mathbf{C}^n .

It follows from the first two properties that $(\alpha x, y) = \overline{\alpha}(x, y)$. They say that (x, y) is linear with respect to the second argument and *anti-linear* with respect to the first one.

An *Hermitian transposition* is the combination of two operations: ordinary transposition and complex conjugation. It is denoted by star, $A^* = \overline{A}^T$, where the bar is the complex conjugation. So the standard Hermitian product can be written as $(x, y) = x^*y$.

Two vectors are called *orthogonal* if $(x, y) = 0$. Vectors orthogonal to some given set of vectors form a subspace. If V' is a subspace of V then its *orthogonal complement* consists of all vectors orthogonal to each vector of V' . Two subspaces are called orthogonal if each vector of one of them is orthogonal to each vector of another one.

A square matrix A is called *Hermitian* if

$$A^* = A.$$

A real matrix is Hermitian if and only if it is symmetric. Hermitian matrices are characterized by the property

$$(Ax, y) = (x, Ay), \quad \text{for all } x, y \text{ in } V, \quad (1)$$

where $(., .)$ is the standard Hermitian product. Indeed, $A^* = A$ is equivalent to

$$(Ax, y) = (Ax)^*y = x^*Ay = (x, Ay), \quad \text{for all } x, y \text{ in } V.$$

A square matrix U is called *unitary* if

$$U^*U = I,$$

which is the same as $U^* = U^{-1}$. In other words, a unitary matrix is such that its columns are orthonormal. Unitary matrices are characterized by the property

$$(Ux, Uy) = (x, y) \quad \text{for all } x, y \text{ in } V. \quad (2)$$

Indeed,

$$(Ux, Uy) = (Ux)^*Uy = x^*U^*Uy = x^*y = (x, y).$$

A real matrix is unitary if and only if it is orthogonal.

2. Spectral theorem for Hermitian matrices. *For an Hermitian matrix:*

- a) all eigenvalues are real,*
- b) eigenvectors corresponding to distinct eigenvalues are orthogonal,*
- c) there exists an orthogonal basis of the whole space, consisting of eigenvectors.*

Thus all Hermitian matrices are diagonalizable. Moreover, for every Hermitian matrix A , there exists a unitary matrix U such that

$$AU = U\Lambda,$$

where Λ is a real diagonal matrix. The diagonal entries of Λ are the eigenvalues of A , and columns of U are eigenvectors of A .

Proof of Theorem 2. a). Let λ be an eigenvalue, then

$$Ax = \lambda x, \quad x \neq 0$$

for some vector x . Multiply both sides on x :

$$(Ax, x) = (\lambda x, x) = \bar{\lambda}(x, x).$$

Property (1) shows that (Ax, x) equals

$$(x, Ax) = (x, \lambda x) = \lambda(x, x).$$

As $(x, x) \neq 0$, we conclude that $\lambda = \bar{\lambda}$, that is λ is real. This proves a).

Proof of b). Suppose we have two distinct eigenvalues $\lambda \neq \mu$. Then

$$Ax = \lambda x, \quad Ay = \mu y, \tag{3}$$

where x, y are eigenvectors. Multiply the first equation on y , use (1) and the fact that λ is real which was just established.

$$\lambda(x, y) = (\lambda x, y) = (Ax, y) = (x, Ay) = (x, \mu y) = \mu(x, y).$$

As $\lambda \neq \mu$, we conclude that $(x, y) = 0$, which proves b).

Proof of c). Let λ_1 be an eigenvalue, and x_1 an eigenvector corresponding to λ_1 (every square matrix has an eigenvalue and an eigenvector). Let V_1 be the set of all vectors orthogonal to x_1 . Then A maps V_1 into itself: for every $x \in V_1$ we also have $Ax \in V_1$. Indeed, $x \in V_1$ means that $(x_1, x) = 0$, then we have using (1):

$$(x_1, Ax) = (Ax_1, x) = \lambda_1(x_1, x) = 0,$$

so $x \in V_1$. Now the linear operator $L(x) = Ax$ when restricted to V_1 is also Hermitian, and it has an eigenvalue λ_2 and an eigenvector $x_2 \in V_1$. By definition of V_1 , x_2 is orthogonal to x_1 . Let V_2 be the orthogonal complement of the span of x_1, x_2 . Then A also maps V_2 into itself, as before. Continuing this way, we find a sequence λ_k, x_k and subspaces V_k containing x_k such that V_k is orthogonal to x_1, \dots, x_{k-1} . The sequence must terminate on the n -th step because $\dim V_k = n - k$: on every step dimension decreases by 1. This completes the proof.

3. Spectral theorem for unitary matrices. *For a unitary matrix:*

- a) all eigenvalues have absolute value 1.*
- b) eigenvectors corresponding to distinct eigenvalues are orthogonal,*
- c) there is an orthogonal basis of the whole space, consisting of eigenvectors.*

Thus unitary matrices are diagonalizable. Moreover, for each unitary matrix A there exists a unitary matrix U such that

$$AU = U\Lambda$$

where U is a diagonal matrix whose diagonal entries have absolute value 1. The columns of U are eigenvectors of A .

Proof of Theorem 2. a) Let λ be an eigenvalue. Then

$$Ax = \lambda x, \quad x \neq 0.$$

Using (2) we obtain

$$(x, x) = (Ax, Ax) = \bar{\lambda}\lambda(x, x).$$

As $(x, x) \neq 0$, we conclude that $\bar{\lambda}\lambda = |\lambda|^2 = 1$, which proves a).

Proof of b). Begin with (3), and multiply the two equations in (3):

$$(x, y) = (Ax, Ay) = \bar{\lambda}\mu(x, x).$$

As $|\lambda| = 1$, we conclude that $\bar{\lambda} = \lambda^{-1}$, so the multiple in the RHS is $\mu/\lambda \neq 1$ by our assumption that $\mu \neq \lambda$. So $(x, y) = 0$, which proves b).

Proof of c). Let λ_1 be an eigenvalue, and x_1 an eigenvector corresponding to this eigenvalue, Let V_1 be the set of all vectors orthogonal to x_1 . As in the proof in section 2, we show that $x \in V_1$ implies that $Ax \in V_1$. Indeed

$$(Ax, x_1) = (x, A^*x_1) = (x, A^{-1}x_1) = \lambda^{-1}(x, x_1) = 0,$$

where we used (2) which is equivalent to $A^* = A^{-1}$. The proof is now completed in exactly the same way as in the previous section.

4. Exponentials of Hermitian matrices. *Let A be an Hermitian matrix. Then e^{iA} is unitary, and conversely, every unitary matrix has the form e^{iA} for some Hermitian matrix A .*

Let B be a real matrix, and $A = iB$. Then A is Hermitian if and only if B is skew symmetric ($B^T = -B$):

$$A^* = (-i)B^T = iB = A.$$

So we obtain a

Corollary: For a real matrix B , e^B is orthogonal if and only if B is skew-symmetric.

Proof. Let $U = e^{iA}$, where A is Hermitian. Then

$$UU^* = e^{iA}e^{-iA^*} = e^{iA}e^{-iA} = I.$$

Conversely, let U be a unitary matrix. Then, by the Spectral Theorem for unitary matrices (section 3), there is another unitary matrix B such that $U = B\Lambda B^{-1}$, and $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$. As all $|\lambda_k| = 1$, we write them as $\lambda_k = e^{i\theta_k}$, where θ_k are real numbers. Then set

$$A = B \text{diag}(\theta_1, \dots, \theta_n) B^{-1} = B\Lambda_1 B^{-1}.$$

Then A is Hermitian:

$$A^* = (B^{-1})^* \Lambda_1 B^* = B\Lambda_1 B^{-1} = A,$$

and evidently $\exp(iA) = U$.

5. These three theorems can be generalized to infinite-dimensional spaces. Unlike the Jordan form theorem. One can say that we understand well Hermitian and unitary operators, but not arbitrary linear operators.

These three theorems and their infinite-dimensional generalizations make the mathematical basis of the most fundamental theory about the real world that we possess, namely quantum mechanics.