

MA 266

Review Topics - Exam # 2 (updated)**(1) First Order Differential Equations.** (Separable, 1st Order Linear, Homogeneous, Exact)**(2) Second Order Linear Homogeneous with Equations Constant Coefficients .**

The differential equation $ay'' + by' + cy = 0$ has *Characteristic Equation* $ar^2 + br + c = 0$. Call the roots r_1 and r_2 . The general solution of $ay'' + by' + cy = 0$ is as follows:

- (a) If r_1, r_2 are real and distinct $\Rightarrow y = C_1 e^{r_1 t} + C_2 e^{r_2 t}$
- (b) If $r_1 = \lambda + i\mu$ (hence $r_2 = \lambda - i\mu$) $\Rightarrow y = C_1 e^{\lambda t} \cos \mu t + C_2 e^{\lambda t} \sin \mu t$
- (c) If $r_1 = r_2$ (repeated roots) $\Rightarrow y = C_1 e^{r_1 t} + C_2 t e^{r_1 t}$

(3) Theory of 2nd Linear Order Equations.

Wronskian of y_1, y_2 is $W(y_1, y_2)(t) = \begin{vmatrix} y_1(t) & y_2(t) \\ y_1'(t) & y_2'(t) \end{vmatrix}$.

- (a) The functions $y_1(t)$ and $y_2(t)$ are linearly independent over $a < t < b$ if $W(y_1, y_2) \neq 0$ for at least one point in the interval.

- (b) **THEOREM (Existence & Uniqueness)** If $p(t), q(t)$ and $g(t)$ are continuous in an open interval $\alpha < t < \beta$ containing t_0 , then the IVP $\begin{cases} y'' + p(t)y' + q(t)y = g(t) \\ y(t_0) = y_0 \\ y'(t_0) = y_1 \end{cases}$ has a unique solution $y = \phi(t)$ defined in the open interval $\alpha < t < \beta$.

- (c) **Superposition Principle** If $y_1(t)$ and $y_2(t)$ are solutions of the 2nd order linear homogeneous equation $P(t)y'' + Q(t)y' + R(t)y = 0$ over the interval $a < t < b$, then $y = C_1 y_1(t) + C_2 y_2(t)$ is also a solution for any constants C_1 and C_2 .

- (d) **THEOREM (Homogeneous)** If $y_1(t)$ and $y_2(t)$ are solutions of the linear homogeneous equation $P(t)y'' + Q(t)y' + R(t)y = 0$ in some interval I and $W(y_1, y_2) \neq 0$ for some t_1 in I , then the general solution is $y_c(t) = C_1 y_1(t) + C_2 y_2(t)$. This is usually called the *complementary solution* and we say that $y_1(t), y_2(t)$ form a *Fundamental Set of Solutions* (FSS) to the differential equation.

- (e) **THEOREM (Nonhomogeneous)** The general solution of the nonhomogeneous equation

$$P(t)y'' + Q(t)y' + R(t)y = G(t)$$

is $y(t) = y_c(t) + y_p(t)$, where $y_c(t) = C_1 y_1(t) + C_2 y_2(t)$ is the general solution of the corresponding homogeneous equation $P(t)y'' + Q(t)y' + R(t)y = 0$ and $y_p(t)$ is a particular solution of the nonhomogeneous equation $P(t)y'' + Q(t)y' + R(t)y = G(t)$.

- (f) **Useful Remark** : If $y_{p_1}(t)$ is a particular solution of $P(t)y'' + Q(t)y' + R(t)y = G_1(t)$ and if $y_{p_2}(t)$ is a particular solution of $P(t)y'' + Q(t)y' + R(t)y = G_2(t)$, then

$$y_p(t) = y_{p_1}(t) + y_{p_2}(t)$$

is a particular solution of $P(t)y'' + Q(t)y' + R(t)y = [G_1(t) + G_2(t)]$.

(4) Reduction of Order. If $y_1(t)$ is one solution of $P(t)y'' + Q(t)y' + R(t)y = 0$, then a second solution may be obtained using the substitution $y = v(t)y_1(t)$. This reduces the original 2^{nd} order equation to a 1^{st} equation using the substitution $w = \frac{dv}{dt}$. Solve that first order equation for w , then since $w = \frac{dv}{dt}$, solve this 1^{st} order equation to determine the function v .

(5) Finding A Particular Solution $y_p(t)$ to Nonhomogeneous Equations.

You can always use the method of Variation of Parameters to find a particular solution $y_p(t)$ of the linear nonhomogeneous equation $y'' + p(t)y' + q(t)y = g(t)$. Variation of Parameters may require integration techniques.

If the coefficients of the differential equation are constants rather than functions **and** if $g(t)$ has a very special form (see table below), it is usually easier to use Undetermined Coefficients :

(a) UNDETERMINED COEFFICIENTS - **IF** $ay'' + by' + cy = g(t)$ **AND** $g(t)$ is as below:

$g(t)$	Form of $y_p(t)$
$P_m(t) = a_mt^m + a_{m-1}t^{m-1} + \dots + a_0$	$t^s \{A_mt^m + A_{m-1}t^{m-1} + \dots + A_0\}$
$e^{\alpha t} P_m(t)$	$t^s \{e^{\alpha t} (A_mt^m + A_{m-1}t^{m-1} + \dots + A_0)\}$
$e^{\alpha t} P_m(t) \cos \beta t$ or $e^{\alpha t} P_m(t) \sin \beta t$	$t^s \{e^{\alpha t} [F_m(t) \cos \beta t + G_m(t) \sin \beta t]\}$

where $s =$ the smallest nonnegative integer ($s = 0, 1$ or 2) such that no term of $y_p(t)$ is a solution of the corresponding homogeneous equation. In other words, no term of $y_p(t)$ is a term of $y_c(t)$. ($F_m(t)$, $G_m(t)$ are both polynomials of degree m .)

(b) VARIATION OF PARAMETERS - If $y_1(t)$ and $y_2(t)$ are two independent solutions of the homogeneous equation $y'' + p(t)y' + q(t)y = 0$, then a particular solution $y_p(t)$ of the nonhomogeneous equation

$$y'' + p(t)y' + q(t)y = g(t) \quad (*)$$

has the form

$$y_p(t) = u_1(t)y_1(t) + u_2(t)y_2(t)$$


where

$$u_1' = \frac{\begin{vmatrix} 0 & y_2 \\ g(t) & y_2' \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}}, \quad u_2' = \frac{\begin{vmatrix} y_1 & 0 \\ y_1' & g(t) \end{vmatrix}}{\begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}}.$$

Remember: Coefficient of y'' in $(*)$ must be “1” in order to use the above formulas.

(6) Spring-Mass Systems $\begin{cases} m u'' + \gamma u' + k u = F(t) \\ u(0) = u_0, \quad u'(0) = u_1 \end{cases}$

m = mass of object, γ = damping constant, k = spring constant, $F(t)$ = external force Weight $w = m g$, Hooke's Law: $F_s = k d$,



I Undamped Free Vibrations : $m u'' + k u = 0$ (Simple Harmonic Motion)

Note that $A \cos \omega_0 t + B \sin \omega_0 t = R \cos(\omega_0 t - \delta)$, where $R = \sqrt{A^2 + B^2}$ = *amplitude*, ω_0 = *frequency*, $\frac{2\pi}{\omega_0}$ = *period* and δ = *phase shift* determined by $\tan \delta = \frac{B}{A}$.

II Damped Free Vibrations : $m u'' + \gamma u' + k u = 0$

- (i) $\gamma^2 - 4km > 0$ (*overdamped*) \iff distinct real roots to CE
- (ii) $\gamma^2 - 4km = 0$ (*critically damped*) \iff repeated roots to CE
- (iii) $\gamma^2 - 4km < 0$ (*underdamped*) \iff complex roots to CE (motion is *oscillatory*)

III Forced Vibrations : ($F(t) = F_0 \cos \omega t$ or $F(t) = F_0 \sin \omega t$, for example)

- (i) $m u'' + \gamma u' + k u = F(t)$ (Damped) In this case if you write the general solution as $u(t) = u_T(t) + u_\infty(t)$, then $u_T(t)$ = *Transient Solution* (i.e. the part of $u(t)$ such that $u_T(t) \rightarrow 0$ as $t \rightarrow \infty$) and $u_\infty(t)$ = *Steady-State Solution* (the solution behaves like this function in the long run).

- (ii) $m u'' + k u = F_0 \cos \omega t$ (Undamped) If $\omega = \omega_0 = \sqrt{\frac{k}{m}} \Rightarrow$ Resonance occurs and the solution is unbounded; while if $\omega \neq \omega_0$ then motion is a series of *beats* (solution is bounded)

(7) n^{th} Order Linear Homogeneous Equations With Constant Coefficients

$$a_0 y^{(n)} + a_1 y^{(n-1)} + \cdots + a_{n-1} y' + a_n y = 0 \quad (*)$$

This differential equation has n independent solutions.

Characteristic Equation : $a_0 r^n + a_1 r^{n-1} + \cdots + a_{n-1} r + a_n = 0$ will have n characteristic roots that may be real and distinct, repeated, complex, or complex and repeated.

- (a) For each real root r that is not repeated \Rightarrow get a solution of (*): e^{rt}
 (b) For each real root r that is repeated $\underline{\mathbf{m}}$ times \Rightarrow get $\underline{\mathbf{m}}$ independent solutions of (*):

$$e^{rt}, te^{rt}, t^2e^{rt}, \dots, t^{m-1}e^{rt}$$

- (c) For each complex root $r = \lambda + i\mu$ repeated $\underline{\mathbf{m}}$ times \Rightarrow get $\underline{\mathbf{2m}}$ solutions of (*):

$$e^{\lambda t} \cos \mu t, te^{\lambda t} \cos \mu t, \dots, t^{m-1}e^{\lambda t} \cos \mu t \quad \underline{\text{and}} \quad e^{\lambda t} \sin \mu t, te^{\lambda t} \sin \mu t, \dots, t^{m-1}e^{\lambda t} \sin \mu t$$

(don't need to consider its conjugate root $\lambda - i\mu$)

(8) Undetermined Coefficients for n^{th} Order Linear Equations

This can only be used to find $y_p(t)$ of $a_0y^{(n)} + a_1y^{(n-1)} + \dots + a_{n-1}y' + a_ny = g(t)$ and $g(t)$ one of the 3 very SPECIAL FORMS in table in (5) above. The particular solution has the same form as before: $y_p(t) = t^s[\dots]$, where s = the smallest nonnegative integer such that no term of $y_p(t)$ is a term of $y_c(t)$, except this time $s = 0, 1, 2, \dots, n$.

(9) Laplace Transforms

- (a) Be able to compute Laplace transforms using definition:

$$\mathcal{L}\{f(t)\} = F(s) = \int_0^\infty e^{-st} f(t) dt$$

and using a table of Laplace transforms (see table on page 317) and using linearity: $\mathcal{L}\{f(t) + g(t)\} = \mathcal{L}\{f(t)\} + \mathcal{L}\{g(t)\}$, $\mathcal{L}\{cf(t)\} = c\mathcal{L}\{f(t)\}$.

- (b) Computing Inverse Laplace Transforms: Must be able to use a table of Laplace transforms usually together with *Partial Fractions* or *Completing the Square*, to find inverse Laplace transforms: $f(t) = \mathcal{L}^{-1}\{F(s)\}$.

- (c) Solving Initial Value Problems: Recall that

$$\mathcal{L}\{y'\} = s\mathcal{L}\{y\} - y(0)$$


$$\mathcal{L}\{y''\} = s^2\mathcal{L}\{y\} - sy(0) - y'(0)$$

$$\mathcal{L}\{y'''\} = s^3\mathcal{L}\{y\} - s^2y(0) - sy'(0) - y''(0)$$

$$\vdots$$


- (d) Discontinuous Functions:

- (i) Unit Step Function (Heaviside Function): If $c \geq 0$, $u_c(t) = \begin{cases} 0, & t < c \\ 1, & t \geq c \end{cases}$




$$\mathcal{L}\{u_c(t)\} = \frac{e^{-cs}}{s}$$

(ii) **Unit “Pulse” Function** : $u_a(t) - u_b(t) = \begin{cases} 1, & a \leq t < b \\ 0, & \text{otherwise} \end{cases}$



(iii) **Translated Functions** : $y = g(t) = \begin{cases} 0, & t < c \\ f(t - c), & t \geq c \end{cases} = u_c(t) f(t - c).$



$$\mathcal{L}\{u_c(t) f(t - c)\} = e^{-cs} F(s), \text{ where } F(s) = \mathcal{L}\{f(t)\}$$

Thus,

$$\mathcal{L}^{-1}\{e^{-cs} F(s)\} = u_c(t) f(t - c), \quad \text{where } f(t) = \mathcal{L}^{-1}\{F(s)\}$$

A useful formula **NOT** in the book :

$$\mathcal{L}\{u_c(t) h(t)\} = e^{-cs} \mathcal{L}\{h(t + c)\}$$

(iv) **Unit Impulse Functions** : If $y = \delta(t - c)$ ($c \geq 0$), then

$$\mathcal{L}\{\delta(t - c)\} = e^{-cs}$$

(e) **Convolutions**:

$$\mathcal{L}\{(f * g)(t)\} = \mathcal{L}\left\{\int_0^t f(t - \tau) g(\tau) d\tau\right\} = \mathcal{L}\{f(t)\} \mathcal{L}\{g(t)\}$$

(10) Systems of Linear Differential Equations : $\mathbf{x}'(t) = \mathbf{A} \mathbf{x}(t)$

- (a) Rewrite a single n^{th} order equation $p_0(t)y^{(n)} + p_1(t)y^{(n-1)} + \dots + p_n(t)y = g(t)$ as a system of 1^{st} order equations. Use the substitution :

$$\text{Let } \begin{matrix} \mathbf{x}_1 = y \\ \mathbf{x}_2 = y' \\ \vdots \\ \mathbf{x}_n = y^{(n-1)} \end{matrix} \quad \text{to get } 1^{st} \text{ Order System : } \begin{cases} x'_1 = x_2 \\ x'_2 = x_3 \\ \vdots \\ x'_{n-1} = x_n \\ x'_n = \frac{1}{p_0} \{-p_n x_1 - p_{n-1} x_2 - \dots - p_1 x_n + g(t)\} \end{cases}$$

- (b) **Existence & Uniqueness Theorem for Systems**. If $\mathbf{P}(t)$ and $\mathbf{g}(t)$ are continuous on an interval $\alpha < t < \beta$ containing t_0 , then the IVP $\begin{cases} \mathbf{x}'(t) = \mathbf{P}(t) \mathbf{x}(t) \\ \mathbf{x}(t_0) = \mathbf{x}_0 \end{cases}$ has a unique solution $\mathbf{x}(t)$ defined on the interval $\alpha < t < \beta$.
- (c) The set of vectors $\{\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(m)}\}$ is *linearly independent* if the equation

$$k_1 \mathbf{x}^{(1)} + k_2 \mathbf{x}^{(2)} + \dots + k_m \mathbf{x}^{(m)} = \mathbf{0}$$

is satisfied only for $k_1 = k_2 = \dots = k_m = 0$. This means you cannot write any one of these vectors as a linear combination of the others.

- (d) Solve 2×2 systems of 1^{st} order equations $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}' = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$ i.e., $\mathbf{x}' = A \mathbf{x}$ using :
- (i) **Elimination Method** : Basic idea - eliminate one of the unknowns (either x_1 or x_2) from the original system to get an equivalent single 2^{nd} order differential equation.
- (ii) **Eigenvalues & Eigenvectors Method** : See (11) below for solutions via this method and corresponding phase portraits.

Eigenvalue : If $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, then the eigenvalues of A are the roots of

$$|A - \lambda I| = \begin{vmatrix} (a - \lambda) & b \\ c & (d - \lambda) \end{vmatrix} = 0$$

Eigenvector : $\vec{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \neq \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ is a nonzero solution to $(A - \lambda I) \vec{v} = \vec{0}$.

- (e) If $\mathbf{x}^{(1)}(t) = \begin{pmatrix} x_{11}(t) \\ x_{21}(t) \end{pmatrix}$ and if $\mathbf{x}^{(2)}(t) = \begin{pmatrix} x_{12}(t) \\ x_{22}(t) \end{pmatrix}$, then the *Wronskian* is

$$W[\mathbf{x}^{(1)}, \mathbf{x}^{(2)}] = \begin{vmatrix} x_{11}(t) & x_{12}(t) \\ x_{21}(t) & x_{22}(t) \end{vmatrix}.$$

If $\mathbf{x}^{(1)}(t)$ and $\mathbf{x}^{(2)}(t)$ are solutions of $\mathbf{x}' = \mathbf{A} \mathbf{x}$ and $W[\mathbf{x}^{(1)}, \mathbf{x}^{(2)}](t_1) \neq 0$, then the set $\{\mathbf{x}^{(1)}(t), \mathbf{x}^{(2)}(t)\}$ forms a *Fundamental Set of Solutions* of the system and a *Fundamental Matrix* is

$$\Phi(t) = \begin{pmatrix} x_{11}(t) & x_{12}(t) \\ x_{21}(t) & x_{22}(t) \end{pmatrix}.$$

(11) Eigenvalue & Eigenvector Method and Phase Portraits : $\mathbf{x}' = \mathbf{A} \mathbf{x}$




The following describes how to find the general solution to (*) and plot solutions (trajectories). A plot of the trajectories of a given homogeneous system

$$\mathbf{x}' = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mathbf{x} \quad (*)$$

is called a **phase portrait**. To sketch the phase portrait, we need to find the corresponding eigenvalues and eigenvectors of the matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and then consider 3 cases :

- (a) $\lambda_1 < \lambda_2$, real and distinct : If $\mathbf{v}^{(1)}, \mathbf{v}^{(2)}$ are e-vectors corresponding to λ_1 and λ_2 , respectively $\Rightarrow \mathbf{x}^{(1)}(t) = e^{\lambda_1 t} \mathbf{v}^{(1)}$ and $\mathbf{x}^{(2)}(t) = e^{\lambda_2 t} \mathbf{v}^{(2)}$ are solutions and hence general solution of (*) is $\mathbf{x}(t) = C_1 \mathbf{x}^{(1)}(t) + C_2 \mathbf{x}^{(2)}(t)$ and hence if $\lambda_1 < \lambda_2$:

$$\mathbf{x}(t) = \underbrace{C_1 e^{\lambda_1 t} \mathbf{v}^{(1)}}_{\substack{\text{dominates} \\ \text{as } t \rightarrow -\infty}} + \underbrace{C_2 e^{\lambda_2 t} \mathbf{v}^{(2)}}_{\substack{\text{dominates} \\ \text{as } t \rightarrow \infty}}$$




(b) $\lambda_1 = \alpha + i\beta$: If $\mathbf{w} = \mathbf{a} + i\mathbf{b}$ is a complex e-vector corresponding to λ_1 then \Rightarrow

$$\mathbf{x}^{(1)}(t) = \Re \left\{ e^{\lambda_1 t} \mathbf{w} \right\} = e^{\alpha t} (\mathbf{a} \cos \beta t - \mathbf{b} \sin \beta t) \quad \text{and}$$

$\mathbf{x}^{(2)}(t) = \Im \left\{ e^{\lambda_1 t} \mathbf{w} \right\} = e^{\alpha t} (\mathbf{a} \sin \beta t + \mathbf{b} \cos \beta t)$ are real-valued solutions and hence general solution of (*) is $\mathbf{x}(t) = C_1 \mathbf{x}^{(1)}(t) + C_2 \mathbf{x}^{(2)}(t)$.

If say $\alpha < 0$:



(Test a point to decide which)

(c) $\lambda_1 = \lambda_2$: If there is only *one* linearly independent eigenvector corresponding to λ_1 , then solutions to $\mathbf{x}' = \mathbf{A} \mathbf{x}$ are $\mathbf{x}^{(1)}(t) = e^{\lambda_1 t} \mathbf{v}$ and $\mathbf{x}^{(2)}(t) = t e^{\lambda_1 t} \mathbf{v} + e^{\lambda_1 t} \mathbf{a}$, where

$$(\mathbf{A} - \lambda_1 I) \mathbf{v} = \mathbf{0}$$


$$(\mathbf{A} - \lambda_1 I) \mathbf{a} = \mathbf{v}$$

(\mathbf{v} is an eigenvector of \mathbf{A} , while \mathbf{a} is called a “generalized eigenvector” of \mathbf{A})

The general solution of the system (*) is $\mathbf{x}(t) = C_1 \mathbf{x}^{(1)}(t) + C_2 \mathbf{x}^{(2)}(t)$ and hence:

$$\mathbf{x}(t) = C_1 e^{\lambda_1 t} \mathbf{v} + C_2 \left[\underbrace{t e^{\lambda_1 t} \mathbf{v}}_{\substack{\text{dominates} \\ \text{as } t \rightarrow \pm\infty}} + e^{\lambda_1 t} \mathbf{a} \right]$$

If say $\lambda_1 < 0$:



(Test a point to decide which)

(12) Particular Solutions to Nonhomogeneous Linear Systems :

$$\mathbf{x}' = \mathbf{A} \mathbf{x} + \mathbf{g}(t)$$

- (a) Undetermined Coefficients for Systems The column vector $\vec{\mathbf{g}}(t) = \begin{pmatrix} g_1(t) \\ g_2(t) \end{pmatrix}$ must have each component function $g_1(t)$ and $g_2(t)$ as one of the three special forms like those for Undetermined Coefficients for regular 2^{nd} order equations and \mathbf{A} must be a constant matrix. The main difference is if say $\mathbf{g}(t) = \mathbf{u} e^{\lambda t}$ and λ is also an eigenvalue of \mathbf{A} , then try a particular solution of the form $\mathbf{x}_p = \mathbf{a} t e^{\lambda t} + \mathbf{b} e^{\lambda t}$.
- (b) Variation of Parameters for Systems : $\mathbf{x}' = \mathbf{A}(t) \mathbf{x} + \mathbf{g}(t)$:

$$\mathbf{x}_p(t) = \Phi(t) \int \Phi^{-1}(t) \mathbf{g}(t) dt,$$

where $\Phi(t)$ is a Fundamental Matrix of the homogeneous system $\mathbf{x}' = \mathbf{A}(t) \mathbf{x} + \mathbf{g}(t)$ can have any form and \mathbf{A} need not be a constant matrix.

PRACTICE PROBLEMS

- [1] For what value of α will the solution to the IVP $\begin{cases} y'' - y' - 2y = 0 \\ y(0) = \alpha \\ y'(0) = 2 \end{cases}$ satisfy $y \rightarrow 0$ as $t \rightarrow \infty$?
- [2] (a) Show that $y_1 = x$ and $y_2 = x^{-1}$ are solutions of the differential equation $x^2 y'' + xy' - y = 0$.
(b) Evaluate the Wronskian $W(y_2, y_1)$ at $x = \frac{1}{2}$.
(c) Find the solution of the initial value problem $x^2 y'' + xy' - y = 0$, $y(1) = 2$, $y'(1) = 4$.
- [3] Find the largest open interval for which the initial value problem $3x^2 y'' + y' + \frac{1}{x-2}y = \frac{1}{x-3}$, $y(1) = 3$, $y'(1) = 2$, has a solution.

In Problems 4, 5, and 6 find the general solution of the homogeneous differential equations in (a) and use the method of **Undetermined Coefficients** to find a particular solution y_p in (b) and find the FORM of a particular solution (c).

- [4] (a) $y'' - 5y' + 6y = 0$ (b) $y'' - 5y' + 6y = t^2$ (c) $y'' - 5y' + 6y = e^{2t} + \cos(3t)$
- [5] (a) $y'' - 6y' + 9y = 0$ (b) $y'' - 6y' + 9y = te^{3t}$ (c) $y'' - 6y' + 9y = e^t + \cos(3t)$
- [6] (a) $y'' - 2y' + 10y = 0$ (b) $y'' - 2y' + 10y = e^x + \cos(3x)$ (c) $y'' - 2y' + 10y = e^x \cos(3x)$

- [7] Find the general solution to (a) $y'' + y' - 6y = 7e^{4t}$ (b) $y'' + y' - 6y = 7e^{4t} - 100 \sin t$

- [8] Solve this IVP: $y'' - y' = 4t$, $y(0) = 0$, $y'(0) = 0$.

- [9] Find the general solution to $y'' + y = \tan t$, $0 < x < \frac{\pi}{2}$.

- [10] The differential equation $x^2 y'' - 2xy' + 2y = 0$ has solutions $y_1(x) = x$ and $y_2 = x^2$. Use the method of **Variation of Parameters** to find a solution of $x^2 y'' - 2xy' + 2y = 2x^2$.

- [11] The differential equation $x^2 y'' + xy' - y = 0$ has one solution $y_1(x) = x$. Use the method of **Reduction of Order** to find a second (linearly independent) solution of $x^2 y'' + xy' - y = 0$.

- [12] For what nonnegative values of γ will the the solution of the initial value problem $u'' + \gamma u' + 4u = 0$, $u(0) = 4$, $u'(0) = 0$ oscillate?

- [13] (a) For what positive values of k does the solution of the initial value problem $2u'' + ku = 3 \cos(2t)$, $u(0) = 0$, $u'(0) = 0$, become *unbounded* (Resonance)?

- (b) For what positive values of k does the solution of the initial value problem $2u'' + u' + ku = 3\cos(2t)$, $u(0) = 0$, $u'(0) = 0$, become *unbounded* (Resonance) ?

[14] Find the steady-state solution of the IVP $y'' + 4y' + 4y = \sin t$, $y(0) = 0$, $y'(0) = 0$.

[15] A 4-kg mass stretches a spring 0.392 m. If the mass is released from 1 m below the equilibrium position with a downward velocity of 10 m/sec, what is the maximum displacement ?

In Problems 16 and 17 find the general solution of the homogeneous differential equations in (a) and use the method of **Undetermined Coefficients** to find the FORM of a particular solution of the nonhomogeneous equation in (b).

[16] (a) $y''' - y' = 0$ (b) $y''' - y' = t + e^t$

[17] (a) $y''' - y'' - y' + y = 0$ (b) $y''' - y'' - y' + y = e^t + \cos t$

[18] Find the solution of the initial value problem $y''' - 2y'' + y' = 0$, $y(0) = 2$, $y'(0) = 0$, $y''(0) = 1$.

[19] Find the general solution of the differential equation $y''' + y' = t^2$.

[20] Find the general solution of $y'' + 4y' = -10\cos 2t$.

[21] Find a fundamental set of solutions of $y^{(5)} - 4y''' = 0$.

[22] Find the Laplace transform of these functions:

(a) $f(t) = 3 - e^{2t}$ (b) $g(t) = 100t^5$ (c) $h(t) = \cosh \pi t$ (d) $k(t) = -10t^3e^{5t}$

[23] Find the inverse Laplace transform of


(a) $F(s) = \frac{9}{s^2 - s - 2}$ (b) $F(s) = \frac{s}{(s-1)^2}$ (c) $F(s) = \frac{8}{(s+1)^4}$ (d) $F(s) = \frac{3s+2}{s^2 + 2s + 5}$

[24] Solve these initial value problems: (a) $\begin{cases} y'' - y' - 6y = 0 \\ y(0) = 1 \\ y'(0) = -1 \end{cases}$ (b) $\begin{cases} y'' - 2y' + 2y = \cos t \\ y(0) = 1 \\ y'(0) = 0 \end{cases}$

(c) $y'' - y = \begin{cases} 1, & t < 5 \\ 2, & 5 \leq t < \infty \end{cases}$; $y(0) = y'(0) = 0$.

(d) $y'' + 4y = \begin{cases} t, & t < 1 \\ 0, & 1 < t < \infty \end{cases}$; $y(0) = y'(0) = 0$.

(e) $y' + y = g(t)$, $y(0) = 0$ and where $g(t)$:



(f) $y'' + 4y = \delta(t-3)$, $y(0) = y'(0) = 0$

[25] $\mathcal{L} \left\{ \int_0^t 100 e^{-2\tau} \cos \pi(t-\tau) d\tau \right\} = ?$

[26] If $g(t) = \mathcal{L}^{-1}\{G(s)\}$, then $\mathcal{L}^{-1} \left\{ \frac{G(s)}{(s-3)^2} \right\} = ?$

[27] Use the Elimination Method to solve the system $\begin{cases} x_1' = x_1 + x_2 \\ x_2' = 4x_1 + x_2 \end{cases}$

[28] Rewrite the 2^{nd} order differential equation $y'' + 2y' + 3ty = \cos t$ with $y(0) = 1$, $y'(0) = 4$ as a system of 1^{st} order differential equations.

[29] Find eigenvalues and corresponding eigenvectors of (a) $A = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix}$ (b) $A = \begin{pmatrix} -2 & 0 \\ 1 & -1 \end{pmatrix}$

[30] Find the solution of the IVP $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}' = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad \begin{pmatrix} x_1(0) \\ x_2(0) \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}.$

Find a fundamental matrix $\Phi(t)$.

[31] Solve $\begin{pmatrix} x_1' \\ x_2' \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad \vec{x}(0) = \begin{pmatrix} -1 \\ 2 \end{pmatrix}.$

[32] Find the general solution of the system $\vec{x}'(t) = A\vec{x}(t)$, where $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$

[33] Tank # 1 initially holds 50 gals of brine with a concentration of 1 lb/gal, while Tank # 2 initially holds 25 gals of brine with a concentration of 3 lb/gal. Pure H_2O flows into Tank # 1 at 5 gal/min. The well-stirred solution from Tank # 1 then flows into Tank # 2 at 5 gal/min. The solution in Tank # 2 flows out at 5 gal/min. Set up and solve an IVP that gives $x_1(t)$ and $x_2(t)$, the amount of salt in Tanks # 1 and # 2, respectively, at time t .

[34] Tank # 1 initially holds 50 gals of brine with concentration of 1 lb/gal and Tank # 2 initially holds 25 gals of brine with concentration 3 lb/gal. The solution in Tank # 1 flows at 5 gal/min into Tank # 2, while the solution in Tank # 2 flows back into Tank # 1 at 5 gal/min. Set up an IVP that gives $x_1(t)$ and $x_2(t)$, the amount of salt in Tanks # 1 and # 2, respectively, at time t .

[35] Find the general solution of $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}' = \begin{pmatrix} -2 & 0 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} 3 \\ 1 \end{pmatrix} e^t.$

[36] Find a particular solution of $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}' = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} - \begin{pmatrix} 2 \\ 3 \end{pmatrix}.$

[37] Find the general solution of $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}' = \begin{pmatrix} 2 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} - \begin{pmatrix} 6e^{-t} \\ 1 \end{pmatrix}.$


[38] Match the phase portraits shown below that best corresponds to each of the given systems of differential equations:

(i) $\vec{x}' = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \vec{x}$; Solution : $\vec{x}(t) = C_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} e^t + C_2 \begin{pmatrix} 1 \\ -1 \end{pmatrix} e^{-t}$


(ii) $\vec{x}' = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \vec{x}$; Solution : $\vec{x}(t) = C_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} e^t + C_2 \begin{pmatrix} 1 \\ -1 \end{pmatrix} e^{3t}$

(iii) $\vec{x}' = \begin{pmatrix} 2 & -1 \\ 1 & 0 \end{pmatrix} \vec{x}$; Solution : $\vec{x}(t) = C_1 \begin{pmatrix} 2 \\ 2 \end{pmatrix} e^t + C_2 e^t \left\{ \begin{pmatrix} 2 \\ 2 \end{pmatrix} t + \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\}$



(iv) $\vec{x}' = \begin{pmatrix} -1 & 1 \\ -1 & -1 \end{pmatrix} \vec{x}$; Solution : $\vec{x}(t) = C_1 \begin{pmatrix} \cos t \\ -\sin t \end{pmatrix} e^{-t} + C_2 \begin{pmatrix} \sin t \\ \cos t \end{pmatrix} e^{-t}$





(A)





(B)





(C)





(D)





(E)



(F)



(G)



(H)

ANSWERS

- [1] $\alpha = -2$ [2] (b) $W(x^{-1}, x)(\frac{1}{2}) = 4$; (c) $y = 3x - x^{-1}$ [3] $0 < x < 2$
- [4] (a) $y = C_1 e^{2t} + C_2 e^{3t}$ (b) $y = At^2 + Bt + C$ (c) $y = Ate^{2t} + B \cos(3t) + C \sin(3t)$
- [5] (a) $y = C_1 e^{3t} + C_2 te^{3t}$ (b) $y = t^2(At + B)e^{3t}$ (c) $y = Ae^t + B \cos(3t) + C \sin(3t)$
- [6] (a) $y = C_1 e^x \cos(3x) + C_2 e^x \sin(3x)$ (b) $y = Ae^x + B \cos(3x) + C \sin(3x)$
(c) $y = x(A \cos(3x) + B \sin(3x))e^x$
- [7] (a) $y = C_1 e^{-3t} + C_2 e^{2t} + \frac{1}{2}e^{4t}$ (b) $y = C_1 e^{-3t} + C_2 e^{2t} + \frac{1}{2}e^{4t} + 2 \cos t + 14 \sin t$
- [8] $y = -4 + 4e^t - 2t^2 - 4t$
- [9] $y = C_1 \cos t + C_2 \sin t - (\cos t) \ln(\sec t + \tan t)$
- [10] $y = 2x^2 \ln x$ or $y = 2x^2 \ln x + (C_1 x + C_2 x^2)$
- [11] $y = x^{-1}$ or $y = Ax^{-1} + Bx$, $A \neq 0$
- [12] $0 \leq \gamma < 4$
- [13] (a) $k = 8$ (resonance) (b) NO value of k , all solutions are bounded.
- [14] $y = \frac{1}{25}(3 \sin t - 4 \cos t)$
- [15] $u(t) = \cos 5t + 2 \sin 5t = \sqrt{5} \cos(5t - \delta)$, $\delta = \tan^{-1} 2 \approx 1.1$ Thus amplitude $= \sqrt{5}$.
- [16] (a) $y = C_1 + C_2 e^{-t} + C_3 e^t$ (b) $y = t(At + B) + Cte^t$
- [17] (a) $y = C_1 e^t + C_2 te^t + C_3 e^{-t}$ (b) $y = At^2 e^t + B \cos t + C \sin t$
- [18] $y = 3 - e^t + te^t$
- [19] $y = C_1 + C_2 \cos t + C_3 \sin t + \frac{1}{3}t^3 - 2t$
- [20] $y = C_1 + C_2 e^{-4t} + (\frac{1}{2} \cos 2t - \sin 2t)$
- [21] $\{1, t, t^2, e^{2t}, e^{-2t}\}$
- [22] (a) $\frac{2s-6}{s^2-2s}$ (b) $\frac{12000}{s^6}$ (c) $\frac{s}{s^2-\pi^2}$ (d) $-\frac{60}{(s-5)^4}$
- [23] (a) $3(e^{2t} - e^{-t})$ (b) $e^t + te^t$ (c) $\frac{4}{3}t^3 e^{-t}$ (d) $3e^{-t} \cos 2t - \frac{1}{2}e^{-t} \sin 2t$
- [24] (a) $y = \frac{1}{5}(e^{3t} + 4e^{-2t})$ (b) $y = \frac{1}{5}(\cos t - 2 \sin t + 4e^t \cos t - 2e^t \sin t)$
(c) $y = -1 + \frac{1}{2}(e^t + e^{-t}) + u_5(t)(-1 + \frac{1}{2}(e^{(t-5)} + e^{-(t-5)}))$,
or $y = -1 + \cosh t + u_5(t)(-1 + \cosh(t-5))$
(d) $y = (-\frac{1}{8} \sin 2t + \frac{t}{4}) - u_1(t)(-\frac{1}{8} \sin 2(t-1) + \frac{t-1}{4}) - u_1(t)(\frac{1}{4} - \frac{1}{4} \cos 2(t-1))$
(e) $y = 3(1 - e^{-t}) - 3u_2(t)(1 - e^{-(t-2)}) + 3u_4(t)(1 - e^{-(t-4)})$ (f) $y = \frac{1}{2}u_3(t)(t) \sin 2(t-3)$
- [25] $\frac{100s}{(s+2)(s^2+\pi^2)}$ [26] $\int_0^t (t-\tau) e^{3(t-\tau)} g(\tau) d\tau$ or $\int_0^t \tau e^{3\tau} g(t-\tau) d\tau$
- [27] $x_1(t) = C_1 e^{3t} + C_2 e^{-t}$, $x_2(t) = 2C_1 e^{3t} - 2C_2 e^{-t}$
- [28] Let $x_1 = y$, $x_2 = y'$, then $\begin{cases} x'_1 = x_2 \\ x'_2 = -3tx_1 - 2x_2 + \cos t \end{cases}$, where $x_1(0) = 1$, $x_2(0) = 4$
- [29] (a) $\lambda_1 = 3$, $\mathbf{v}^{(1)} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$; $\lambda_2 = -1$, $\mathbf{v}^{(2)} = \begin{pmatrix} 1 \\ -2 \end{pmatrix}$
- [29] (b) $\lambda_1 = -1$, $\mathbf{v}^{(1)} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$; $\lambda_2 = -2$, $\mathbf{v}^{(2)} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$
- [30] $\mathbf{x}(t) = 2e^{3t} \begin{pmatrix} 1 \\ 2 \end{pmatrix} + e^{-t} \begin{pmatrix} 1 \\ -2 \end{pmatrix}$, $\Phi(t) = \begin{pmatrix} e^{3t} & e^{-t} \\ 2e^{3t} & -2e^{-t} \end{pmatrix}$
- [31] $\mathbf{x}(t) = 2e^t \begin{pmatrix} \sin t \\ \cos t \end{pmatrix} - e^t \begin{pmatrix} \cos t \\ -\sin t \end{pmatrix}$ [32] $\mathbf{x}(t) = C_1 e^t \begin{pmatrix} 1 \\ 0 \end{pmatrix} + C_2 \left\{ e^t \begin{pmatrix} 0 \\ 1 \end{pmatrix} + te^t \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\}$
- [33] $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix}' = \begin{pmatrix} -\frac{1}{10} & 0 \\ \frac{1}{10} & -\frac{1}{5} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$, $\begin{pmatrix} x_1(0) \\ x_2(0) \end{pmatrix} = \begin{pmatrix} 50 \\ 75 \end{pmatrix}$
- Solution: $\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = 50e^{-\frac{t}{10}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + 25e^{-\frac{t}{5}} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

$$[34] \quad \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}' = \begin{pmatrix} -\frac{1}{10} & \frac{1}{5} \\ \frac{1}{10} & -\frac{1}{5} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad \begin{pmatrix} x_1(0) \\ x_2(0) \end{pmatrix} = \begin{pmatrix} 50 \\ 75 \end{pmatrix}$$

$$\text{Solution : } \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \frac{125}{3} \begin{pmatrix} 2 \\ 1 \end{pmatrix} - \frac{100}{3} e^{-\frac{3t}{10}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

$$[35] \quad \mathbf{x}(t) = C_1 e^{-t} \begin{pmatrix} 0 \\ 1 \end{pmatrix} + C_2 e^{-2t} \begin{pmatrix} 1 \\ -1 \end{pmatrix} + e^t \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad [36] \quad \mathbf{x}_p(t) = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$$

$$[37] \quad \mathbf{x}(t) = C_1 e^t \begin{pmatrix} 0 \\ 1 \end{pmatrix} + C_2 e^{2t} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + e^{-t} \begin{pmatrix} 2 \\ -1 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$[38] \quad \text{(i) } \mathbf{C} \quad \text{(ii) } \mathbf{A} \quad \text{(iii) } \mathbf{B} \quad \text{(iv) } \mathbf{D}$$

$f(t) = \mathcal{L}^{-1}\{F(s)\}$		$F(s) = \mathcal{L}\{f(t)\}$
1.	1	$\frac{1}{s}$
2.	e^{at}	$\frac{1}{s-a}$
3.	t^n	$\frac{n!}{s^{n+1}}$
4.	$t^p \ (p > -1)$	$\frac{\Gamma(p+1)}{s^{p+1}}$
5.	$\sin at$	$\frac{a}{s^2 + a^2}$
6.	$\cos at$	$\frac{s}{s^2 + a^2}$
7.	$\sinh at$	$\frac{a}{s^2 - a^2}$
8.	$\cosh at$	$\frac{s}{s^2 - a^2}$
9.	$e^{at} \sin bt$	$\frac{b}{(s-a)^2 + b^2}$
10.	$e^{at} \cos bt$	$\frac{s-a}{(s-a)^2 + b^2}$
11.	$t^n e^{at}$	$\frac{n!}{(s-a)^{n+1}}$
11.	$t^n e^{at}$	$\frac{n!}{(s-a)^{n+1}}$
12.	$u_c(t)$	$\frac{e^{-cs}}{s}$
13.	$u_c(t)f(t-c)$	$e^{-cs}F(s)$
14.	$e^{ct}f(t)$	$F(s-c)$
15.	$f(ct)$	$\frac{1}{c}F\left(\frac{s}{c}\right), \ c > 0$
16.	$\int_0^t f(t-\tau)g(\tau) d\tau$	$F(s)G(s)$
17.	$\delta(t-c)$	e^{-cs}
18.	$f^{(n)}(t)$	$s^n F(s) - s^{n-1}f(0) - \dots - sf^{(n-2)}(0) - f^{(n-1)}(0)$
19.	$(-t)^n f(t)$	$F^{(n)}(s)$