



Continuous fields of C^* -algebras over finite dimensional spaces

Marius Dadarlat¹

Department of Mathematics, Purdue University, West Lafayette, IN 47907, USA

Received 10 July 2007; accepted 24 June 2009

Available online 17 July 2009

Communicated by Dan Voiculescu

Abstract

Let X be a finite dimensional compact metrizable space. We study a technique which employs semiprojectivity as a tool to produce approximations of $C(X)$ -algebras by $C(X)$ -subalgebras with controlled complexity. The following applications are given. All unital separable continuous fields of C^* -algebras over X with fibers isomorphic to a fixed Cuntz algebra \mathcal{O}_n , $n \in \{2, 3, \dots, \infty\}$, are locally trivial. They are trivial if $n = 2$ or $n = \infty$. For $n \geq 3$ finite, such a field is trivial if and only if $(n - 1)[1_A] = 0$ in $K_0(A)$, where A is the C^* -algebra of continuous sections of the field. We give a complete list of the Kirchberg algebras D satisfying the UCT and having finitely generated K -theory groups for which every unital separable continuous field over X with fibers isomorphic to D is automatically locally trivial or trivial. In a more general context, we show that a separable unital continuous field over X with fibers isomorphic to a KK -semiprojective Kirchberg C^* -algebra is trivial if and only if it satisfies a K -theoretical Fell type condition.

© 2009 Elsevier Inc. All rights reserved.

Keywords: C^* -algebras; Continuous fields; K -theory; Kirchberg algebras

Contents

1. Introduction	1851
2. $C(X)$ -algebras	1853

E-mail address: mdd@math.purdue.edu.

¹ The author was partially supported by NSF grant #DMS-0500693.

3. Semiprojectivity	1858
4. Approximation of $C(X)$ -algebras	1862
5. Representing $C(X)$ -algebras as inductive limits	1868
6. When is a fibered product locally trivial	1870
7. When is a $C(X)$ -algebra locally trivial	1874
References	1880

1. Introduction

The Gelfand–Naimark characterization of commutative C^* -algebras has suggested the problem of representing noncommutative C^* -algebras as sections of bundles. By a result of Fell [15], if the primitive spectrum X of a separable C^* -algebra A is Hausdorff, then A is isomorphic to the C^* -algebra of continuous sections vanishing at infinity of a continuous field of simple C^* -algebras over X . In particular A is a continuous $C(X)$ -algebra in the sense of Kasparov [18]. This description is very satisfactory, since as explained in [4], the continuous fields of C^* -algebras are in natural correspondence with the bundles of C^* -algebras in the sense of topology. Nevertheless, only a tiny fraction of the continuous fields of C^* -algebras corresponds to locally trivial bundles.

In this paper we prove automatic and conditional local/global trivialization results for continuous fields of Kirchberg algebras. By a Kirchberg algebra we mean a purely infinite simple nuclear separable C^* -algebra [29]. Notable examples include the simple Cuntz–Krieger algebras [8]. The following theorem illustrates our results.

Theorem 1.1. *A separable unital $C(X)$ -algebra A over a finite dimensional compact Hausdorff space X all of whose fibers are isomorphic to the same Cuntz algebra \mathcal{O}_n , $n \in \{2, 3, \dots, \infty\}$, is locally trivial. If $n = 2$ or $n = \infty$, then $A \cong C(X) \otimes \mathcal{O}_n$. If $3 \leq n < \infty$, then A is isomorphic to $C(X) \otimes \mathcal{O}_n$ if and only if $(n - 1)[1_A] = 0$ in $K_0(A)$.*

The case $X = [0, 1]$ of Theorem 1.1 was proved in a joint paper with G. Elliott [10]. We parametrize the homotopy classes

$$[X, \text{Aut}(\mathcal{O}_n)] \cong \begin{cases} K_1(C(X) \otimes \mathcal{O}_n) & \text{if } 3 \leq n < \infty, \\ \{*\} & \text{if } n = 2, \infty \end{cases}$$

(see Theorem 7.4), and hence classify the unital separable $C(SX)$ -algebras A with fiber \mathcal{O}_n over the suspension SX of a finite dimensional metrizable Hausdorff space X .

To put our results in perspective, let us recall that none of the general basic properties of a continuous field implies any kind of local triviality. An example of a continuous field of Kirchberg algebras over $[0, 1]$ which is not locally trivial at any point even though all of its fibers are mutually isomorphic is exhibited in [10, Ex. 8.4]. Examples of nonexact continuous fields with similar properties were found by S. Wassermann [37].

A separable C^* -algebra D is KK -semiprojective if the functor $KK(D, -)$ is continuous, see Section 3. The class of KK -semiprojective C^* -algebras includes the nuclear semiprojective C^* -algebras and also the C^* -algebras which satisfy the Universal Coefficient Theorem in KK -theory (abbreviated UCT [31]) and whose K -theory groups are finitely generated. It is very interesting

that the *only obstruction* to local or global triviality for a continuous field of Kirchberg algebras is of purely K-theoretical nature.

Theorem 1.2. *Let A be a separable C^* -algebra whose primitive spectrum X is compact Hausdorff and of finite dimension. Suppose that each primitive quotient $A(x)$ of A is nuclear, purely infinite and stable. Then A is isomorphic to $C(X) \otimes D$ for some KK -semiprojective stable Kirchberg algebra D if and only if there is $\sigma \in KK(D, A)$ such that $\sigma_x \in KK(D, A(x))^{-1}$ for all $x \in X$. For any such σ there is an isomorphism of $C(X)$ -algebras $\Phi : C(X) \otimes D \rightarrow A$ such that $KK(\Phi|_D) = \sigma$.*

We have an entirely similar result covering the unital case: Theorem 7.3. The required existence of σ is a KK -theoretical analog of the classical condition of Fell that appears in the trivialization theorem of Dixmier and Douady [12] of continuous fields with fibers isomorphic to the compact operators. An important feature of our condition is that it is a priori much weaker than the condition that A is $KK_{C(X)}$ -equivalent to $C(X) \otimes D$. In particular, we do not need to worry at all about the potentially hard issue of constructing elements in $KK_{C(X)}(A, C(X) \otimes D)$. To illustrate this point, let us note that it is almost trivial to verify that the local existence of σ is automatic for unital $C(X)$ -algebras with fiber \mathcal{O}_n and hence to derive Theorem 1.1. A C^* -algebra D has the *automatic local triviality property* if any separable $C(X)$ -algebra over a finite dimensional compact Hausdorff space X all of whose fibers are isomorphic to D is locally trivial. A unital C^* -algebra D has the *automatic local triviality property in the unital sense* if any separable unital $C(X)$ -algebra over a finite dimensional compact Hausdorff space X all of whose fibers are isomorphic to D is locally trivial. The *automatic triviality property* is defined similarly.

Theorem 1.3 (Automatic triviality). *A separable continuous $C(X)$ -algebra over a finite dimensional compact Hausdorff space X all of whose fibers are isomorphic to $\mathcal{O}_2 \otimes \mathcal{K}$ is isomorphic to $C(X) \otimes \mathcal{O}_2 \otimes \mathcal{K}$. The C^* -algebra $\mathcal{O}_2 \otimes \mathcal{K}$ is the only Kirchberg algebra satisfying the automatic local triviality property and hence the automatic triviality property.*

Theorem 1.4 (Automatic local triviality in the unital sense). *A unital KK -semiprojective Kirchberg algebra D has the automatic local triviality property in the unital sense if and only if all unital $*$ -endomorphisms of D are KK -equivalences. In that case, if A is a separable unital $C(X)$ -algebra over a finite dimensional compact Hausdorff space X all of whose fibers are isomorphic to D , then $A \cong C(X) \otimes D$ if and only if there is $\sigma \in KK(D, A)$ such that the induced homomorphism $K_0(\sigma) : K_0(D) \rightarrow K_0(A)$ maps $[1_D]$ to $[1_A]$.*

It is natural to ask if there are other unital Kirchberg algebras besides the Cuntz algebras which have the automatic local triviality property in the unital sense. Consider the following list \mathcal{G} of pointed abelian groups:

- (a) $(\{0\}, 0)$;
- (b) (\mathbb{Z}, k) with $k > 0$;
- (c) $(\mathbb{Z}/p^{e_1} \oplus \dots \oplus \mathbb{Z}/p^{e_n}, p^{s_1} \oplus \dots \oplus p^{s_n})$ where p is a prime, $n \geq 1$, $0 \leq s_i < e_i$ for $1 \leq i \leq n$ and $0 < s_{i+1} - s_i < e_{i+1} - e_i$ for $1 \leq i < n$. If $n = 1$ the latter condition is vacuous. Note that if the integers $1 \leq e_1 \leq \dots \leq e_n$ are given then there exists integers s_1, \dots, s_n satisfying the conditions above if and only if $e_{i+1} - e_i \geq 2$ for each $1 \leq i \leq n$. If that is the case one can choose $s_i = i - 1$ for $1 \leq i \leq n$.

- (d) $(G(p_1) \oplus \cdots \oplus G(p_m), g_1 \oplus \cdots \oplus g_m)$ where p_1, \dots, p_m are distinct primes and each $(G(p_j), g_j)$ is a pointed group as in (c).
- (e) $(\mathbb{Z} \oplus G(p_1) \oplus \cdots \oplus G(p_m), k \oplus g_1 \oplus \cdots \oplus g_m)$ where $(G(p_j), g_j)$ are as in (d). Moreover we require that $k > 0$ is divisible by $p_1^{s_{n(1)}+1} \cdots p_m^{s_{n(m)}+1}$ where $s_{n(j)}$ is defined as in (c) corresponding to the prime p_j .

Theorem 1.5 (*Automatic local triviality in the unital sense – the UCT case*). *Let D be a unital Kirchberg algebra which satisfies the UCT and has finitely generated K -theory groups. (i) D has the automatic triviality property in the unital sense if and only if D is isomorphic to either \mathcal{O}_2 or \mathcal{O}_∞ . (ii) D has the automatic local triviality property in the unital sense if and only if $K_1(D) = 0$ and $(K_0(D), [1_D])$ is isomorphic to one of the pointed groups from the list \mathcal{G} . (iii) If D is as in (ii), then a separable unital $C(X)$ -algebra A over a finite dimensional compact Hausdorff space X all of whose fibers are isomorphic to D is trivial if and only if there exists a homomorphism of groups $K_0(D) \rightarrow K_0(A)$ which maps $[1_D]$ to $[1_A]$.*

We use semiprojectivity (in various flavors) to approximate and represent continuous $C(X)$ -algebras as inductive limits of fibered products of n locally trivial $C(X)$ -subalgebras where $n \leq \dim(X) < \infty$. This clarifies the local structure of many $C(X)$ -algebras (see Theorem 5.2) and gives a new understanding of the K -theory of separable continuous $C(X)$ -algebras with arbitrary nuclear fibers.

A remarkable isomorphism result for separable nuclear strongly purely infinite stable C^* -algebras was announced (with an outline of the proof) by Kirchberg in [20]: two such C^* -algebras A and B with the same primitive spectrum X are isomorphic if and only if they are $KK_{C(X)}$ -equivalent. This is always the case after tensoring with \mathcal{O}_2 . However the problem of recognizing when A and B are $KK_{C(X)}$ -equivalent is open even for very simple spaces X such as the unit interval or non-Hausdorff spaces with more than two points.

The proof of Theorem 4.6 (one of our main results) generalizes and refines a technique that was pioneered for fields over zero dimensional spaces in joint work with Pasnicu [11] and for fields over an interval in joint work with G. Elliott [10]. We shall rely heavily on the classification theorem (and related results) of Kirchberg [19] and Phillips [28], and on the work on nonsimple nuclear purely infinite C^* -algebras of Blanchard and Kirchberg [5,4] and Kirchberg and Rørdam [21,22].

The author is grateful to E. Blanchard, L.G. Brown and N.C. Phillips for useful discussions and comments.

2. $C(X)$ -algebras

Let X be a locally compact Hausdorff space. A $C(X)$ -algebra is a C^* -algebra A endowed with a $*$ -homomorphism θ from $C_0(X)$ to the center $ZM(A)$ of the multiplier algebra $M(A)$ of A such that $C_0(X)A$ is dense in A ; see [18,3]. We write fa rather than $\theta(f)a$ for $f \in C_0(X)$ and $a \in A$. If $Y \subseteq X$ is a closed set, we let $C_0(X, Y)$ denote the ideal of $C_0(X)$ consisting of functions vanishing on Y . Then $C_0(X, Y)A$ is a closed two-sided ideal of A (by Cohen factorization). The quotient of A by this ideal is a $C(X)$ -algebra denoted by $A(Y)$ and is called the restriction of $A = A(X)$ to Y . The quotient map is denoted by $\pi_Y : A(X) \rightarrow A(Y)$. If Z is a closed subset of Y we have a natural restriction map $\pi_Z^Y : A(Y) \rightarrow A(Z)$ and $\pi_Z = \pi_Z^Y \circ \pi_Y$. If Y reduces to a point x , we write $A(x)$ for $A(\{x\})$ and π_x for $\pi_{\{x\}}$. The C^* -algebra $A(x)$ is called the fiber of A at x . The image $\pi_x(a) \in A(x)$ of $a \in A$ is denoted by $a(x)$. A morphism of $C(X)$ -algebras

$\eta : A \rightarrow B$ induces a morphism $\eta_Y : A(Y) \rightarrow B(Y)$. If $A(x) \neq 0$ for x in a dense subset of X , then θ is injective. If X is compact, then $\theta(1) = 1_{M(A)}$. Let A be a C^* -algebra, $a \in A$ and $\mathcal{F}, \mathcal{G} \subseteq A$. Throughout the paper we will assume that X is a compact Hausdorff space unless stated otherwise. If $\varepsilon > 0$, we write $a \in_\varepsilon \mathcal{F}$ if there is $b \in \mathcal{F}$ such that $\|a - b\| < \varepsilon$. Similarly, we write $\mathcal{F} \subset_\varepsilon \mathcal{G}$ if $a \in_\varepsilon \mathcal{G}$ for every $a \in \mathcal{F}$. The following lemma collects some basic properties of $C(X)$ -algebras.

Lemma 2.1. *Let A be a $C(X)$ -algebra and let $B \subset A$ be a $C(X)$ -subalgebra. Let $a \in A$ and let Y be a closed subset of X .*

- (i) *The map $x \mapsto \|a(x)\|$ is upper semicontinuous.*
- (ii) $\|\pi_Y(a)\| = \max\{\|\pi_x(a)\| : x \in Y\}$.
- (iii) *If $a(x) \in \pi_x(B)$ for all $x \in X$, then $a \in B$.*
- (iv) *If $\delta > 0$ and $a(x) \in_\delta \pi_x(B)$ for all $x \in X$, then $a \in_\delta B$.*
- (v) *The restriction of $\pi_x : A \rightarrow A(x)$ to B induces an isomorphism $B(x) \cong \pi_x(B)$ for all $x \in X$.*

Proof. (i), (ii) are proved in [3] and (iii) follows from (iv). (iv): By assumption, for each $x \in X$, there is $b_x \in B$ such that $\|\pi_x(a - b_x)\| < \delta$. Using (i) and (ii), we find a closed neighborhood U_x of x such that $\|\pi_{U_x}(a - b_x)\| < \delta$. Since X is compact, there is a finite subcover (U_{x_i}) . Let (α_i) be a partition of unity subordinated to this cover. Setting $b = \sum_i \alpha_i b_{x_i} \in B$, one checks immediately that $\|\pi_x(a - b)\| \leq \sum_i \alpha_i(x) \|\pi_x(a - b_{x_i})\| < \delta$, for all $x \in X$. Thus $\|a - b\| < \delta$ by (ii). (v): If $\iota : B \hookrightarrow A$ is the inclusion map, then $\pi_x(B)$ coincides with the image of $\iota_x : B/C(X, x)B \rightarrow A/C(X, x)A$. Thus it suffices to check that ι_x is injective. If $\iota_x(b + C(X, x)B) = \pi_x(b) = 0$ for some $b \in B$, then $b = fa$ for some $f \in C(X, x)$ and some $a \in A$. If (f_λ) is an approximate unit of $C(X, x)$, then $b = \lim_\lambda f_\lambda fa = \lim_\lambda f_\lambda b$ and hence $b \in C(X, x)B$. \square

A $C(X)$ -algebra such that the map $x \mapsto \|a(x)\|$ is continuous for all $a \in A$ is called a *continuous $C(X)$ -algebra* or a C^* -bundle [3,23,4]. A C^* -algebra A is a continuous $C(X)$ -algebra if and only if A is the C^* -algebra of continuous sections of a continuous field of C^* -algebras over X in the sense of [12, Def. 10.3.1] (see [3,4,27]).

Lemma 2.2. *Let A be a separable continuous $C(X)$ -algebra over a locally compact Hausdorff space X . If all the fibers of A are nonzero, then X has a countable basis of open sets. Thus the compact subspaces of X are metrizable.*

Proof. Since A is separable, its primitive spectrum $\text{Prim}(A)$ has a countable basis of open sets by [12, 3.3.4]. The continuous map $\eta : \text{Prim}(A) \rightarrow X$ (induced by $\theta : C_0(X) \rightarrow ZM(A) \cong C_b(\text{Prim}(A))$) is open since the $C(X)$ -algebra A is continuous and surjective since $A(x) \neq 0$ for all $x \in X$ (see [4, p. 388] and [27, Prop. 2.1, Thm. 2.3]). \square

Lemma 2.3. *Let X be a compact metrizable space. A $C(X)$ -algebra A all of whose fibers are nonzero and simple is continuous if and only if there is $e \in A$ such that $\|e(x)\| \geq 1$ for all $x \in X$.*

Proof. By Lemma 2.1(i) it suffices to prove that $\liminf_{n \rightarrow \infty} \|a(x_n)\| \geq \|a(x_0)\|$ for any $a \in A$ and any sequence (x_n) converging to x_0 in X . Set $D = A(x_0)$ and let e be as in the statement. Let $\psi : D \rightarrow A$ be a set-theoretical lifting of id_D such that $\|\psi(d)\| = \|d\|$ for all $d \in D$. Then $\lim_{n \rightarrow \infty} \|\pi_{x_n} \psi(a(x_0)) - a(x_n)\| = 0$ for all $a \in A$, by Lemma 2.1(i). By applying this

to e , since $\|e(x_n)\| \geq 1$, we see that $\liminf_{n \rightarrow \infty} \|\pi_{x_n} \psi(e(x_0))\| \geq 1$. Since D is a simple C^* -algebra, if $\varphi_n : D \rightarrow B_n$ is a sequence of contractive maps such that $\lim_{n \rightarrow \infty} \|\varphi_n(\lambda c + d) - \lambda \varphi_n(c) - \varphi_n(d)\| = 0$, $\lim_{n \rightarrow \infty} \|\varphi_n(cd) - \varphi_n(c)\varphi_n(d)\| = 0$, $\lim_{n \rightarrow \infty} \|\varphi_n(c^*) - \varphi_n(c)^*\| = 0$, for all $c, d \in D$, $\lambda \in \mathbb{C}$, and $\liminf_{n \rightarrow \infty} \|\varphi_n(c)\| > 0$ for some $c \in D$, then $\lim_{n \rightarrow \infty} \|\varphi_n(c)\| = \|c\|$ for all $c \in D$. In particular this observation applies to $\varphi_n = \pi_{x_n} \psi$ by Lemma 2.1(i). Therefore

$$\liminf_{n \rightarrow \infty} \|a(x_n)\| \geq \liminf_{n \rightarrow \infty} (\|\pi_{x_n} \psi(a(x_0))\| - \|\pi_{x_n} \psi(a(x_0)) - a(x_n)\|) = \|a(x_0)\|.$$

Conversely, if A is continuous, take e to be a large multiple of some full element of A . \square

Let $\eta : B \rightarrow A$ and $\psi : E \rightarrow A$ be $*$ -homomorphisms. The pullback of these maps is

$$B \oplus_{\eta, \psi} E = \{(b, e) \in B \oplus E : \eta(b) = \psi(e)\}.$$

We are going to use pullbacks in the context of $C(X)$ -algebras. Let X be a compact space and let Y, Z be closed subsets of X such that $X = Y \cup Z$. The following result is proved in [12, Prop. 10.1.13] for continuous $C(X)$ -algebras.

Lemma 2.4. *If A is a $C(X)$ -algebra, then A is isomorphic to $A(Y) \oplus_{\pi, \pi} A(Z)$, the pullback of the restriction maps $\pi_{Y \cap Z}^Y : A(Y) \rightarrow A(Y \cap Z)$ and $\pi_{Y \cap Z}^Z : A(Z) \rightarrow A(Y \cap Z)$.*

Proof. By the universal property of pullbacks, the maps π_Y and π_Z induce a map $\eta : A \rightarrow A(Y) \oplus_{\pi, \pi} A(Z)$, $\eta(a) = (\pi_Y(a), \pi_Z(a))$, which is injective by Lemma 2.1(ii). Thus it suffices to show that the range of η is dense. Let $b, c \in A$ be such that $\pi_{Y \cap Z}(b - c) = 0$ and let $\varepsilon > 0$. We shall find $a \in A$ such that $\|\eta(a) - (\pi_Y(b), \pi_Z(c))\| < \varepsilon$. By Lemma 2.1(i), there is an open neighborhood V of $Y \cap Z$ such that $\|\pi_x(b - c)\| < \varepsilon$ for all $x \in V$. Let $\{\lambda, \mu\}$ be a partition of unity on X subordinated to the open cover $\{Y \cup V, Z \cup V\}$. Then $a = \lambda b + \mu c$ is an element of A which has the desired property. \square

Let $B \subset A(Y)$ and $E \subset A(Z)$ be $C(X)$ -subalgebras such that $\pi_{Y \cap Z}^Z(E) \subseteq \pi_{Y \cap Z}^Y(B)$. As an immediate consequence of Lemma 2.4 we see that the pullback $B \oplus_{\pi_{Y \cap Z}^Z, \pi_{Y \cap Z}^Y} E$ is isomorphic to the $C(X)$ -subalgebra $B \oplus_{Y \cap Z} E$ of A defined as

$$B \oplus_{Y \cap Z} E = \{a \in A : \pi_Y(a) \in B, \pi_Z(a) \in E\}.$$

Lemma 2.5. *The fibers of $B \oplus_{Y \cap Z} E$ are given by*

$$\pi_x(B \oplus_{Y \cap Z} E) = \begin{cases} \pi_x(B), & \text{if } x \in X \setminus Z, \\ \pi_x(E), & \text{if } x \in Z, \end{cases}$$

and there is an exact sequence of C^* -algebras

$$0 \longrightarrow \{b \in B : \pi_{Y \cap Z}(b) = 0\} \longrightarrow B \oplus_{Y \cap Z} E \xrightarrow{\pi_Z} E \longrightarrow 0. \tag{1}$$

Proof. Let $x \in X \setminus Z$. The inclusion $\pi_x(B \oplus_{Y \cap Z} E) \subset \pi_x(B)$ is obvious by definition. Given $b \in B$, let us choose $f \in C(X)$ vanishing on Z and such that $f(x) = 1$. Then $a = (fb, 0)$ is an element of A by Lemma 2.4. Moreover $a \in B \oplus_{Y \cap Z} E$ and $\pi_x(a) = \pi_x(b)$. We have $\pi_Z(B \oplus_{Y \cap Z} E) \subset E$, by definition. Conversely, given $e \in E$, let us observe that $\pi_{Y \cap Z}^Z(e) \in \pi_{Y \cap Z}^Y(B)$ (by assumption) and hence $\pi_{Y \cap Z}^Z(e) = \pi_{Y \cap Z}^Y(b)$ for some $b \in B$. Then $a = (b, e)$ is an element of A by Lemma 2.4 and $\pi_Z(a) = e$. This completes the proof for the first part of the lemma and also it shows that the map π_Z from the sequence (1) is surjective. Its kernel is identified using Lemma 2.1(iii). \square

Let X, Y, Z and A be as above. Let $\eta : B \hookrightarrow A(Y)$ be a $C(Y)$ -linear $*$ -monomorphism and let $\psi : E \hookrightarrow A(Z)$ be a $C(Z)$ -linear $*$ -monomorphism. Assume that

$$\pi_{Y \cap Z}^Z(\psi(E)) \subseteq \pi_{Y \cap Z}^Y(\eta(B)). \tag{2}$$

This gives a map $\gamma = \eta_{Y \cap Z}^{-1} \psi_{Y \cap Z} : E(Y \cap Z) \rightarrow B(Y \cap Z)$. To simplify notation we let π stand for both $\pi_{Y \cap Z}^Y$ and $\pi_{Y \cap Z}^Z$ in the following lemma.

Lemma 2.6.

(a) *There are isomorphisms of $C(X)$ -algebras*

$$B \oplus_{\pi, \gamma \pi} E \cong B \oplus_{\pi \eta, \pi \psi} E \cong \eta(B) \oplus_{Y \cap Z} \psi(E),$$

where the second isomorphism is given by the map $\chi : B \oplus_{\pi \eta, \pi \psi} E \rightarrow A$ induced by the pair (η, ψ) . Its components χ_x can be identified with ψ_x for $x \in Z$ and with η_x for $x \in X \setminus Z$.

- (b) *Condition (2) is equivalent to $\psi(E) \subset \pi_Z(A \oplus_Y \eta(B))$.*
- (c) *If \mathcal{F} is a finite subset of A such that $\pi_Y(\mathcal{F}) \subset_\varepsilon \eta(B)$ and $\pi_Z(\mathcal{F}) \subset_\varepsilon \psi(E)$, then $\mathcal{F} \subset_\varepsilon \eta(B) \oplus_{Y \cap Z} \psi(E) = \chi(B \oplus_{\pi \eta, \pi \psi} E)$.*

Proof. This is an immediate corollary of Lemmas 2.1, 2.4, 2.5. For illustration, let us verify (c). By assumption $\pi_x(\mathcal{F}) \subset_\varepsilon \eta_x(B)$ for all $x \in X \setminus Z$ and $\pi_z(\mathcal{F}) \subset_\varepsilon \psi_z(E)$ for all $z \in Z$. We deduce from Lemma 2.5 that $\pi_x(\mathcal{F}) \subset_\varepsilon \pi_x(\eta(B) \oplus_{Y \cap Z} \psi(E))$ for all $x \in X$. Therefore $\mathcal{F} \subset_\varepsilon \eta(B) \oplus_{Y \cap Z} \psi(E)$ by Lemma 2.1(iv). \square

Definition 2.7. Let \mathcal{C} be a class of C^* -algebras. A $C(Z)$ -algebra E is called \mathcal{C} -elementary if there is a finite partition of Z into closed subsets Z_1, \dots, Z_r ($r \geq 1$) and there exist C^* -algebras D_1, \dots, D_r in \mathcal{C} such that $E \cong \bigoplus_{i=1}^r C(Z_i) \otimes D_i$. The notion of *category* of a $C(X)$ -algebra with respect to a class \mathcal{C} is defined inductively: if A is \mathcal{C} -elementary then $\text{cat}_{\mathcal{C}}(A) = 0$; $\text{cat}_{\mathcal{C}}(A) \leq n$ if there are closed subsets Y and Z of X with $X = Y \cup Z$ and there exist a $C(Y)$ -algebra B such that $\text{cat}_{\mathcal{C}}(B) \leq n - 1$, a \mathcal{C} -elementary $C(Z)$ -algebra E and a $*$ -monomorphism of $C(Y \cap Z)$ -algebras $\gamma : E(Y \cap Z) \rightarrow B(Y \cap Z)$ such that A is isomorphic to

$$B \oplus_{\pi, \gamma \pi} E = \{(b, d) \in B \oplus E : \pi_{Y \cap Z}^Y(b) = \gamma \pi_{Y \cap Z}^Z(d)\}.$$

By definition $\text{cat}_{\mathcal{C}}(A) = n$ if n is the smallest number with the property that $\text{cat}_{\mathcal{C}}(A) \leq n$. If no such n exists, then $\text{cat}_{\mathcal{C}}(A) = \infty$.

Definition 2.8. Let \mathcal{C} be a class of C^* -algebras and let A be a $C(X)$ -algebra. An n -fibred \mathcal{C} -monomorphism (ψ_0, \dots, ψ_n) into A consists of $(n + 1)$ $*$ -monomorphisms of $C(X)$ -algebras $\psi_i : E_i \rightarrow A(Y_i)$, where Y_0, \dots, Y_n is a closed cover of X , each E_i is a \mathcal{C} -elementary $C(Y_i)$ -algebra and

$$\pi_{Y_i \cap Y_j}^{Y_i} \psi_i(E_i) \subseteq \pi_{Y_i \cap Y_j}^{Y_j} \psi_j(E_j), \quad \text{for all } i \leq j. \tag{3}$$

Given an n -fibred morphism into A we have an associated *continuous* $C(X)$ -algebra defined as the fibred product (or pullback) of the $*$ -monomorphisms ψ_i :

$$A(\psi_0, \dots, \psi_n) = \{(d_0, \dots, d_n) : d_i \in E_i, \pi_{Y_i \cap Y_j}^{Y_i} \psi_i(d_i) = \pi_{Y_i \cap Y_j}^{Y_j} \psi_j(d_j) \text{ for all } i, j\} \tag{4}$$

and an induced $C(X)$ -monomorphism (defined by using Lemma 2.4)

$$\eta = \eta_{(\psi_0, \dots, \psi_n)} : A(\psi_0, \dots, \psi_n) \rightarrow A \subset \bigoplus_{i=0}^n A(Y_i),$$

$$\eta(d_0, \dots, d_n) = (\psi_0(d_0), \dots, \psi_n(d_n)).$$

There are natural coordinate maps $p_i : A(\psi_0, \dots, \psi_n) \rightarrow E_i$, $p_i(d_0, \dots, d_n) = d_i$. Let us set $X_k = Y_k \cup \dots \cup Y_n$. Then (ψ_k, \dots, ψ_n) is an $(n - k)$ -fibred \mathcal{C} -monomorphism into $A(X_k)$. Let $\eta_k : A(X_k)(\psi_k, \dots, \psi_n) \rightarrow A(X_k)$ be the induced map and set $B_k = A(X_k)(\psi_k, \dots, \psi_n)$. Let us note that $B_0 = A(\psi_0, \dots, \psi_n)$ and that there are natural $C(X_{k-1})$ -isomorphisms

$$B_{k-1} \cong B_k \oplus_{\pi_{\eta_k, \pi_{\psi_{k-1}}}} E_{k-1} \cong B_k \oplus_{\pi, \gamma_k \pi} E_{k-1}, \tag{5}$$

where π stands for $\pi_{X_k \cap Y_{k-1}}$ and $\gamma_k : E_{k-1}(X_k \cap Y_{k-1}) \rightarrow B_k(X_k \cap Y_{k-1})$ is defined by $(\gamma_k)_x = (\eta_k)_x^{-1}(\psi_{k-1})_x$, for all $x \in X_k \cap Y_{k-1}$. In particular, this decomposition shows that $\text{cat}_{\mathcal{C}}(A(\psi_0, \dots, \psi_n)) \leq n$.

Lemma 2.9. Suppose that the class \mathcal{C} from Definition 2.7 consists of stable Kirchberg algebras. If A is a $C(X)$ -algebra over a compact metrizable space X such that $\text{cat}_{\mathcal{C}}(A) < \infty$, then A contains a full properly infinite projection and $A \cong A \otimes \mathcal{O}_{\infty} \otimes \mathcal{K}$.

Proof. We prove this by induction on $n = \text{cat}_{\mathcal{C}}(A)$. The case $n = 0$ is immediate since $D \cong D \otimes \mathcal{O}_{\infty}$ for any Kirchberg algebra D [19]. Let $A = B \oplus_{\pi, \gamma \pi} E$ where B, E and γ are as in Definition 2.7 with $\text{cat}_{\mathcal{C}}(B) = n - 1$ and $\text{cat}_{\mathcal{C}}(E) = 0$. Let us consider the exact sequence $0 \rightarrow J \rightarrow A \rightarrow E \rightarrow 0$, where $J = \{b \in B : \pi_{Y \cap Z}(b) = 0\}$. Since J is an ideal of $B \cong B \otimes \mathcal{O}_{\infty} \otimes \mathcal{K}$, J absorbs $\mathcal{O}_{\infty} \otimes \mathcal{K}$ by [22, Prop. 8.5]. Since both E and J are stable and purely infinite, it follows that A is stable by [30, Prop. 6.12] and purely infinite by [22, Prop. 3.5]. Since A has Hausdorff primitive spectrum, A is strongly purely infinite by [5, Thm. 5.8]. It follows that $A \cong A \otimes \mathcal{O}_{\infty}$ by [22, Thm. 9.1]. Finally A contains a full properly infinite projection since there is a full embedding of \mathcal{O}_2 into A by [5, Prop. 5.6]. \square

3. Semiprojectivity

In this section we study the notion of KK -semiprojectivity. The main result is Theorem 3.12. Let A and B be C^* -algebras. Two $*$ -homomorphisms $\varphi, \psi : A \rightarrow B$ are approximately unitarily equivalent, written $\varphi \approx_u \psi$, if there is a sequence of unitaries (u_n) in the C^* -algebra $B^+ = B + \mathbb{C}1$ obtained by adjoining a unit to B , such that $\lim_{n \rightarrow \infty} \|u_n \varphi(a) u_n^* - \psi(a)\| = 0$ for all $a \in A$. We say that φ and ψ are asymptotically unitarily equivalent, written $\varphi \approx_{uh} \psi$, if there is a norm continuous unitary valued map $t \rightarrow u_t \in B^+, t \in [0, 1)$, such that $\lim_{t \rightarrow 1} \|u_t \varphi(a) u_t^* - \psi(a)\| = 0$ for all $a \in A$. A $*$ -homomorphism $\varphi : D \rightarrow A$ is full if $\varphi(d)$ is not contained in any proper two-sided closed ideal of A if $d \in D$ is nonzero.

We shall use several times Kirchberg’s Theorem [29, Thm. 8.3.3] and the following theorem of Phillips [28].

Theorem 3.1. *Let A and B be separable C^* -algebras such that A is simple and nuclear, $B \cong B \otimes \mathcal{O}_\infty$, and there exist full projections $p \in A$ and $q \in B$. For any $\sigma \in KK(A, B)$ there is a full $*$ -homomorphism $\varphi : A \rightarrow B$ such that $KK(\varphi) = \sigma$. If $K_0(\sigma)[p] = [q]$ then we may arrange that $\varphi(p) = q$. If $\psi : A \rightarrow B$ is another $*$ -homomorphism such that $KK(\psi) = KK(\varphi)$ and $\psi(p) = q$, then $\varphi \approx_{uh} \psi$ via a path of unitaries $t \mapsto u_t \in U(qBq)$.*

Theorem 3.1 does not appear in this form in [28] but it is an immediate consequence of [28, Thm. 4.1.1]. Since $pAp \otimes \mathcal{K} \cong A \otimes \mathcal{K}$ and $qBq \otimes \mathcal{K} \cong B \otimes \mathcal{K}$ by [6], and $qBq \otimes \mathcal{O}_\infty \cong qBq$ by [22, Prop. 8.5], it suffices to discuss the case when p and q are the units of A and B . If σ is given, [28, Thm. 4.1.1] yields a full $*$ -homomorphism $\varphi : A \rightarrow B \otimes \mathcal{K}$ such that $KK(\varphi) = \sigma$. Let $e \in \mathcal{K}$ be a rank-one projection and suppose that $[\varphi(1_A)] = [1_B \otimes e]$ in $K_0(B)$. Since both $\varphi(1_A)$ and $1_B \otimes e$ are full projections and $B \cong B \otimes \mathcal{O}_\infty$, it follows by [28, Lemma 2.1.8] that $u\varphi(1_A)u^* = 1_B \otimes e$ for some unitary in $(B \otimes \mathcal{K})^+$. Replacing φ by $u\varphi u^*$ we can arrange that $KK(\varphi) = \sigma$ and φ is unital. For the second part of the theorem let us note that any unital $*$ -homomorphism $\varphi : A \rightarrow B$ is full and if two unital $*$ -homomorphisms $\varphi, \psi : A \rightarrow B$ are asymptotically unitarily equivalent when regarded as maps into $B \otimes \mathcal{K}$, then $\varphi \approx_{uh} \psi$ when regarded as maps into B , by an argument from the proof of [28, Thm. 4.1.4].

A separable nonzero C^* -algebra D is *semiprojective* [1] if for any separable C^* -algebra A and any increasing sequence of two-sided closed ideals (J_n) of A with $J = \overline{\bigcup_n J_n}$, the natural map $\varinjlim \text{Hom}(D, A/J_n) \rightarrow \text{Hom}(D, A/J)$ (induced by $\pi_n : A/J_n \rightarrow A/J$) is surjective. If we weaken this condition and require only that the above map has dense range, where $\text{Hom}(D, A/J)$ is given the point-norm topology, then D is called *weakly semiprojective* [14]. These definitions do not change if we drop the separability of A . We shall use (weak) semiprojectivity in the following context. Let A be a $C(X)$ -algebra (with X metrizable), let $x \in X$ and set $U_n = \{y \in X : d(y, x) \leq 1/n\}$. Then $J_n = C(X, U_n)A$ is an increasing sequence of ideals of A such that $J = C(X, x)A, A/J_n \cong A(U_n)$ and $A/J \cong A(x)$.

Examples 3.2 (*Weakly semiprojective C^* -algebras*). Any finite dimensional C^* -algebra is semiprojective. A Kirchberg algebra D satisfying the UCT and having finitely generated K -theory groups is weakly semiprojective by work of Neubüser [26], H. Lin [24] and Spielberg [32]. This also follows from Theorem 3.12 and Proposition 3.14 below. If in addition $K_1(D)$ is torsion free, then D is semiprojective as proved by Spielberg [33] who extended the foundational work of Blackadar [1] and Szymański [34].

The following generalizations of two results of Loring [25] are used in Section 5; see [10].

Proposition 3.3. *Let D be a separable semiprojective C^* -algebra. For any finite set $\mathcal{F} \subset D$ and any $\varepsilon > 0$, there exist a finite set $\mathcal{G} \subset D$ and $\delta > 0$ with the following property. Let $\pi : A \rightarrow B$ be a surjective $*$ -homomorphism, and let $\varphi : D \rightarrow B$ and $\gamma : D \rightarrow A$ be $*$ -homomorphisms such that $\|\pi\gamma(d) - \varphi(d)\| < \delta$ for all $d \in \mathcal{G}$. Then there is a $*$ -homomorphism $\psi : D \rightarrow A$ such that $\pi\psi = \varphi$ and $\|\gamma(c) - \psi(c)\| < \varepsilon$ for all $c \in \mathcal{F}$.*

Proposition 3.4. *Let D be a separable semiprojective C^* -algebra. For any finite set $\mathcal{F} \subset D$ and any $\varepsilon > 0$, there exist a finite set $\mathcal{G} \subset D$ and $\delta > 0$ with the following property. For any two $*$ -homomorphisms $\varphi, \psi : D \rightarrow B$ such that $\|\varphi(d) - \psi(d)\| < \delta$ for all $d \in \mathcal{G}$, there is a homotopy $\Phi \in \text{Hom}(D, C[0, 1] \otimes B)$ such that $\Phi_0 = \varphi$ to $\Phi_1 = \psi$ and $\|\varphi(c) - \Phi_t(c)\| < \varepsilon$ for all $c \in \mathcal{F}$ and $t \in [0, 1]$.*

Definition 3.5. A separable C^* -algebra D is *KK-stable* if there is a finite set $\mathcal{G} \subset D$ and there is $\delta > 0$ with the property that for any two $*$ -homomorphisms $\varphi, \psi : D \rightarrow A$ such that $\|\varphi(a) - \psi(a)\| < \delta$ for all $a \in \mathcal{G}$, one has $KK(\varphi) = KK(\psi)$.

Corollary 3.6. *Any semiprojective C^* -algebra is weakly semiprojective and KK-stable.*

Proof. This follows from Proposition 3.4. \square

Proposition 3.7. *Let D be a separable weakly semiprojective C^* -algebra. For any finite set $\mathcal{F} \subset D$ and any $\varepsilon > 0$ there exist a finite set $\mathcal{G} \subset D$ and $\delta > 0$ such that for any C^* -algebras $B \subset A$ and any $*$ -homomorphism $\varphi : D \rightarrow A$ with $\varphi(\mathcal{G}) \subset_\delta B$, there is a $*$ -homomorphism $\psi : D \rightarrow B$ such that $\|\varphi(c) - \psi(c)\| < \varepsilon$ for all $c \in \mathcal{F}$. If in addition D is KK-stable, then we can choose \mathcal{G} and δ such that we also have $KK(\psi) = KK(\varphi)$.*

Proof. This follows from [14, Thms. 3.1, 4.6]. Since the result is essential to us we include a short proof. Fix \mathcal{F} and ε . Let (\mathcal{G}_n) be an increasing sequence of finite subsets of D whose union is dense in D . If the statement is not true, then there are sequences of C^* -algebras $C_n \subset A_n$ and $*$ -homomorphisms $\varphi_n : D \rightarrow A_n$ satisfying $\varphi_n(\mathcal{G}_n) \subset_{1/n} C_n$ and with the property that for any $n \geq 1$ there is no $*$ -homomorphism $\psi_n : D \rightarrow C_n$ such that $\|\varphi_n(c) - \psi_n(c)\| < \varepsilon$ for all $c \in \mathcal{F}$. Set $B_i = \prod_{n \geq i} A_n$ and $E_i = \prod_{n \geq i} C_n \subset B_i$. If $v_i : B_i \rightarrow B_{i+1}$ is the natural projection, then $v_i(E_i) = E_{i+1}$. Let us observe that if we define $\Phi_i : D \rightarrow B_i$ by $\Phi_i(d) = (\varphi_i(d), \varphi_{i+1}(d), \dots)$, then the image of $\Phi = \varinjlim \Phi_i : D \rightarrow \varinjlim (B_i, v_i)$ is contained in $\varinjlim (E_i, v_i)$. Since D is weakly semiprojective, there is i and a $*$ -homomorphism $\Psi_i : D \rightarrow E_i$, of the form $\Psi_i(d) = (\psi_i(d), \psi_{i+1}(d), \dots)$ such that $\|\Phi_i(c) - \Psi_i(c)\| < \varepsilon$ for all $c \in \mathcal{F}$. Therefore $\|\varphi_i(c) - \psi_i(c)\| < \varepsilon$ for all $c \in \mathcal{F}$ which gives a contradiction. \square

It is useful to combine Propositions 3.7 and 3.3 in a single statement.

Proposition 3.8. *Let D be a separable semiprojective C^* -algebra. For any finite set $\mathcal{F} \subset D$ and any $\varepsilon > 0$, there exist a finite set $\mathcal{G} \subset D$ and $\delta > 0$ with the following property. Let $\pi : A \rightarrow B$ be a surjective $*$ -homomorphism which maps a C^* -subalgebra A' of A onto a C^* -subalgebra B' of B . Let $\varphi : D \rightarrow B'$ and $\gamma : D \rightarrow A$ be $*$ -homomorphisms such that $\gamma(\mathcal{G}) \subset_\delta A'$*

and $\|\pi\gamma(d) - \varphi(d)\| < \delta$ for all $d \in \mathcal{G}$. Then there is a $*$ -homomorphism $\psi : D \rightarrow A'$ such that $\pi\psi = \varphi$ and $\|\gamma(c) - \psi(c)\| < \varepsilon$ for all $c \in \mathcal{F}$.

Proof. Let \mathcal{G}_L and δ_L be given by Proposition 3.3 applied to the input data \mathcal{F} and $\varepsilon/2$. We may assume that $\mathcal{F} \subset \mathcal{G}_L$ and $\varepsilon > \delta_L$. Next, let \mathcal{G}_P and δ_P be given by Proposition 3.7 applied to the input data \mathcal{G}_L and $\delta_L/2$. We show now that $\mathcal{G} := \mathcal{G}_L \cup \mathcal{G}_P$ and $\delta := \min\{\delta_P, \delta_L/2\}$ have the desired properties. We have $\gamma(\mathcal{G}_P) \subset_{\delta_P} A'$ since $\mathcal{G}_P \subset \mathcal{G}$ and $\delta \leq \delta_P$. By Proposition 3.7 there is a $*$ -homomorphism $\gamma' : D \rightarrow A'$ such that $\|\gamma'(d) - \gamma(d)\| < \delta_L/2$ for all $d \in \mathcal{G}_L$. Then, since $\mathcal{G}_L \subset \mathcal{G}$ and $\delta \leq \delta_L/2$,

$$\|\pi\gamma'(d) - \varphi(d)\| \leq \|\pi\gamma'(d) - \pi\gamma(d)\| + \|\pi\gamma(d) - \varphi(d)\| < \delta_L/2 + \delta \leq \delta_L$$

for all $d \in \mathcal{G}_L$. Therefore we can invoke Proposition 3.3 to perturb γ' to a $*$ -homomorphism $\psi : D \rightarrow A'$ such that $\pi\psi = \varphi$ and $\|\gamma'(d) - \psi(d)\| < \varepsilon/2$ for all $d \in \mathcal{F}$. Finally we observe that for $d \in \mathcal{F} \subset \mathcal{G}_L$

$$\|\gamma(d) - \psi(d)\| \leq \|\gamma(d) - \gamma'(d)\| + \|\gamma'(d) - \psi(d)\| < \delta_L/2 + \varepsilon/2 < \varepsilon. \quad \square$$

Definition 3.9. (a) A separable C^* -algebra D is *KK-semiprojective* if for any separable C^* -algebra A and any increasing sequence of two-sided closed ideals (J_n) of A with $J = \bigcup_n J_n$, the natural map $\varinjlim KK(D, A/J_n) \rightarrow KK(D, A/J)$ is surjective.

(b) We say that the functor $KK(D, -)$ is *continuous* if for any inductive system $B_1 \rightarrow B_2 \rightarrow \dots$ of separable C^* -algebras, the induced map $\varinjlim KK(D, B_n) \rightarrow KK(D, \varinjlim B_n)$ is bijective.

Proposition 3.10. Any separable *KK-semiprojective* C^* -algebra is *KK-stable*.

Proof. We shall prove the statement by contradiction. Let D be separable *KK-semiprojective* C^* -algebra. Let (\mathcal{G}_n) be an increasing sequence of finite subsets of D whose union is dense in D . If the statement is not true, then there are sequences of $*$ -homomorphisms $\varphi_n, \psi_n : D \rightarrow A_n$ such that $\|\varphi_n(d) - \psi_n(d)\| < 1/n$ for all $d \in \mathcal{G}_n$ and yet $KK(\varphi_n) \neq KK(\psi_n)$ for all $n \geq 1$. Set $B_i = \prod_{n \geq i} A_n$ and let $v_i : B_i \rightarrow B_{i+1}$ be the natural projection. Let us define $\Phi_i, \Psi_i : D \rightarrow B_i$ by $\Phi_i(d) = (\varphi_i(d), \varphi_{i+1}(d), \dots)$ and $\Psi_i(d) = (\psi_i(d), \psi_{i+1}(d), \dots)$, for all d in D . Let B'_i be the separable C^* -subalgebra of B_i generated by the images of Φ_i and Ψ_i . Then $v_i(B'_i) = B'_{i+1}$ and one verifies immediately that $\varinjlim \Phi_i = \varinjlim \Psi_i : D \rightarrow \varinjlim (B'_i, v_i)$. Since D is *KK-semiprojective*, we must have $KK(\Phi_i) = KK(\Psi_i)$ for some i and hence $KK(\varphi_n) = KK(\psi_n)$ for all $n \geq i$. This gives a contradiction. \square

Proposition 3.11. A unital Kirchberg algebra D is *KK-stable* if and only if $D \otimes \mathcal{K}$ is *KK-stable*. D is *weakly semiprojective* if and only if $D \otimes \mathcal{K}$ is *weakly semiprojective*.

Proof. Since $KK(D, A) \cong KK(D, A \otimes \mathcal{K}) \cong KK(D \otimes \mathcal{K}, A \otimes \mathcal{K})$ the first part of the proposition is immediate. Suppose now that $D \otimes \mathcal{K}$ is weakly semiprojective. Then D is weakly semiprojective as shown in the proof of [32, Thm. 2.2]. Conversely, assume that D is weakly semiprojective. It suffices to find $\alpha \in \text{Hom}(D \otimes \mathcal{K}, D)$ and a sequence (β_n) in $\text{Hom}(D, D \otimes \mathcal{K})$ such that $\beta_n \alpha$ converges to $\text{id}_{D \otimes \mathcal{K}}$ in the point-norm topology. Let s_i be the canonical generators of \mathcal{O}_∞ . If (e_{ij}) is a system of matrix units for \mathcal{K} , then $\lambda(e_{ij}) = s_i s_j^*$ defines a $*$ -homomorphism $\mathcal{K} \rightarrow \mathcal{O}_\infty$

such that $KK(\lambda) \in KK(\mathcal{K}, \mathcal{O}_\infty)^{-1}$. Therefore, by composing $\text{id}_D \otimes \lambda$ with some isomorphism $D \otimes \mathcal{O}_\infty \cong D$ (given by [29, Thm. 7.6.6]) we obtain a $*$ -monomorphism $\alpha : D \otimes \mathcal{K} \rightarrow D$ which induces a KK -equivalence. Let $\beta : D \rightarrow D \otimes \mathcal{K}$ be defined by $\beta(d) = d \otimes e_{11}$. Then $\beta\alpha \in \text{End}(D \otimes \mathcal{K})$ induces a KK -equivalence and hence after replacing β by $\theta\beta$ for some automorphism θ of $D \otimes \mathcal{K}$, we may arrange that $KK(\beta\alpha) = KK(\text{id}_D)$. By Theorem 3.1, $\beta\alpha \approx_u \text{id}_{D \otimes \mathcal{K}}$, so that there is a sequence of unitaries $u_n \in (D \otimes \mathcal{K})^+$ such that $u_n\beta\alpha(-)u_n^*$ converges to $\text{id}_{D \otimes \mathcal{K}}$. \square

Theorem 3.12. *For a separable C^* -algebra D consider the following properties:*

- (i) D is KK -semiprojective.
- (ii) The functor $KK(D, -)$ is continuous.
- (iii) D is weakly semiprojective and KK -stable.

Then (i) \Leftrightarrow (ii). Moreover, (iii) \Rightarrow (i) if D is nuclear and (i) \Rightarrow (iii) if D is a Kirchberg algebra. Thus (i) \Leftrightarrow (ii) \Leftrightarrow (iii) for any Kirchberg algebra D .

Proof. The implication (ii) \Rightarrow (i) is obvious. (i) \Rightarrow (ii): Let (B_n, γ_n, m) be an inductive system with inductive limit B and let $\gamma_n : B_n \rightarrow B$ be the canonical maps. We have an induced map $\beta : \varinjlim KK(D, B_n) \rightarrow KK(D, B)$. First we show that β is surjective. The mapping telescope construction of L.G. Brown (as described in the proof of [1, Thm. 3.1]) produces an inductive system of C^* -algebras (T_n, η_n, m) with inductive limit B such that each $\eta_{n,n+1}$ is surjective, and each canonical map $\eta_n : T_n \rightarrow B$ is homotopic to $\gamma_n\alpha_n$ for some $*$ -homomorphism $\alpha_n : T_n \rightarrow B_n$. In particular $KK(\eta_n) = KK(\gamma_n)KK(\alpha_n)$. Let $x \in KK(D, B)$. By (i) there are n and $y \in KK(D, T_n)$ such that $KK(\eta_n)y = x$ and hence $KK(\gamma_n)KK(\alpha_n)y = x$. Thus $z = KK(\alpha_n)y \in KK(D, B_n)$ is a lifting of x . Let us show now that the map β is injective. Let x be an element in the kernel of the map $KK(D, B_n) \rightarrow KK(D, B)$. Consider the commutative diagram whose exact rows are portions of the Puppe sequence in K -theory [2, Thm. 19.4.3] and with vertical maps induced by $\gamma_m : B_m \rightarrow B, m \geq n$.

$$\begin{array}{ccccc}
 KK(D, C_{\gamma_n}) & \longrightarrow & KK(D, B_n) & \longrightarrow & KK(D, B) \\
 \uparrow & & \parallel & & \uparrow \\
 KK(D, C_{\gamma_{n,m}}) & \longrightarrow & KK(D, B_n) & \longrightarrow & KK(D, B_m)
 \end{array}$$

By exactness, x is the image of some element $y \in KK(D, C_{\gamma_n})$. Since $C_{\gamma_n} = \varinjlim C_{\gamma_{n,m}}$, the map $\varinjlim KK(D, C_{\gamma_{n,m}}) \rightarrow KK(D, C_{\gamma_n})$ is surjective by the first part of the proof. Therefore there is $m \geq n$ such that y lifts to some $z \in KK(D, C_{\gamma_{n,m}})$. The image of z in $KK(D, B_m)$ equals $KK(\gamma_{n,m})x$ and vanishes by exactness of the bottom row.

(iii) \Rightarrow (i): Let $A, (J_n)$ and J be as in Definition 3.9. Using the five-lemma and the split exact sequence $0 \rightarrow KK(D, A) \rightarrow KK(D, A^+) \rightarrow KK(D, \mathbb{C}) \rightarrow 0$, we reduce the proof to the case when A is unital. Let $x \in KK(D, A/J)$. Since the map $KK(D^+, A/J) \rightarrow KK(D, A/J)$ is surjective, x lifts to some element $x^+ \in KK(D^+, A/J)$. By [29, Thm. 8.3.3], since D^+ is nuclear, there is a $*$ -homomorphism $\Phi : D^+ \rightarrow A/J \otimes \mathcal{O}_\infty \otimes \mathcal{K}$ such that $KK(\Phi) = x^+$ and hence if set $\varphi = \Phi|_D$, then $KK(\varphi) = x$. Since D is weakly semiprojective, there are n and a $*$ -homomorphism $\psi : D \rightarrow A/J_n \otimes \mathcal{O}_\infty \otimes \mathcal{K}$ such that $\|\pi_n\psi(d) - \varphi(d)\| < \delta$ for all $d \in \mathcal{G}$, where

\mathcal{G} and δ are as in the definition of KK -stability. Therefore $KK(\pi_n\psi) = KK(\varphi)$ and hence $KK(\psi)$ is a lifting of x to $KK(D, A/J_n)$.

(i) \Rightarrow (iii): D is KK -stable by Proposition 3.10. It remains to show that D is weakly semiprojective. Since any nonunital Kirchberg algebra is isomorphic to the stabilization of a unital one (see [29, Prop. 4.1.3]) and since by Proposition 3.11 D is KK -semiprojective if and only if $D \otimes \mathcal{K}$ is KK -semiprojective, we may assume that D is unital. Let $A, (J_n), \pi_{m,n} : A/J_m \rightarrow A/J_n$ ($m \leq n$) and $\pi_n : A/J_n \rightarrow A/J$ be as in the definition of weak semiprojectivity. By [1, Cor. 2.15], we may assume that A and the $*$ -homomorphism $\varphi : D \rightarrow A$ (for which we want to construct an approximative lifting) are unital. In particular φ is injective since D is simple. Set $B = \varphi(D) \subset A/J$ and $B_n = \pi_n^{-1}(B) \subset A/J_n$. The corresponding maps $\pi_{m,n} : B_m \rightarrow B_n$ ($m \leq n$) and $\pi_n : B_n \rightarrow B$ are surjective and they induce an isomorphism $\varinjlim(B_n, \pi_{n,n+1}) \cong B$.

Given $\varepsilon > 0$ and $\mathcal{F} \subset D$ (a finite set) we are going to produce an approximate lifting $\varphi_n : D \rightarrow B_n$ for φ . Since 1_B is a properly infinite projection, it follows by [1, Props. 2.18 and 2.23] that the unit 1_n of B_n is a properly infinite projection, for all sufficiently large n . Since D is KK -semiprojective, there exist m and an element $h \in KK(D, B_m)$ which lifts $KK(\varphi)$ such that $K_0(h)[1_D] = [1_m]$. By [29, Thm. 8.3.3], there is a full $*$ -homomorphism $\eta : D \rightarrow B_m \otimes \mathcal{K}$ such that $KK(\eta) = h$. By [29, Prop. 4.1.4], since both $\eta(1_D)$ and 1_m are full and properly infinite projections in $B_m \otimes \mathcal{K}$, there is a partial isometry $w \in B_m \otimes \mathcal{K}$ such that $w^*w = \eta(1_D)$ and $ww^* = 1_m$. Replacing η by $w\eta(-)w^*$, we may assume that $\eta : D \rightarrow B_m$ is unital. Then $KK(\pi_m\eta) = KK(\pi_m)h = KK(\varphi)$. By Theorem 3.1, $\pi_m\eta \approx_{uh} \varphi$. Thus there is a unitary $u \in B$ such that $\|u\pi_m\eta(d)u^* - \varphi(d)\| < \varepsilon$ for all $d \in \mathcal{F}$. Since $C(\mathbb{T})$ is semiprojective, there is $n \geq m$ such that u lifts to a unitary $u_n \in B_n$. Then $\varphi_n := u_n\pi_{m,n}\eta(-)u_n^*$ is a $*$ -homomorphism from D to B_n such that $\|\pi_n\varphi_n(d) - \varphi(d)\| < \varepsilon$ for all $d \in \mathcal{F}$. \square

Corollary 3.13. *Any separable nuclear semiprojective C^* -algebra is KK -semiprojective.*

Proof. This is very similar to the proof of the implication (iii) \Rightarrow (i) of Theorem 3.12. Alternatively, the statement follows from Corollary 3.6 and Theorem 3.12. \square

Blackadar has shown that a semiprojective Kirchberg algebra satisfying the UCT has finitely generated K -theory groups [29, Prop. 8.4.15]. A similar argument gives the following:

Proposition 3.14. *Let D be a separable C^* -algebra satisfying the UCT. Then D is KK -semiprojective if and only if $K_*(D)$ is finitely generated.*

Proof. If $K_*(D)$ is finitely generated, then D is KK -semiprojective by [31]. Conversely, assume that D is KK -semiprojective. Since D satisfies the UCT, we infer that if $G = K_i(D)$ ($i = 0, 1$), then G is semiprojective in the category of countable abelian groups, in the sense that if $H_1 \rightarrow H_2 \rightarrow \dots$ is an inductive system of countable abelian groups with inductive limit H , then the natural map $\varinjlim \text{Hom}(G, H_n) \rightarrow \text{Hom}(G, H)$ is surjective. This implies that G is finitely generated. Indeed, taking $H = G$, we see that id_G lifts to $\text{Hom}(G, H_n)$ for some finitely generated subgroup H_n of G and hence G is a quotient of H_n . \square

4. Approximation of $C(X)$ -algebras

In this section we use weak semiprojectivity to approximate a continuous $C(X)$ -algebra A by $C(X)$ -subalgebras given by pullbacks of n -fibered monomorphisms into A .

Lemma 4.1. *Let D be a finite direct sum of simple C^* -algebras and let $\varphi, \psi : D \rightarrow A$ be $*$ -homomorphisms. Suppose that $\mathcal{H} \subset D$ contains a nonzero element from each simple direct summand of D . If $\|\psi(d) - \varphi(d)\| \leq \|d\|/2$ for all $d \in \mathcal{H}$, then φ is injective if and only if ψ is injective.*

Proof. Let us note that φ is injective if and only if $\|\varphi(d)\| = \|d\|$ for all $d \in \mathcal{H}$. Therefore if φ is injective, then $\|\psi(d)\| \geq \|\varphi(d)\| - \|\psi(d) - \varphi(d)\| \geq \|d\|/2$ for all $d \in \mathcal{H}$ and hence ψ is injective. \square

A sequence (A_n) of subalgebras of A is called *exhaustive* if for any finite subset \mathcal{F} of A and any $\varepsilon > 0$ there is n such that $\mathcal{F} \subset_\varepsilon A_n$.

Lemma 4.2. *Let \mathcal{C} be a class consisting of finite direct sums of separable simple weakly semiprojective C^* -algebras. Let X be a compact metrizable space and let A be a $C(X)$ -algebra. Let $\mathcal{F} \subset A$ be a finite set, let $\varepsilon > 0$ and suppose that $A(x)$ admits an exhaustive sequence of C^* -algebras isomorphic to C^* -algebras in \mathcal{C} for some $x \in X$. Then there exist a compact neighborhood U of x and a $*$ -homomorphism $\varphi : D \rightarrow A(U)$ for some $D \in \mathcal{C}$ such that $\pi_U(\mathcal{F}) \subset_\varepsilon \varphi(D)$. If A is a continuous $C(X)$ -algebra, then we may arrange that φ_z is injective for all $z \in U$.*

Proof. Let $\mathcal{F} = \{a_1, \dots, a_r\}$ and ε be given. By hypothesis there exist $D \in \mathcal{C}$, $\{c_1, \dots, c_r\} \subset D$ and a $*$ -monomorphism $\iota : D \rightarrow A(x)$ such that $\|\pi_x(a_i) - \iota(c_i)\| < \varepsilon/2$, for all $i = 1, \dots, r$. Set $U_n = \{y \in X : d(x, y) \leq 1/n\}$. Choose a full element d_j in each direct summand of D . Since D is weakly semiprojective, there is a $*$ -homomorphism $\varphi : D \rightarrow A(U_n)$ (for some n) such that $\|\pi_x \varphi(c_i) - \iota(c_i)\| < \varepsilon/2$ for all $i = 1, \dots, r$, and $\|\pi_x \varphi(d_j) - \iota(d_j)\| \leq \|d_j\|/2$ for all d_j . Therefore

$$\|\pi_x \varphi(c_i) - \pi_x(a_i)\| \leq \|\pi_x \varphi(c_i) - \iota(c_i)\| + \|\pi_x(a_i) - \iota(c_i)\| < \varepsilon/2 + \varepsilon/2 = \varepsilon$$

and φ_x is injective by Lemma 4.1. By Lemma 2.1(i), after increasing n and setting $U = U_n$ and $\varphi = \pi_U \varphi$, we have

$$\|\varphi(c_i) - \pi_U(a_i)\| = \|\pi_U(\varphi(c_i) - a_i)\| < \varepsilon,$$

for all $i = 1, \dots, r$. This shows that $\pi_U(\mathcal{F}) \subset_\varepsilon \varphi(D)$. If A is continuous, then after shrinking U we may arrange that $\|\varphi_z(d_j)\| \geq \|\varphi_x(d_j)\|/2 = \|d_j\|/2$ for all d_j and all $z \in U$. This implies that φ_z is injective for all $z \in U$. \square

Lemma 4.3. *Let X be a compact metrizable space and let A be a separable continuous $C(X)$ -algebra the fibers of which are stable Kirchberg algebras. Let $\mathcal{F} \subset A$ be a finite set and let $\varepsilon > 0$. Suppose that there exist a KK -semiprojective stable Kirchberg algebra D and $\sigma \in KK(D, A)$ such that $\sigma_x \in KK(D, A(x))^{-1}$ for some $x \in X$. Then there exist a closed neighborhood U of x and a full $*$ -homomorphism $\psi : D \rightarrow A(U)$ such that $KK(\psi) = \sigma_U$ and $\pi_U(\mathcal{F}) \subset_\varepsilon \psi(D)$.*

Proof. By [29, Thm. 8.4.1] there is an isomorphism $\psi_0 : D \rightarrow A(x)$ such that $KK(\psi_0) = \sigma_x$. Let $\mathcal{H} \subset D$ be such that $\psi_0(\mathcal{H}) = \pi_x(\mathcal{F})$. Set $U_n = \{y \in X : d(x, y) \leq 1/n\}$. By Theorem 3.12 D is KK -stable and weakly semiprojective. By Proposition 3.7 there exists a $*$ -homomorphism $\psi_n : D \rightarrow A(U_n)$ (for some n) such that $\|\pi_x \psi_n(d) - \psi_0(d)\| < \varepsilon$ for all $d \in \mathcal{H}$ and $KK(\pi_x \psi_n) =$

$KK(\psi_0) = \sigma_x$. Since $\varinjlim_m KK(D, A(U_m)) = KK(D, A(x))$, we deduce that there is $m \geq n$ such that $KK(\pi_{U_m} \psi_n) = \sigma_{U_m}$. By increasing m we may arrange that $\pi_{U_m}(\mathcal{F}) \subset_\varepsilon \pi_{U_m} \psi_n(D)$ since we have seen that $\pi_x(\mathcal{F}) = \psi_0(\mathcal{H}) \subset_\varepsilon \pi_x \psi_n(D)$. We can arrange that ψ_z is injective for all $z \in U$ by reasoning as in the proof of Lemma 4.2. We conclude by setting $U = U_m$ and $\psi = \pi_{U_m} \psi_n$. \square

The following lemma is useful for constructing fibered morphisms.

Lemma 4.4. *Let $(D_j)_{j \in J}$ be a finite family consisting of finite direct sums of weakly semiprojective simple C^* -algebras. Let $\varepsilon > 0$ and for each $j \in J$ let $\mathcal{H}_j \subset D_j$ be a finite set such that for each direct summand of D_j there is an element of \mathcal{H}_j of norm $\geq \varepsilon$ which is contained and is full in that summand. Let $\mathcal{G}_j \subset D_j$ and $\delta_j > 0$ be given by Proposition 3.7 applied to D_j , \mathcal{H}_j and $\varepsilon/2$. Let X be a compact metrizable space, let $(Z_j)_{j \in J}$ be disjoint nonempty closed subsets of X and let Y be a closed nonempty subset of X such that $X = Y \cup (\bigcup_j Z_j)$. Let A be a continuous $C(X)$ -algebra and let \mathcal{F} be a finite subset of A . Let $\eta : B(Y) \rightarrow A(Y)$ be a $*$ -monomorphism of $C(Y)$ -algebras and let $\varphi_j : D_j \rightarrow A(Z_j)$ be $*$ -homomorphisms such that $(\varphi_j)_x$ is injective for all $x \in Z_j$ and $j \in J$, and which satisfy the following conditions:*

- (i) $\pi_{Z_j}(\mathcal{F}) \subset_{\varepsilon/2} \varphi_j(\mathcal{H}_j)$, for all $j \in J$,
- (ii) $\pi_Y(\mathcal{F}) \subset_\varepsilon \eta(B)$,
- (iii) $\pi_{Y \cap Z_j}^Z \varphi_j(\mathcal{G}_j) \subset_{\delta_j} \pi_{Y \cap Z_j}^Y \eta(B)$, for all $j \in J$.

Then, there are $C(Z_j)$ -linear $*$ -monomorphisms $\psi_j : C(Z_j) \otimes D_j \rightarrow A(Z_j)$, satisfying

$$\|\varphi_j(c) - \psi_j(c)\| < \varepsilon/2, \quad \text{for all } c \in \mathcal{H}_j, \text{ and } j \in J, \tag{6}$$

and such that if we set $E = \bigoplus_j C(Z_j) \otimes D_j$, $Z = \bigcup_j Z_j$, and $\psi : E \rightarrow A(Z) = \bigoplus_j A(Z_j)$, $\psi = \bigoplus_j \psi_j$, then $\pi_{Y \cap Z}^Z(\psi(E)) \subseteq \pi_{Y \cap Z}^Y(\eta(B))$, $\pi_Z(\mathcal{F}) \subset_\varepsilon \psi(E)$ and hence

$$\mathcal{F} \subset_\varepsilon \eta(B) \oplus_{Y \cap Z} \psi(E) = \chi(B \oplus_{\pi \eta, \pi \psi} E),$$

where χ is the isomorphism induced by the pair (η, ψ) . If we assume that each D_j is KK -stable, then we also have $KK(\varphi_j) = KK(\psi_j|_{D_j})$ for all $j \in J$.

Proof. Let $\mathcal{F} = \{a_1, \dots, a_r\} \subset A$ be as in the statement. By (i), for each $j \in J$ we find $\{c_1^{(j)}, \dots, c_r^{(j)}\} \subseteq \mathcal{H}_j$ such that $\|\varphi_j(c_i^{(j)}) - \pi_{Z_j}(a_i)\| < \varepsilon/2$ for all i . Consider the $C(X)$ -algebra $A \oplus_Y \eta(B) \subset A$. From (iii), Lemma 2.1(iv) and Lemma 2.5 we obtain

$$\varphi_j(\mathcal{G}_j) \subset_{\delta_j} \pi_{Z_j}(A \oplus_Y \eta(B)).$$

Applying Proposition 3.7 we perturb φ_j to a $*$ -homomorphism $\psi_j : D_j \rightarrow \pi_{Z_j}(A \oplus_Y \eta(B))$ satisfying (6), and hence such that $\|\varphi_j(c_i^{(j)}) - \psi_j(c_i^{(j)})\| < \varepsilon/2$, for all i, j . Therefore

$$\|\psi_j(c_i^{(j)}) - \pi_{Z_j}(a_i)\| \leq \|\psi_j(c_i^{(j)}) - \varphi_j(c_i^{(j)})\| + \|\varphi_j(c_i^{(j)}) - \pi_{Z_j}(a_i)\| < \varepsilon.$$

This shows that $\pi_{Z_j}(\mathcal{F}) \subset_\varepsilon \psi_j(D_j)$. From (6) and Lemma 4.1 we obtain that each $(\psi_j)_x$ is injective. We extend ψ_j to a $C(Z_j)$ -linear $*$ -monomorphism $\psi_j : C(Z_j) \otimes D_j \rightarrow$

$\pi_{Z_j}(A \oplus_Y \eta(B))$ and then we define E , ψ and Z as in the statement. In this way we obtain that $\psi : E \rightarrow (A \oplus_Y \eta(B))(Z) \subset A(Z)$ satisfies

$$\pi_Z(\mathcal{F}) \subset_\varepsilon \psi(E). \tag{7}$$

The property $\psi(E) \subset (A \oplus_Y \eta(B))(Z)$ is equivalent to $\pi_{Y \cap Z}^Z(\psi(E)) \subseteq \pi_{Y \cap Z}^Y(\eta(B))$ by Lemma 2.6(b). Finally, from (ii), (7) and Lemma 2.6(c) we get $\mathcal{F} \subset_\varepsilon \eta(B) \oplus_{Y \cap Z} \psi(E)$. \square

Let \mathcal{C} be as in Lemma 4.2. Let A be a $C(X)$ -algebra, let $\mathcal{F} \subset A$ be a finite set and let $\varepsilon > 0$. An $(\mathcal{F}, \varepsilon, \mathcal{C})$ -approximation of A

$$\alpha = \{ \mathcal{F}, \varepsilon, \{ U_i, \varphi_i : D_i \rightarrow A(U_i), \mathcal{H}_i, \mathcal{G}_i, \delta_i \}_{i \in I} \}, \tag{8}$$

is a collection with the following properties: $(U_i)_{i \in I}$ is a finite family of closed subsets of X , whose interiors cover X and $(D_i)_{i \in I}$ are C^* -algebras in \mathcal{C} ; for each $i \in I$, $\varphi_i : D_i \rightarrow A(U_i)$ is a $*$ -homomorphism such that $(\varphi_i)_x$ is injective for all $x \in U_i$; $\mathcal{H}_i \subset D_i$ is a finite set such that $\pi_{U_i}(\mathcal{F}) \subset_{\varepsilon/2} \varphi_i(\mathcal{H}_i)$ and such that for each direct summand of D_i there is an element of \mathcal{H}_i of norm $\geq \varepsilon$ which is contained and is full in that summand; the finite set $\mathcal{G}_i \subset D_i$ and $\delta_i > 0$ are given by Proposition 3.7 applied to the weakly semiprojective C^* -algebra D_i for the input data \mathcal{H}_i and $\varepsilon/2$; if D_i is KK -stable, then \mathcal{G}_i and δ_i are chosen such that the second part of Proposition 3.7 also applies.

Lemma 4.5. *Let A and \mathcal{C} be as in Lemma 4.2. Suppose that each fiber of A admits an exhaustive sequence of C^* -algebras isomorphic to C^* -algebras in \mathcal{C} . Then for any finite subset \mathcal{F} of A and any $\varepsilon > 0$ there is an $(\mathcal{F}, \varepsilon, \mathcal{C})$ -approximation of A . Moreover, if A , D and σ are as in Lemma 4.3 and $\sigma_x \in KK(D, A(x))^{-1}$ for all $x \in X$, then there is an $(\mathcal{F}, \varepsilon, \mathcal{C})$ -approximation of A such that $\mathcal{C} = \{D\}$ and $KK(\varphi_i) = \sigma_{U_i}$ for all $i \in I$.*

Proof. Since X is compact, this is an immediate consequence of Lemmas 4.2, 4.3 and Proposition 3.7. \square

It is useful to consider the following operation of restriction. Suppose that Y is a closed subspace of X and let $(V_j)_{j \in J}$ be a finite family of closed subsets of Y which refines the family $(Y \cap U_i)_{i \in I}$ and such that the interiors of the V_j 's form a cover of Y . Let $\iota : J \rightarrow I$ be a map such that $V_j \subseteq Y \cap U_{\iota(j)}$. Define

$$\iota^*(\alpha) = \{ \pi_Y(\mathcal{F}), \varepsilon, \{ V_j, \pi_{V_j} \varphi_{\iota(j)} : D_{\iota(j)} \rightarrow A(V_j), \mathcal{H}_{\iota(j)}, \mathcal{G}_{\iota(j)}, \delta_{\iota(j)} \}_{j \in J} \}.$$

It is obvious that $\iota^*(\alpha)$ is a $(\pi_Y(\mathcal{F}), \varepsilon, \mathcal{C})$ -approximation of $A(Y)$. The operation $\alpha \mapsto \iota^*(\alpha)$ is useful even in the case $X = Y$. Indeed, by applying this procedure we can refine the cover of X that appears in a given $(\mathcal{F}, \varepsilon, \mathcal{C})$ -approximation of A .

An $(\mathcal{F}, \varepsilon, \mathcal{C})$ -approximation α (as in (8)) is subordinated to an $(\mathcal{F}', \varepsilon', \mathcal{C})$ -approximation, $\alpha' = \{ \mathcal{F}', \varepsilon', \{ U_{i'}, \varphi_{i'} : D_{i'} \rightarrow A(U_{i'}), \mathcal{H}_{i'}, \mathcal{G}_{i'}, \delta_{i'} \}_{i' \in I'} \}$, written $\alpha < \alpha'$, if

- (i) $\mathcal{F} \subseteq \mathcal{F}'$,
- (ii) $\varphi_i(\mathcal{G}_i) \subseteq \pi_{U_i}(\mathcal{F}')$ for all $i \in I$, and
- (iii) $\varepsilon' < \min(\{\varepsilon\} \cup \{\delta_i, i \in I\})$.

Let us note that, with notation as above, we have $\iota^*(\alpha) < \iota^*(\alpha')$ whenever $\alpha < \alpha'$.

The following theorem is the crucial technical result of our paper. It provides an approximation of continuous $C(X)$ -algebras by subalgebras of category $\leq \dim(X)$.

Theorem 4.6. *Let \mathcal{C} be a class consisting of finite direct sums of weakly semiprojective simple C^* -algebras. Let X be a finite dimensional compact metrizable space and let A be a separable continuous $C(X)$ -algebra the fibers of which admit exhaustive sequences of C^* -algebras isomorphic to C^* -algebras in \mathcal{C} . For any finite set $\mathcal{F} \subset A$ and any $\varepsilon > 0$ there exist $n \leq \dim(X)$ and an n -fibered \mathcal{C} -monomorphism (ψ_0, \dots, ψ_n) into A which induces a $*$ -monomorphism $\eta : A(\psi_0, \dots, \psi_n) \rightarrow A$ such that $\mathcal{F} \subset_\varepsilon \eta(A(\psi_0, \dots, \psi_n))$.*

Proof. By Lemma 4.5, for any finite set $\mathcal{F} \subset A$ and any $\varepsilon > 0$ there is an $(\mathcal{F}, \varepsilon, \mathcal{C})$ -approximation of A . Moreover, for any finite set $\mathcal{F} \subset A$, any $\varepsilon > 0$ and any n , there is a sequence $\{\alpha_k : 0 \leq k \leq n\}$ of $(\mathcal{F}_k, \varepsilon_k, \mathcal{C})$ -approximations of A such that $(\mathcal{F}_0, \varepsilon_0) = (\mathcal{F}, \varepsilon)$ and α_k is subordinated to α_{k+1} :

$$\alpha_0 < \alpha_1 < \dots < \alpha_n.$$

Indeed, assume that α_k was constructed. Let us choose a finite set \mathcal{F}_{k+1} which contains \mathcal{F}_k and liftings to A of all the elements in $\bigcup_{i_k \in I_k} \varphi_{i_k}(\mathcal{G}_{i_k})$. This choice takes care of the above conditions (i) and (ii). Next we choose ε_{k+1} sufficiently small such that (iii) is satisfied. Let α_{k+1} be an $(\mathcal{F}_{k+1}, \varepsilon_{k+1}, \mathcal{C})$ -approximation of A given by Lemma 4.5. Then obviously $\alpha_k < \alpha_{k+1}$. Fix a tower of approximations of A as above where $n = \dim(X)$.

By [4, Lemma 3.2], for every open cover \mathcal{V} of X there is a finite open cover \mathcal{U} which refines \mathcal{V} and such that the set \mathcal{U} can be partitioned into $n + 1$ nonempty subsets consisting of elements with pairwise disjoint closures. Since we can refine simultaneously the covers that appear in a finite family $\{\alpha_k : 0 \leq k \leq n\}$ of approximations while preserving subordination, we may arrange not only that all α_k share the same cover $(U_i)_{i \in I}$, but moreover, that the cover $(U_i)_{i \in I}$ can be partitioned into $n + 1$ subsets $\mathcal{U}_0, \dots, \mathcal{U}_n$ consisting of mutually disjoint elements. For definiteness, let us write $\mathcal{U}_k = \{U_{i_k} : i_k \in I_k\}$. Now for each k we consider the closed subset of X

$$Y_k = \bigcup_{i_k \in I_k} U_{i_k},$$

the map $\iota_k : I_k \rightarrow I$ and the $(\pi_{Y_k}(\mathcal{F}_k), \varepsilon_k, \mathcal{C})$ -approximation of $A(Y_k)$, induced by α_k , which is of the form

$$\iota_k^*(\alpha_k) = \left\{ \pi_{Y_k}(\mathcal{F}_k), \varepsilon, \left\{ U_{i_k}, \varphi_{i_k} : D_{i_k} \rightarrow A(U_{i_k}), \mathcal{H}_{i_k}, \mathcal{G}_{i_k}, \delta_{i_k} \right\}_{i_k \in I_k} \right\},$$

where each U_{i_k} is nonempty. We have

$$\pi_{U_{i_k}}(\mathcal{F}_k) \subset_{\varepsilon_k/2} \varphi_{i_k}(\mathcal{H}_{i_k}), \tag{9}$$

by construction. Since $\alpha_k < \alpha_{k+1}$ we obtain

$$\mathcal{F}_k \subseteq \mathcal{F}_{k+1}, \tag{10}$$

$$\varphi_{i_k}(\mathcal{G}_{i_k}) \subseteq \pi_{U_{i_k}}(\mathcal{F}_{k+1}), \quad \text{for all } i_k \in I_k, \tag{11}$$

$$\varepsilon_{k+1} < \min(\{\varepsilon_k\} \cup \{\delta_{i_k}, i_k \in I_k\}). \tag{12}$$

Set $X_k = Y_k \cup \dots \cup Y_n$ and $E_k = \bigoplus_{i_k} C(U_{i_k}) \otimes D_{i_k}$ for $0 \leq k \leq n$. We shall construct a sequence of $C(Y_k)$ -linear $*$ -monomorphisms, $\psi_k : E_k \rightarrow A(Y_k)$, $k = n, \dots, 0$, such that (ψ_k, \dots, ψ_n) is an $(n - k)$ -fibered monomorphism into $A(X_k)$. Each map

$$\psi_k = \bigoplus_{i_k} \psi_{i_k} : E_k \rightarrow A(Y_k) = \bigoplus_{i_k} A(U_{i_k})$$

will have components $\psi_{i_k} : C(U_{i_k}) \otimes D_{i_k} \rightarrow A(U_{i_k})$ whose restrictions to D_{i_k} will be perturbations of $\varphi_{i_k} : D_{i_k} \rightarrow A(U_{i_k})$, $i_k \in I_k$. We shall construct the maps ψ_k by induction on decreasing k such that if $B_k = A(X_k)(\psi_k, \dots, \psi_n)$ and $\eta_k : B_k \rightarrow A(X_k)$ is the map induced by the $(n - k)$ -fibered monomorphism (ψ_k, \dots, ψ_n) , then

$$\pi_{X_{k+1} \cap U_{i_k}}(\psi_{i_k}(D_{i_k})) \subseteq \pi_{X_{k+1} \cap U_{i_k}}(\eta_{k+1}(B_{k+1})), \quad \forall i_k \in I_k, \tag{13}$$

and

$$\pi_{X_k}(\mathcal{F}_k) \subseteq_{\varepsilon_k} \eta_k(B_k). \tag{14}$$

Note that (13) is equivalent to

$$\pi_{X_{k+1} \cap Y_k}(\psi_k(E_k)) \subseteq \pi_{X_{k+1} \cap Y_k}(\eta_{k+1}(B_{k+1})). \tag{15}$$

For the first step of induction, $k = n$, we choose $\psi_n = \bigoplus_{i_n} \tilde{\varphi}_{i_n}$ where $\tilde{\varphi}_{i_n} : C(U_{i_n}) \otimes D_{i_n} \rightarrow A(U_{i_n})$ are $C(U_{i_n})$ -linear extensions of the original φ_{i_n} . Then $B_n = E_n$ and $\eta_n = \psi_n$. Assume that $\psi_n, \dots, \psi_{k+1}$ were constructed and that they have the desired properties. We shall construct now ψ_k . Condition (14) formulated for $k + 1$ becomes

$$\pi_{X_{k+1}}(\mathcal{F}_{k+1}) \subseteq_{\varepsilon_{k+1}} \eta_{k+1}(B_{k+1}). \tag{16}$$

Since $\varepsilon_{k+1} < \delta_{i_k}$, by using (11) and (16) we obtain

$$\pi_{X_{k+1} \cap U_{i_k}}(\varphi_{i_k}(\mathcal{G}_{i_k})) \subseteq_{\delta_{i_k}} \pi_{X_{k+1} \cap U_{i_k}}(\eta_{k+1}(B_{k+1})), \quad \text{for all } i_k \in I_k. \tag{17}$$

Since $\mathcal{F}_k \subseteq \mathcal{F}_{k+1}$ and $\varepsilon_{k+1} < \varepsilon_k$, condition (16) gives

$$\pi_{X_{k+1}}(\mathcal{F}_k) \subseteq_{\varepsilon_k} \eta_{k+1}(B_{k+1}). \tag{18}$$

Conditions (9), (17) and (18) enable us to apply Lemma 4.4 and perturb $\tilde{\varphi}_{i_k}$ to a $*$ -monomorphism $\psi_{i_k} : C(U_{i_k}) \otimes D_{i_k} \rightarrow A(U_{i_k})$ satisfying (13) and (14) and such that

$$KK(\psi_{i_k}|_{D_{i_k}}) = KK(\varphi_{i_k}) \tag{19}$$

if the algebras in \mathcal{C} are assumed to be KK -stable. We set $\psi_k = \bigoplus_{i_k} \psi_{i_k}$ and this completes the construction of (ψ_0, \dots, ψ_n) . Condition (14) for $k = 0$ gives $\mathcal{F} \subseteq_{\varepsilon} \eta_0(B_0) = \eta(A(\psi_0, \dots, \psi_n))$.

Thus (ψ_0, \dots, ψ_n) satisfies the conclusion of the theorem. Finally let us note that it can happen that $X_k = X$ for some $k > 0$. In this case $\mathcal{F} \subset_\varepsilon A(\psi_k, \dots, \psi_n)$ and for this reason we write $n \leq \dim(X)$ in the statement of the theorem. \square

Proposition 4.7. *Let X be a finite dimensional compact metrizable space and let A be a separable continuous $C(X)$ -algebra the fibers of which are stable Kirchberg algebras. Let D be a KK -semiprojective stable Kirchberg algebra and suppose that there exists $\sigma \in KK(D, A)$ such that $\sigma_x \in KK(D, A(x))^{-1}$ for all $x \in X$. For any finite subset \mathcal{F} of A and any $\varepsilon > 0$ there is an n -fibered \mathcal{C} -monomorphism (ψ_0, \dots, ψ_n) into A such that $n \leq \dim(X)$, $\mathcal{C} = \{D\}$, and each component $\psi_i : C(Y_i) \otimes D \rightarrow A(Y_i)$ satisfies $KK(\psi_i) = \sigma_{Y_i}$, $i = 0, \dots, n$. Moreover, if $\eta : A(\psi_0, \dots, \psi_n) \rightarrow A$ is the induced $*$ -monomorphism, then $\mathcal{F} \subset_\varepsilon \eta(A(\psi_0, \dots, \psi_n))$ and $KK(\eta_x)$ is a KK -equivalence for each $x \in X$.*

Proof. We repeat the proof of Theorem 4.6 while using only $(\mathcal{F}_i, \varepsilon_i, \{D\})$ -approximations of A provided by the second part of Lemma 4.5. The outcome will be an n -fibered $\{D\}$ -monomorphism (ψ_0, \dots, ψ_n) into A such that $\mathcal{F} \subset_\varepsilon \eta(A(\psi_0, \dots, \psi_n))$. Moreover we can arrange that $KK(\psi_i) = \sigma_{Y_i}$ for all $i = 0, \dots, n$, by (19), since $KK(\varphi_{i_k}) = \sigma_{U_{i_k}}$ by Lemma 4.5. If $x \in X$, and $i = \min\{k : x \in Y_k\}$, then $\eta_x \equiv (\psi_i)_x$, and hence $KK(\eta_x)$ is a KK -equivalence. \square

Remark 4.8. Let us point out that we can strengthen the conclusion of Theorem 4.6 and Proposition 4.7 as follows. Fix a metric d for the topology of X . Then we may arrange that there is a closed cover $\{Y'_0, \dots, Y'_n\}$ of X and a number $\ell > 0$ such that $\{x : d(x, Y'_i) \leq \ell\} \subset Y_i$ for $i = 0, \dots, n$. Indeed, when we choose the finite closed cover $\mathcal{U} = (U_i)_{i \in I}$ of X in the proof of Theorem 4.6 which can be partitioned into $n + 1$ subsets $\mathcal{U}_0, \dots, \mathcal{U}_n$ consisting of mutually disjoint elements, as given by [4, Lemma 3.2], and which refines all the covers $\mathcal{U}(\alpha_0), \dots, \mathcal{U}(\alpha_n)$ corresponding to $\alpha_0, \dots, \alpha_n$, we may assume that \mathcal{U} also refines the covers given by the interiors of the elements of $\mathcal{U}(\alpha_0), \dots, \mathcal{U}(\alpha_n)$. Since each U_i is compact and I is finite, there is $\ell > 0$ such that if $V_i = \{x : d(x, U_i) \leq \ell\}$, then the cover $\mathcal{V} = (V_i)_{i \in I}$ still refines all of $\mathcal{U}(\alpha_0), \dots, \mathcal{U}(\alpha_n)$ and for each $k = 0, \dots, n$, the elements of $\mathcal{V}_k = \{V_i : U_i \in \mathcal{U}_k\}$, are still mutually disjoint. We shall use the cover \mathcal{V} rather than \mathcal{U} in the proof of the two theorems and observe that $Y'_k \stackrel{def}{=} \bigcup_{i_k \in I_k} U_{i_k} \subset \bigcup_{i_k \in I_k} V_{i_k} = Y_k$ has the desired property. Finally let us note that if we define $\psi'_i : E(Y'_i) \rightarrow A(Y'_i)$ by $\psi'_i = \pi_{Y'_i} \psi_i$, then $(\psi'_0, \dots, \psi'_n)$ is an n -fibered \mathcal{C} -monomorphism into A which satisfies the conclusion of Theorem 4.6 and Proposition 4.7 since $\pi_{Y'_i}(\mathcal{F}) \subset_\varepsilon \psi'_i(E_i)$ for all $i = 0, \dots, n$ and $X = \bigcup_{i=1}^n Y'_i$.

5. Representing $C(X)$ -algebras as inductive limits

We have seen that Theorem 4.6 yields exhaustive sequences for certain $C(X)$ -algebras. In this section we show how to pass from an exhaustive sequence to a nested exhaustive sequence using semiprojectivity. The remainder of the paper does not depend on this section.

Proposition 5.1. *Let X, A and \mathcal{C} be as in Theorem 4.6. Let (ψ_0, \dots, ψ_n) be an n -fibered \mathcal{C} -monomorphism into A with components $\psi_i : E_i \rightarrow A(Y_i)$. Let $\mathcal{F}_i \subset E_i, \mathcal{F} \subset A(\psi_0, \dots, \psi_n)$ be finite sets and let $\varepsilon > 0$. Then there are finite sets $\mathcal{G}_i \subset E_i$ and $\delta_i > 0, i = 0, \dots, n$, such that for any $C(X)$ -subalgebra $A' \subset A$ which satisfies $\psi_i(\mathcal{G}_i) \subset_{\delta_i} A'(Y_i), i = 0, \dots, n$, there is an n -fibered \mathcal{C} -monomorphism $(\psi'_0, \dots, \psi'_n)$ into A' , with $\psi'_i : E_i \rightarrow A'(Y_i)$ and such that (i) $\|\psi_i(a) - \psi'_i(a)\| < \varepsilon$ for all $a \in \mathcal{F}_i$ and all $i \in \{0, \dots, n\}$, (ii) $(\psi_j)_x^{-1}(\psi_i)_x = (\psi'_j)_x^{-1}(\psi'_i)_x$*

for all $x \in Y_i \cap Y_j$ and $0 \leq i \leq j \leq n$. Moreover $A(\psi_0, \dots, \psi_n) = A'(\psi'_0, \dots, \psi'_n)$ and the maps $\eta : A(\psi_0, \dots, \psi_n) \rightarrow A$ and $\eta' : A'(\psi'_0, \dots, \psi'_n) \rightarrow A'$ induced by (ψ_0, \dots, ψ_n) and $(\psi'_0, \dots, \psi'_n)$ satisfy (iii) $\|\eta(a) - \eta'(a)\| < \varepsilon$ for all $a \in \mathcal{F}$.

Proof. Let us observe that if we prove (i) and (ii) then (iii) will follow by enlarging the sets \mathcal{F}_i so that $p_i(\mathcal{F}) \subset \mathcal{F}_i$, where $p_i : A(\psi_0, \dots, \psi_n) \rightarrow E_i$ are the coordinate maps. We proceed now with the proof of (i) and (ii) by making some simplifications. We may assume that $E_0 = C(Y_0) \otimes D_0$ with $D_0 \in \mathcal{C}$ since the perturbations corresponding to disjoint closed sets can be done independently of each other. Without any loss of generality, we may assume that $\mathcal{F}_0 \subset D_0$ since we are working with morphisms on E_0 which are $C(Y_0)$ -linear. We also enlarge \mathcal{F}_0 so that for each direct summand C of D_0 , \mathcal{F}_0 contains an element c which is full in C and such that $\|c\| \geq 2\varepsilon$.

The proof is by induction on n . If $n = 0$ the statement follows from Proposition 3.7 and Lemma 4.1. Assume now that the statement is true for $n - 1$. Let $E_i, \psi_i, A, A', \mathcal{F}_i, 1 \leq i \leq n$ and ε be as in the statement. For $0 \leq i < j \leq n$ let $\eta_{j,i} : E_i(Y_i \cap Y_j) \rightarrow E_j(Y_i \cap Y_j)$ be the $*$ -homomorphism of $C(Y_i \cap Y_j)$ -algebras defined fiberwise by $(\eta_{j,i})_x = (\psi_j)_x^{-1}(\psi_i)_x$.

Let \mathcal{G}_0 and δ_0 be given by Proposition 3.8 applied to the C^* -algebra D_0 for the input data \mathcal{F}_0 and ε . For each $1 \leq j \leq n$ choose a finite subset \mathcal{H}_j of E_j whose restriction to $Y_j \cap Y_0$ contains $\eta_{j,0}(\mathcal{G}_0)$. Consider the sets $\mathcal{F}'_j := \mathcal{F}_j \cup \mathcal{H}_j, 1 \leq j \leq n$ and the number $\varepsilon' = \min\{\delta_0, \varepsilon\}$. Let $\mathcal{G}_1, \dots, \mathcal{G}_n$ and $\delta_1, \dots, \delta_n$ be given by the inductive assumption for $n - 1$ applied to $A(X_1), A'(X_1), \psi_j, \mathcal{F}'_j, 1 \leq j \leq n$ and ε' , where $X_1 = Y_1 \cup \dots \cup Y_n$.

We need to show that $\mathcal{G}_0, \mathcal{G}_1, \dots, \mathcal{G}_n$ and $\delta_0, \delta_1, \dots, \delta_n$ satisfy the statement. By the inductive step there exists an $(n - 1)$ -fibered \mathcal{C} -monomorphism $(\psi'_1, \dots, \psi'_n)$ into $A'(X_1)$ with components $\psi'_j : E_j \rightarrow A'(Y_j)$ such that

- (a) $\|\psi_j(a) - \psi'_j(a)\| < \varepsilon' = \min\{\delta_0, \varepsilon\}$ for all $a \in \mathcal{F}_j \cup \mathcal{H}_j$ and all $1 \leq j \leq n$,
- (b) $(\psi_j)_x^{-1}(\psi_i)_x = (\psi'_j)_x^{-1}(\psi'_i)_x$ for all $x \in Y_i \cap Y_j$ and $1 \leq i \leq j \leq n$.

The condition (b) enables to define a $*$ -homomorphism $\varphi : E_0 \rightarrow A'(Y_0 \cap X_1)$ with fiber maps $\varphi_x = (\psi'_j)_x(\psi_j^{-1})_x(\psi_0)_x$ for $x \in Y_0 \cap Y_j$ and $1 \leq j \leq n$.

Let us observe that $\psi_0 : E_0 \rightarrow A(Y_0)$ is an approximate lifting of φ . More precisely we have $\|\pi_{X_1 \cap Y_0}^{Y_0} \psi_0(a) - \varphi(a)\| < \delta_0$ for all $a \in \mathcal{G}_0$. Indeed, for $x \in Y_0 \cap Y_j, 1 \leq j \leq n$ and $a \in \mathcal{G}_0$ we have

$$\begin{aligned} \|(\psi_0)_x(a(x)) - (\psi'_j)_x(\psi_j^{-1})_x(\psi_0)_x(a(x))\| &= \|(\psi_j)_x(\eta_{j,0})_x(a(x)) - (\psi'_j)_x(\eta_{j,0})_x(a(x))\| \\ &\leq \sup_{h \in \mathcal{H}_j} \|\psi_j(h) - \psi'_j(h)\| < \varepsilon' \leq \delta_0. \end{aligned}$$

Since we also have $\psi_0(\mathcal{G}_0) \subset_{\delta_0} A'(Y_0)$ by hypothesis, it follows from Proposition 3.8 that there exists $\psi'_0 : D_0 \rightarrow A(Y_0)$ such that $\|\psi'_0(a) - \psi_0(a)\| < \varepsilon$ for all $a \in \mathcal{F}_0$ and $\pi_{Y_0 \cap X_1}^{Y_0} \psi'_0 = \varphi$. By Lemma 4.1 each $(\psi'_0)_x$ is injective since each $(\psi_0)_x$ is injective. The $C(Y_0)$ -linear extension of ψ'_0 to E_0 satisfies $(\psi_j)_x^{-1}(\psi_0)_x = (\psi'_j)_x^{-1}(\psi'_0)_x$ for all $x \in Y_0 \cap Y_j$ and $1 \leq j \leq n$ and this completes the proof of (ii). Condition (i) follows from (b). \square

The following result gives an inductive limit representation for continuous $C(X)$ -algebras whose fibers are inductive limits of finite direct sums of simple semiprojective C^* -algebras. For

example the fibers can be arbitrary AF-algebras or Kirchberg algebras which satisfy the UCT and whose K_1 -groups are torsion free. Indeed, by [29, Prop. 8.4.13], these algebras are isomorphic to inductive limits of sequences of Kirchberg algebras (D_n) with finitely generated K-theory groups and torsion free K_1 -groups. The algebras D_n are semiprojective by [33].

Theorem 5.2. *Let \mathcal{C} be a class consisting of finite direct sums of semiprojective simple C^* -algebras. Let X be a finite dimensional compact metrizable space and let A be a separable continuous $C(X)$ -algebra such that all its fibers admit exhaustive sequences consisting of C^* -algebras isomorphic to C^* -algebras in \mathcal{C} . Then A is isomorphic to the inductive limit of a sequence of continuous $C(X)$ -algebras A_k such that $\text{cat}_{\mathcal{C}}(A_k) \leq \dim(X)$.*

Proof. By Theorem 4.6 and Proposition 5.1 we find a sequence $(\psi_0^{(k)}, \dots, \psi_n^{(k)})$ of n -fibered \mathcal{C} -monomorphisms into A which induces $*$ -monomorphisms $\eta^{(k)} : A_k = A(\psi_0^{(k)}, \dots, \psi_n^{(k)}) \rightarrow A$ with the following properties. There is a sequence of finite sets $\mathcal{F}_k \subset A_k$ and a sequence of $C(X)$ -linear $*$ -monomorphisms $\mu_k : A_k \rightarrow A_{k+1}$ such that

- (i) $\|\eta^{(k+1)}\mu_k(a) - \eta^{(k)}(a)\| < 2^{-k}$ for all $a \in \mathcal{F}_k$ and all $k \geq 1$,
- (ii) $\mu_k(\mathcal{F}_k) \subset \mathcal{F}_{k+1}$ for all $k \geq 1$,
- (iii) $\bigcup_{j=k+1}^{\infty} (\mu_{j-1} \circ \dots \circ \mu_k)^{-1}(\mathcal{F}_j)$ is dense in A_k and $\bigcup_{j=k}^{\infty} \eta^{(j)}(\mathcal{F}_j)$ is dense in A for all $k \geq 1$.

Arguing as in the proof of [29, Prop. 2.3.2], one verifies that

$$\varphi_k(a) = \lim_{j \rightarrow \infty} \eta^{(j)} \circ (\mu_{j-1} \circ \dots \circ \mu_k)(a)$$

defines a sequence of $*$ -monomorphisms $\varphi_k : A_k \rightarrow A$ such that $\varphi_{k+1}\mu_k = \varphi_k$ and the induced map $\varphi : \varinjlim_k (A_k, \mu_k) \rightarrow A$ is an isomorphism of $C(X)$ -algebras. \square

Remark 5.3. By similar arguments one proves a unital version of Theorem 5.2.

6. When is a fibered product locally trivial

For C^* -algebras A, B we endow the space $\text{Hom}(A, B)$ of $*$ -homomorphisms with the point-norm topology. If X is a compact Hausdorff space, then $\text{Hom}(A, C(X) \otimes B)$ is homeomorphic to the space of continuous maps from X to $\text{Hom}(A, B)$ endowed with the compact-open topology. We shall identify a $*$ -homomorphism $\varphi \in \text{Hom}(A, C(X) \otimes B)$ with the corresponding continuous map $X \rightarrow \text{Hom}(A, B)$, $x \mapsto \varphi_x$, $\varphi_x(a) = \varphi(a)(x)$ for all $x \in X$ and $a \in A$. Let D be a C^* -algebra and let A be a $C(X)$ -algebra. If $\alpha : D \rightarrow A$ is a $*$ -homomorphism, let us denote by $\tilde{\alpha} : C(X) \otimes D \rightarrow A$ its (unique) $C(X)$ -linear extension and write $\tilde{\alpha} \in \text{Hom}_{C(X)}(C(X) \otimes D, A)$. For C^* -algebras D, B we shall make without further comment the following identifications

$$\text{Hom}_{C(X)}(C(X) \otimes D, C(X) \otimes B) \cong \text{Hom}(D, C(X) \otimes B) \cong C(X, \text{Hom}(D, B)).$$

For a C^* -algebra D we denote by $\text{End}(D)$ the set of full (and unital if D is unital) $*$ -endomorphisms of D and by $\text{End}(D)^0$ the path component of id_D in $\text{End}(D)$. Let us consider

$$\text{End}(D)^* = \{ \gamma \in \text{End}(D) : KK(\gamma) \in KK(D, D)^{-1} \}.$$

Proposition 6.1. *Let X be a compact metrizable space and let D be a KK -semiprojective Kirchberg algebra. Let $\alpha : D \rightarrow C(X) \otimes D$ be a full (and unital, if D is unital) $*$ -homomorphism such that $KK(\alpha_x) \in KK(D, D)^{-1}$ for all $x \in X$. Then there is a full $*$ -homomorphism $\Phi : D \rightarrow C(X \times [0, 1]) \otimes D$ such that $\Phi_{(x,0)} = \alpha_x$ and $\Phi_{(x,t)} \in \text{Aut}(D)$ for all $x \in X$ and $t \in (0, 1]$. Moreover, if $\Phi_1 : D \rightarrow C(X) \otimes D$ is defined by $\Phi_1(d)(x) = \Phi_{(x,1)}(d)$, for all $d \in D$ and $x \in X$, then $\alpha \approx_{uh} \Phi_1$.*

Proof. Since X is a metrizable compact space, X is homeomorphic to the projective limit of a sequence of finite simplicial complexes (X_i) by [13, Thm. 10.1, p. 284]. Since D is KK -semiprojective, $KK(D, \varinjlim C(X_i) \otimes D) = KK(D, C(X) \otimes D)$ by Theorem 3.12. By Theorem 3.1, there is i and a full (and unital if D is unital) $*$ -homomorphism $\varphi : D \rightarrow C(X_i) \otimes D$ whose KK -class maps to $KK(\alpha) \in KK(D, C(X) \otimes D)$. To summarize, we have found a finite simplicial complex Y , a continuous map $h : X \rightarrow Y$ and a continuous map $y \mapsto \varphi_y \in \text{End}(D)$, defined on Y , such that the full (and unital if D is unital) $*$ -homomorphism $h^*\varphi : D \rightarrow C(X) \otimes D$ corresponding to the continuous map $x \mapsto \varphi_{h(x)}$ satisfies $KK(h^*\varphi) = KK(\alpha)$. We may arrange that $h(X)$ intersects all the path components of Y by dropping the path components which are not intersected. Since $\alpha_x \in \text{End}(D)^*$ by hypothesis, and since $KK(\alpha_x) = KK(\varphi_{h(x)})$, we infer that $\varphi_y \in \text{End}(D)^*$ for all $y \in Y$. We shall find a continuous map $y \mapsto \psi_y \in \text{End}(D)^*$ defined on Y , such that the maps $y \mapsto \psi_y \varphi_y$ and $y \mapsto \varphi_y \psi_y$ are homotopic to the constant map ι that takes Y to id_D . It is clear that it suffices to deal separately with each path component of Y , so that for this part of the proof we may assume that Y is connected. Fix a point $z \in Y$. By [29, Thm. 8.4.1] there is $v \in \text{Aut}(D)$ such that $KK(v^{-1}) = KK(\varphi_z)$ and hence $KK(v\varphi_z) = KK(\text{id}_D)$. By Theorem 3.1, there is a unitary $u \in M(D)$ such that $uv\varphi_z(-)u^*$ is homotopic to id_D . Let us set $\theta = uv(-)u^* \in \text{Aut}(D)$ and observe that $\theta\varphi_z \in \text{End}(D)^0$. Since Y is path connected, it follows that the entire image of the map $y \mapsto \theta\varphi_y$ is contained in $\text{End}(D)^0$. Since $\text{End}(D)^0$ is a path connected H -space with unit element, it follows by [38, Thm. 2.4, p. 462] that the homotopy classes $[Y, \text{End}(D)^0]$ (with no condition on basepoints, since the action of the fundamental group $\pi_1(\text{End}(D)^0, \text{id}_D)$ is trivial by [38, 3.6, p. 166]) form a group under the natural multiplication. Therefore we find $y \mapsto \psi'_y \in \text{End}(D)^0$ such that $y \mapsto \psi'_y \theta\varphi_y$ and $y \mapsto \theta\varphi_y \psi'_y$ are homotopic to ι .

It follows that $y \mapsto \psi_y \stackrel{\text{def}}{=} \psi'_y \theta$ is the homotopic inverse of $y \mapsto \varphi_y$ in $[Y, \text{End}(D)^*]$. Composing with h we obtain that the maps $x \mapsto \varphi_{h(x)} \psi_{h(x)}$ and $x \mapsto \psi_{h(x)} \varphi_{h(x)}$ are homotopic to the constant map that takes X to id_D . By the homotopy invariance of KK -theory we obtain that

$$KK(\widetilde{h^*\varphi} h^*\psi) = KK(\widetilde{h^*\psi} h^*\varphi) = KK(\iota_D),$$

where $\widetilde{h^*\varphi}$ and $\widetilde{h^*\psi}$ denote the $C(X)$ -linear extensions of the corresponding maps and $\iota_D : D \rightarrow C(X) \otimes D$ is defined by $\iota_D(d) = 1_{C(X)} \otimes d$ for all $d \in D$. Let us recall that $KK(h^*\varphi) = KK(\alpha)$ and hence $KK(\widetilde{h^*\varphi}) = KK(\widetilde{\alpha})$. If we set $\Psi = h^*\psi$, then

$$KK(\widetilde{\alpha}\Psi) = KK(\widetilde{\Psi}\alpha) = KK(\iota_D).$$

By Theorem 3.1 $\widetilde{\alpha}\Psi \approx_u \iota_D$ and $\widetilde{\Psi}\alpha \approx_u \iota_D$, and hence $\widetilde{\alpha}\widetilde{\Psi} \approx_u \text{id}_{C(X) \otimes D}$ and $\widetilde{\Psi}\widetilde{\alpha} \approx_u \text{id}_{C(X) \otimes D}$. By [29, Cor. 2.3.4], there is an isomorphism $\Gamma : C(X) \otimes D \rightarrow C(X) \otimes D$ such that $\Gamma \approx_u \widetilde{\alpha}$. In particular Γ is $C(X)$ -linear and $\Gamma_x \in \text{Aut}(D)$ for all $x \in X$. Replacing Γ by $u\Gamma(\cdot)u^*$ for

some unitary $u \in M(C(X) \otimes D)$ we can arrange that $\Gamma|_D$ is arbitrarily close to α . Therefore $KK(\Gamma|_D) = KK(\alpha)$ since D is KK -stable. By Theorem 3.1 there is a continuous map $(0, 1] \rightarrow U(M(C(X) \otimes D))$, $t \mapsto u_t$, with the property that

$$\lim_{t \rightarrow 0} \|u_t \Gamma(a) u_t^* - \alpha(a)\| = 0, \quad \text{for all } a \in D.$$

Therefore the equation

$$\Phi_{(x,t)} = \begin{cases} \alpha_x, & \text{if } t = 0, \\ u_t(x) \Gamma_x u_t(x)^*, & \text{if } t \in (0, 1], \end{cases}$$

defines a continuous map $\Phi : X \times [0, 1] \rightarrow \text{End}(D)^*$ which extends α and such that $\Phi(X \times (0, 1]) \subset \text{Aut}(D)$. Since α is homotopic to Φ_1 , we have that $\alpha \approx_{uh} \Phi_1$ by Theorem 3.1. \square

Proposition 6.2. *Let X be a compact metrizable space and let D be a KK -semiprojective Kirchberg algebra. Let Y be a closed subset of X . Assume that a map $\gamma : Y \rightarrow \text{End}(D)^*$ extends to a continuous map $\alpha : X \rightarrow \text{End}(D)^*$. Then there is a continuous extension $\eta : X \rightarrow \text{End}(D)^*$ of γ , such that $\eta(X \setminus Y) \subset \text{Aut}(D)$.*

Proof. Since the map $x \rightarrow \alpha_x$ takes values in $\text{End}(D)^*$, by Proposition 6.1 there exists a continuous map $\Phi : X \times [0, 1] \rightarrow \text{End}(D)^*$ which extends α and such that $\Phi(X \times (0, 1]) \subset \text{Aut}(D)$. Let d be a metric for the topology of X such that $\text{diam}(X) \leq 1$. The equation $\eta(x) = \Phi(x, d(x, Y))$ defines a map on X that satisfies the conclusion of the proposition. \square

Lemma 6.3. *Let X be a compact metrizable space and let D be a KK -semiprojective Kirchberg algebra. Let Y be a closed subset of X . Let $\alpha : Y \times [0, 1] \cup X \times \{0\} \rightarrow \text{End}(D)$ be a continuous map such that $\alpha_{(x,0)} \in \text{End}(D)^*$ for all $x \in X$. Suppose that there is an open set V in X which contains Y and such that α extends to a continuous map $\alpha_V : V \times [0, 1] \cup X \times \{0\} \rightarrow \text{End}(D)$. Then there is $\eta : X \times [0, 1] \rightarrow \text{End}(D)^*$ such that η extends α and $\eta_{(x,t)} \in \text{Aut}(D)$ for all $x \in X \setminus Y$ and $t \in (0, 1]$.*

Proof. By Proposition 6.2 it suffices to find a continuous map $\widehat{\alpha} : X \times [0, 1] \rightarrow \text{End}(D)^*$ which extends α . Fix a metric d for the topology of X and define $\lambda : X \rightarrow [0, 1]$ by $\lambda(x) = d(x, X \setminus V)(d(x, X \setminus V) + d(x, Y))^{-1}$. Let us define $\widehat{\alpha} : X \times [0, 1] \rightarrow \text{End}(D)$ by $\widehat{\alpha}_{(x,t)} = \alpha_V(x, \lambda(x)t)$ and observe that $\widehat{\alpha}$ extends α . Finally, since $\widehat{\alpha}_{(x,t)}$ is homotopic to $\widehat{\alpha}_{(x,0)} = \alpha_{(x,0)}$, we conclude that the image of $\widehat{\alpha}$ is contained in $\text{End}(D)^*$. \square

Proposition 6.4. *Let X be a compact metrizable space and let D be a KK -semiprojective stable Kirchberg algebra. Let A be a separable $C(X)$ -algebra which is locally isomorphic to $C(X) \otimes D$. Suppose that there is $\sigma \in KK(D, A)$ such that $\sigma_x \in KK(D, A(x))^{-1}$ for all $x \in X$. Then there is an isomorphism of $C(X)$ -algebras $\psi : C(X) \otimes D \rightarrow A$ such that $KK(\psi|_D) = \sigma$.*

Proof. Since X is compact and A is locally trivial it follows that $\text{cat}_{\{D\}}(A) < \infty$. By Lemma 2.9, $A \cong pAp \otimes \mathcal{O}_\infty \otimes \mathcal{K}$ for some projection $p \in A$. By Theorem 3.1, there is a full $*$ -homomorphism $\varphi : D \rightarrow A$ such that $KK(\varphi) = \sigma$. We shall construct an isomorphism of $C(X)$ -algebras $\psi : C(X) \otimes D \rightarrow A$ such that ψ is homotopic to $\widetilde{\varphi}$, the $C(X)$ -linear extension

of φ . Moreover the homotopy $(H_t)_{t \in [0,1]}$ will have the property that $H_{(x,t)} : D \rightarrow A(x)$ is an isomorphism for all $x \in X$ and $t > 0$. We prove this by induction on numbers n with the property that there are two closed covers of X , W_1, \dots, W_n and Y_1, \dots, Y_n such that Y_i contained in the interior of W_i and $A(W_i) \cong C(W_i) \otimes D$ for $1 \leq i \leq n$. First we observe that the case $n = 1$ follows from Proposition 6.2. Let us now pass from $n - 1$ to n . Given two covers as above, there is yet another closed cover V_1, \dots, V_n of X such that V_i is a neighborhood of Y_i and W_i is a neighborhood of V_i for all $1 \leq i \leq n$. Set $Y = \bigcup_{i=1}^{n-1} Y_i$, $V = \bigcup_{i=1}^{n-1} V_i$ and $W = \bigcup_{i=1}^{n-1} W_i$. By the inductive hypothesis applied to $A(V)$, and the covers V_1, \dots, V_{n-1} and $W_1 \cap V, \dots, W_{n-1} \cap V$ there is a homotopy $h : D \rightarrow A(V) \otimes C[0, 1]$ such that $h_{(x,0)} = \varphi_x$ and $h_{(x,t)} : D \rightarrow A(x)$ is an isomorphism for all $(x, t) \in V \times (0, 1]$. Fix a trivialization $\nu : A(Y_{n+1}) \rightarrow C(Y_{n+1}) \otimes D$. Define a continuous map $\alpha : (V \cap Y_{n+1}) \times [0, 1] \cup Y_{n+1} \times \{0\} \rightarrow \text{End}(D)$ by setting $\alpha_{(x,t)} = \nu_x h_{(x,t)}$ if $(x, t) \in (V \cap Y_{n+1}) \times [0, 1]$ and $\alpha_{(x,0)} = \nu_x \varphi_x$ if $x \in Y_{n+1}$. Since $V \cap Y_{n+1}$ is a neighborhood of $Y \cap Y_{n+1}$ in Y_{n+1} and since $\nu_x \varphi_x \in \text{End}(D)^*$ for all $x \in Y_{n+1}$, by Lemma 6.3 there is a continuous map $\eta : Y_{n+1} \times [0, 1] \rightarrow \text{End}(D)^*$ which extends the restriction of α to $(Y \cap Y_{n+1}) \times [0, 1] \cup Y_{n+1} \times \{0\}$. We conclude the construction of the desired homotopy by defining $H : D \rightarrow A(X) \otimes C[0, 1]$ by $H_{(x,t)} = h_{(x,t)}$ for $(x, t) \in Y \times [0, 1]$ and $H_{(x,t)} = \nu_x^{-1} \eta_{(x,t)}$ for $(x, t) \in Y_{n+1} \times [0, 1]$. \square

Lemma 6.5. *Let D be a KK-semiprojective stable Kirchberg algebra. Let X be a compact metrizable space and Y, Z be closed subsets of X such that $X = Y \cup Z$. Suppose that $\gamma : D \rightarrow C(Y \cap Z) \otimes D$ is a full $*$ -homomorphism which admits a lifting to a full $*$ -homomorphism $\alpha : D \rightarrow C(Y) \otimes D$ such that $\alpha_x \in \text{End}(D)^*$ for all $x \in Y$. Then the pullback $C(Y) \otimes D \oplus_{\pi_{Y \cap Z}, \tilde{\gamma}_{\pi_{Y \cap Z}}} C(Z) \otimes D$ is isomorphic to $C(X) \otimes D$.*

Proof. By Proposition 6.2 there is a $*$ -homomorphism $\eta : D \rightarrow C(Y) \otimes D$ such that $\eta_x = \gamma_x$ for $x \in Y \cap Z$ and such that $\eta_x \in \text{Aut}(D)$ for $x \in Y \setminus Z$. Using the short five-lemma one checks immediately that the triplet $(\tilde{\eta}, \tilde{\gamma}, \text{id}_{C(Z) \otimes D})$ defines a $C(X)$ -linear isomorphism:

$$C(X) \otimes D = C(Y) \otimes D \oplus_{\pi_{Y \cap Z}, \pi_{Y \cap Z}} C(Z) \otimes D \rightarrow C(Y) \otimes D \oplus_{\pi_{Y \cap Z}, \tilde{\gamma}_{\pi_{Y \cap Z}}} C(Z) \otimes D. \quad \square$$

Lemma 6.6. *Let D be a KK-semiprojective stable Kirchberg algebra. Let Y, Z and Z' be closed subsets of a compact metrizable space X such that Z' is a neighborhood of Z and $X = Y \cup Z$. Let B be a $C(Y)$ -algebra locally isomorphic to $C(Y) \otimes D$ and let E be a $C(Z')$ -algebra locally isomorphic to $C(Z') \otimes D$. Let $\alpha : E(Y \cap Z') \rightarrow B(Y \cap Z')$ be a $*$ -monomorphism of $C(Y \cap Z')$ -algebras such that $\text{KK}(\alpha_x) \in \text{KK}(E(x), B(x))^{-1}$ for all $x \in Y \cap Z'$. If $\gamma = \alpha_{Y \cap Z}$, then $B(Y) \oplus_{\pi_{Y \cap Z}, \gamma_{\pi_{Y \cap Z}}} E(Z)$ is locally isomorphic to $C(X) \otimes D$.*

Proof. Since we are dealing with a local property, we may assume that $B = C(Y) \otimes D$ and $E = C(Z') \otimes D$. To simplify notation we let π stand for both $\pi_{Y \cap Z}^Y$ and $\pi_{Y \cap Z}^Z$ in the sequel. Let us denote by H the $C(X)$ -algebra $C(Y) \otimes D \oplus_{\pi, \gamma \pi} C(Z) \otimes D$. We must show that H is locally trivial. Let $x \in X$. If $x \notin Z$, then there is a closed neighborhood V of x which does not intersect Z , and hence the restriction of H to V is isomorphic to $C(V) \otimes D$, as it follows immediately from the definition of H . It remains to consider the case when $x \in Z$. Now Z' is a closed neighborhood of x in X and the restriction of H to Z' is isomorphic to $C(Y \cap Z') \otimes D \oplus_{\pi, \gamma \pi} C(Z) \otimes D$. Since $\gamma : Y \cap Z \rightarrow \text{End}(D)^*$ admits a continuous extension $\alpha : Y \cap Z' \rightarrow \text{End}(D)^*$, it follows that $H(Z')$ is isomorphic to $C(Z') \otimes D$ by Lemma 6.5. \square

Proposition 6.7. *Let X, A, D and σ be as in Proposition 4.7. For any finite subset \mathcal{F} of A and any $\varepsilon > 0$ there is a $C(X)$ -algebra B which is locally isomorphic to $C(X) \otimes D$ and there exists a $C(X)$ -linear $*$ -monomorphism $\eta : B \rightarrow A$ such that $\mathcal{F} \subset_\varepsilon \eta(B)$ and $KK(\eta_x) \in KK(B(x), A(x))^{-1}$ for all $x \in X$.*

Proof. Let $\psi_k : E_k = C(Y_k) \otimes D \rightarrow A(Y_k), k = 0, \dots, n$, be as in the conclusion of Proposition 4.7, strengthen as in Remark 4.8. Therefore we may assume that there is another n -fibered $\{D\}$ -monomorphism $(\psi'_0, \dots, \psi'_n)$ into A such that $\psi'_k : C(Y'_k) \otimes D \rightarrow A(Y'_k), Y'_k$ is a closed neighborhood of Y_i , and $\pi_{Y_k} \psi'_k = \psi_k, k = 0, \dots, n$. Let X_k, B_k, η_k and γ_k be as in Definition 2.8. B_0 and η_0 satisfy the conclusion of the proposition, except that we need to prove that B_0 is locally isomorphic to $C(X) \otimes D$. We prove by induction on decreasing k that the $C(X_k)$ -algebras B_k are locally trivial. Indeed $B_n = C(X_n) \otimes D$ and assuming that B_k is locally trivial, it follows by Lemma 6.6 that B_{k-1} is locally trivial, since by (5)

$$B_{k-1} \cong B_k \oplus_{\pi_{\eta_k, \pi_{\psi_{k-1}}}} E_{k-1} \cong B_k \oplus_{\pi, \gamma_k \pi} E_{k-1} \quad (\pi = \pi_{X_k \cap Y_{k-1}})$$

and $\gamma_k : E_{k-1}(X_k \cap Y_{k-1}) \rightarrow B_k(X_k \cap Y_{k-1}), (\gamma_k)_x = (\eta_k)_x^{-1}(\psi_{k-1})_x$, extends to a $*$ -monomorphism $\alpha : E_{k-1}(X_k \cap Y'_{k-1}) \rightarrow B_k(X_k \cap Y'_{k-1}), \alpha_x = (\eta_k)_x^{-1}(\psi'_{k-1})_x$ and $KK(\alpha_x)$ is a KK -equivalence since both $KK((\eta_k)_x)$ and $KK((\psi_{k-1})_x)$ are KK -equivalences. \square

7. When is a $C(X)$ -algebra locally trivial

In this section we prove Theorems 1.1–1.5 and some of their consequences.

7.1. Proof of Theorem 1.2

Proof. Let X denote the primitive spectrum of A . Then A is a continuous $C(X)$ -algebra and its fibers are stable Kirchberg algebras (see [5, 2.2.2]). Since A is separable, X is metrizable by Lemma 2.2. By Proposition 6.7 there is a sequence of $C(X)$ -algebras $(A_k)_{k=1}^\infty$ locally isomorphic to $C(X) \otimes D$ and a sequence of $C(X)$ -linear $*$ -monomorphisms $(\eta_k : A_k \rightarrow A)_{k=1}^\infty$, such that $KK(\eta_k)_x$ is a KK -equivalence for each $x \in X$ and $(\eta_k(A_k))_{k=1}^\infty$ is an exhaustive sequence of $C(X)$ -subalgebras of A . Since D is weakly semiprojective and KK -stable, after passing to a subsequence of (A_k) if necessary, we find a sequence $(\sigma_k)_{k=1}^\infty, \sigma_k \in KK(D, A_k)$ such that $KK(\eta_k)\sigma_k = \sigma$ for all $k \geq 1$. Since both $KK(\eta_k)_x$ and σ_x are KK -equivalences, we deduce that $(\sigma_k)_x \in KK(D, A_k(x))^{-1}$ for all $x \in X$. By Proposition 6.4, for each $k \geq 1$ there is an isomorphism of $C(X)$ -algebras $\varphi_k : C(X) \otimes D \rightarrow A_k$ such that $KK(\varphi_k) = \sigma_k$. Therefore if we set $\theta_k = \eta_k \varphi_k$, then θ_k is a $C(X)$ -linear $*$ -monomorphism from $B \stackrel{def}{=} C(X) \otimes D$ to A such that $KK(\theta_k) = \sigma$ and $(\theta_k(B))_{k=1}^\infty$ is an exhaustive sequence of $C(X)$ -subalgebras of A . Using again the weak semiprojectivity and the KK -stability of D , and Lemma 4.1, after passing to a subsequence of $(\theta_k)_{k=1}^\infty$, we construct a sequence of finite sets $\mathcal{F}_k \subset B$ and a sequence of $C(X)$ -linear $*$ -monomorphisms $\mu_k : B \rightarrow B$ such that

- (i) $KK(\theta_{k+1}\mu_k) = KK(\theta_k)$ for all $k \geq 1$,
- (ii) $\|\theta_{k+1}\mu_k(a) - \theta_k(a)\| < 2^{-k}$ for all $a \in \mathcal{F}_k$ and all $k \geq 1$,
- (iii) $\mu_k(\mathcal{F}_k) \subset \mathcal{F}_{k+1}$ for all $k \geq 1$,
- (iv) $\bigcup_{j=k+1}^\infty (\mu_{j-1} \circ \dots \circ \mu_k)^{-1}(\mathcal{F}_j)$ is dense in B and $\bigcup_{j=k}^\infty \theta_j(\mathcal{F}_j)$ is dense in A for all $k \geq 1$.

Arguing as in the proof of [29, Prop. 2.3.2], one verifies that

$$\Delta_k(a) = \lim_{j \rightarrow \infty} \theta_j \circ (\mu_{j-1} \circ \dots \circ \mu_k)(a)$$

defines a sequence of $*$ -monomorphisms $\Delta_k : B \rightarrow A$ such that $\Delta_{k+1}\mu_k = \Delta_k$ and the induced map $\Delta : \varinjlim_k (B, \mu_k) \rightarrow A$ is an isomorphism of $C(X)$ -algebras. Let us show that $\varinjlim_k (B, \mu_k)$ is isomorphic to B . To this purpose, in view of Elliott’s intertwining argument, it suffices to show that each map μ_k is approximately unitarily equivalent to a $C(X)$ -linear automorphism of B . Since $KK(\theta_k) = \sigma$, we deduce from (i) that $KK((\mu_k)_x) = KK(\text{id}_D)$ for all $x \in X$. By Proposition 6.1, this property implies that each map μ_k is approximately unitarily equivalent to a $C(X)$ -linear automorphism of B . Therefore there is an isomorphism of $C(X)$ -algebras $\Delta : B \rightarrow A$. Let us show that we can arrange that $KK(\Delta|_D) = \sigma$. By Theorem 3.1, there is a full $*$ -homomorphism $\alpha : D \rightarrow B$ such that $KK(\alpha) = KK(\Delta^{-1})\sigma$. Since $KK(\Delta_x^{-1})\sigma_x \in KK(D, D)^{-1}$, by Proposition 6.1 there is $\Phi_1 : D \rightarrow C(X) \otimes D$ such that $\tilde{\Phi}_1 \in \text{Aut}_{C(X)}(B)$ and $KK(\Phi_1) = KK(\Delta^{-1})\sigma$. Then $\Phi = \Delta\tilde{\Phi}_1 : B \rightarrow A$ is an isomorphism such that $KK(\Phi|_D) = KK(\Delta\Phi_1) = \sigma$. \square

Dixmier and Douady [12] proved that a continuous field with fibers \mathcal{K} over a finite dimensional locally compact Hausdorff space is locally trivial if and only if it verifies Fell’s condition, i.e. for each $x_0 \in X$ there is a continuous section a of the field such that $a(x)$ is a rank one projection for each x in a neighborhood of x_0 . We have an analogous result:

Corollary 7.1. *Let A be a separable C^* -algebra whose primitive spectrum X is Hausdorff and of finite dimension. Suppose that for each $x \in X$, $A(x)$ is KK -semiprojective, nuclear, purely infinite and stable. Then A is locally trivial if and only if for each $x \in X$ there exist a closed neighborhood V of x , a Kirchberg algebra D and $\sigma \in KK(D, A(V))$ such that $\sigma_v \in KK(D, A(v))^{-1}$ for each $v \in V$.*

Proof. One applies Theorem 1.2 for $D \otimes \mathcal{K}$ and $A(V)$. \square

Proposition 7.2. *Let ψ be a full endomorphism of a Kirchberg algebra D . If D is unital we assume that $\psi(1) = 1$ as well. Then the continuous $C[0, 1]$ -algebra $E = \{f \in C[0, 1] \otimes D : f(0) \in \psi(D)\}$ is locally trivial if and only if ψ is homotopic to an automorphism of D .*

Proof. Suppose that E is trivial on some neighborhood of 0. Thus there is $s \in (0, 1]$ and an isomorphism $\theta : C[0, s] \otimes D \xrightarrow{\cong} E[0, s]$. Since $E[0, s] \subset C[0, s] \otimes D$, there is a continuous path $(\theta_t)_{t \in [0, s]}$ in $\text{End}(D)$ such that $\theta_t \in \text{Aut}(D)$ for $0 < t \leq s$ and $\theta_0(D) = \psi(D)$. Set $\beta = \theta_0^{-1}\psi \in \text{Aut}(D)$. Then ψ is homotopic to an automorphism via the path $(\theta_t\beta)_{t \in [0, s]}$. Conversely, if ψ is homotopic to an automorphism α , then by Theorem 3.1 there is a continuous path $(u_t)_{t \in (0, 1]}$ of unitaries in D^+ such that $\lim_{t \rightarrow 0} \|\psi(d) - u_t\alpha(d)u_t^*\| = 0$ for all $d \in D$. The path $(\theta_t)_{t \in [0, 1]}$ defined by $\theta_0 = \psi$ and $\theta_t = u_t\alpha u_t^*$ for $t \in (0, 1]$ induces a $C[0, 1]$ -linear $*$ -endomorphism of $C[0, 1] \otimes D$ which maps injectively $C[0, 1] \otimes D$ onto E . \square

7.2. Proof of Theorem 1.3

Proof. For the first part we apply Theorem 1.2 for $D = \mathcal{O}_2 \otimes \mathcal{K}$ and $\sigma = 0$. For the second part we assert that if D is a Kirchberg such that all continuous $C[0, 1]$ -algebras with fibers isomorphic

to D are locally trivial then D is stable and $KK(D, D) = 0$. Thus D is KK -equivalent to \mathcal{O}_2 and hence that $D \cong \mathcal{O}_2 \otimes \mathcal{K}$ by [29, Thm. 8.4.1]. The Kirchberg algebra D is either unital or stable [29, Prop. 4.1.3]. Let $\psi : D \rightarrow D$ be a $*$ -monomorphism such that $KK(\psi) = 0$ and such that $\psi(1_D) < 1_D$ if D is unital. By Proposition 7.2 ψ is homotopic to an automorphism of θ of D . Therefore D must be nonunital (and hence stable), since otherwise 1_D would be homotopic to its proper subprojection $\psi(1_D)$. Moreover $KK(\theta) = KK(\psi) = 0$ and hence $KK(D, D) = 0$ since θ is an automorphism. \square

We turn now to unital $C(X)$ -algebras.

Theorem 7.3. *Let A be a separable unital $C(X)$ -algebra over a finite dimensional compact Hausdorff space X . Suppose that each fiber $A(x)$ is nuclear simple and purely infinite. Then A is isomorphic to $C(X) \otimes D$, for some KK -semiprojective unital Kirchberg algebra D , if and only if there is $\sigma \in KK(D, A)$ such that $K_0(\sigma)[1_D] = [1_A]$ and $\sigma_x \in KK(D, A(x))^{-1}$ for all $x \in X$. For any such σ there is an isomorphism of $C(X)$ -algebras $\Phi : C(X) \otimes D \rightarrow A$ such that $KK(\Phi|_D) = \sigma$.*

Proof. We verify the nontrivial implication. X is metrizable by Lemma 2.2. A is a continuous $C(X)$ -algebra by Lemma 2.3. By Theorem 1.2, there is an isomorphism $\Phi : C(X) \otimes D \otimes \mathcal{K} \rightarrow A \otimes \mathcal{K}$ such that $KK(\Phi) = \sigma$. Since $K_0(\sigma)[1_D] = [1_A]$, and since $A \otimes \mathcal{K}$ contains a full properly infinite projection, we may arrange that $\Phi(1_{C(X) \otimes D} \otimes e_{11}) = 1_A \otimes e_{11}$ after conjugating Φ by some unitary $u \in M(A \otimes \mathcal{K})$. Then $\varphi = \Phi|_{C(X) \otimes D \otimes e_{11}}$ satisfies the conclusion of the theorem. \square

7.3. Proof of Theorem 1.4

Proof. Let D be a KK -semiprojective unital Kirchberg algebra D such that every unital $*$ -endomorphism of D is a KK -equivalence. Suppose that A is a separable unital $C(X)$ -algebra over a finite dimensional compact Hausdorff space the fibers of which are isomorphic to D . We shall prove that A is locally trivial. By Theorem 7.3, it suffices to show that each point $x_0 \in X$ has a closed neighborhood V for which there is $\sigma \in KK(D, A(V))$ such that $K_0(\sigma)[1_D] = [1_{A(V)}]$ and $\sigma_x \in KK(D, A(x))^{-1}$ for all $x \in V$.

Let $(V_n)_{n=1}^\infty$ be a decreasing sequence of closed neighborhoods of x_0 whose intersection is $\{x_0\}$. Then $A(x_0) \cong \varprojlim A(V_n)$. By assumption, there is an isomorphism $\eta : D \rightarrow A(x_0)$. Since D is KK -semiprojective, there is $m \geq 1$ such that $KK(\eta)$ lifts to some $\sigma \in KK(D, A(V_m))$ such that $K_0(\sigma)[1_D] = [1_{A(V_m)}]$. Let $x \in V_m$. By assumption, there is an isomorphism $\varphi : A(x) \rightarrow D$. The K_0 -morphism induced by $KK(\varphi)\sigma_x$ maps $[1_D]$ to itself. By Theorem 3.1 there is a unital $*$ -homomorphism $\psi : D \rightarrow D$ such that $KK(\psi) = KK(\varphi)\sigma_x$. By assumption we must have $KK(\psi) \in KK(D, D)^{-1}$ and hence $\sigma_x \in KK(D, A(x))^{-1}$ since φ is an isomorphism. Therefore $A(V_m) \cong C(V_m) \otimes D$ by Theorem 7.3.

Conversely, let us assume that all separable unital continuous $C[0, 1]$ -algebras with fibers isomorphic to D are locally trivial. Let ψ be any unital $*$ -endomorphism of D . By Proposition 7.2 ψ is homotopic to an automorphism of D and hence $KK(\psi)$ is invertible. \square

7.4. Proof of Theorem 1.1

Proof. Let A be as in Theorem 1.1 and let $n \in \{2, 3, \dots\} \cup \{\infty\}$. It is known that \mathcal{O}_n satisfies the UCT. Moreover $K_0(\mathcal{O}_n)$ is generated by $[1_{\mathcal{O}_n}]$ and $K_1(\mathcal{O}_n) = 0$. Therefore any unital $*$ -

endomorphism of \mathcal{O}_n is a KK -equivalence. It follows that A is locally trivial by Theorem 1.4. Suppose now that $n = 2$. Since $KK(\mathcal{O}_2, \mathcal{O}_2) = KK(\mathcal{O}_2, A) = 0$, we may apply Theorem 1.4 with $\sigma = 0$ and obtain that $A \cong C(X) \otimes \mathcal{O}_2$. Suppose now that $n = \infty$. Let us define $\theta : K_0(\mathcal{O}_\infty) \rightarrow K_0(A)$ by $\theta(k[1_{\mathcal{O}_\infty}]) = k[1_A]$, $k \in \mathbb{Z}$. Since \mathcal{O}_∞ satisfies the UCT, θ lifts to some element $\sigma \in KK(\mathcal{O}_\infty, A)$. By Theorem 1.4 it follows that $A \cong C(X) \otimes \mathcal{O}_\infty$. Finally let us consider the case $n \in \{3, 4, \dots\}$. Then $K_0(\mathcal{O}_n) = \mathbb{Z}/(n - 1)$. Since \mathcal{O}_n satisfies the UCT, the existence of an element $\sigma \in KK(\mathcal{O}_n, A)$ such that $K_0(\sigma)[1_{\mathcal{O}_n}] = [1_A]$ is equivalent to the existence of a morphism of groups $\theta : \mathbb{Z}/(n - 1) \rightarrow K_0(A)$ such that $\theta(\bar{1}) = [1_A]$. This is equivalent to requiring that $(n - 1)[1_A] = 0$. \square

As a corollary of Theorem 1.1 we have that $[X, \text{Aut}(\mathcal{O}_\infty)]$ reduces to a point. The homotopy groups of the endomorphisms of the stable Cuntz–Krieger algebras were computed in [7]. Let v_1, \dots, v_n be the canonical generators of \mathcal{O}_n , $2 \leq n < \infty$.

Theorem 7.4. *For any compact metrizable space X there is a bijection $[X, \text{Aut}(\mathcal{O}_n)] \rightarrow K_1(C(X) \otimes \mathcal{O}_n)$. The k th-homotopy group $\pi_k(\text{Aut}(\mathcal{O}_n))$ is isomorphic to $\mathbb{Z}/(n - 1)$ if k is odd and it vanishes if k is even. In particular $\pi_1(\text{Aut}(\mathcal{O}_n))$ is generated by the class of the canonical action of \mathbb{T} on \mathcal{O}_n , $\lambda_z(v_i) = zv_i$.*

Proof. Since \mathcal{O}_n satisfies the UCT, we deduce that $\text{End}(\mathcal{O}_n)^* = \text{End}(\mathcal{O}_n)$. An immediate application of Proposition 6.1 shows that the natural map $\text{Aut}(\mathcal{O}_n) \hookrightarrow \text{End}(\mathcal{O}_n)$ induces an isomorphism of groups $[X, \text{Aut}(\mathcal{O}_n)] \cong [X, \text{End}(\mathcal{O}_n)]$. Let $\iota : \mathcal{O}_n \rightarrow C(X) \otimes \mathcal{O}_n$ be defined by $\iota(v_i) = 1_{C(X)} \otimes v_i$, $i = 1, \dots, n$. The map $\psi \mapsto u(\psi) = \psi(v_1)\iota(v_1)^* + \dots + \psi(v_n)\iota(v_n)^*$ is known to be a homeomorphism from $\text{Hom}(\mathcal{O}_n, C(X) \otimes \mathcal{O}_n)$ to the unitary group of $C(X) \otimes \mathcal{O}_n$. Its inverse maps a unitary w to the $*$ -homomorphism ψ uniquely defined by $\psi(v_i) = w\iota(v_i)$, $i = 1, \dots, n$. Therefore

$$[X, \text{Aut}(\mathcal{O}_n)] \cong [X, \text{End}(\mathcal{O}_n)] \cong \pi_0(U(C(X) \otimes \mathcal{O}_n)) \cong K_1(C(X) \otimes \mathcal{O}_n).$$

The last isomorphism holds since $\pi_0(U(B)) \cong K_1(B)$ if $B \cong B \otimes \mathcal{O}_\infty$, by [28, Lemma 2.1.7]. One verifies easily that if $\varphi \in \text{Hom}(\mathcal{O}_n, C(X) \otimes \mathcal{O}_n)$, then $u(\tilde{\psi}\varphi) = \tilde{\psi}(u(\varphi))u(\psi)$. Therefore the bijection $\chi : [X, \text{End}(\mathcal{O}_n)] \rightarrow K_1(C(X) \otimes \mathcal{O}_n)$ is an isomorphism of groups whenever $K_1(\tilde{\psi}) = \text{id}$ for all $\psi \in \text{Hom}(\mathcal{O}_n, C(X) \otimes \mathcal{O}_n)$. Using the $C(X)$ -linearity of $\tilde{\psi}$ one observes that this holds if the $n - 1$ torsion of $K_0(C(X))$ reduces to $\{0\}$, since in that case the map $K_1(C(X)) \rightarrow K_1(C(X) \otimes \mathcal{O}_n)$ is surjective by the Künneth formula. \square

Corollary 7.5. *Let X be a finite dimensional compact metrizable space. The isomorphism classes of unital separable $C(SX)$ -algebras with all fibers isomorphic to \mathcal{O}_n are parameterized by $K_1(C(X) \otimes \mathcal{O}_n)$.*

Proof. This follows from Theorems 1.1 and 7.4, since the locally trivial principal H -bundles over $SX = X \times [0, 1]/X \times \{0, 1\}$ are parameterized by the homotopy classes $[X, H]$ if H is a path connected group [17, Cor. 8.4]. Here we take $H = \text{Aut}(\mathcal{O}_n)$. \square

Examples of nontrivial unital $C(X)$ -algebras with fiber \mathcal{O}_n over a $2m$ -sphere arising from vector bundles were exhibited in [36], see also [35].

We need some preparation for the proof of Theorem 1.5. Let G be a group, let $g \in G$ and set $\text{End}(G, g) = \{\alpha \in \text{End}(G) : \alpha(g) = g\}$. The pair (G, g) is called *weakly rigid* if $\text{End}(G, g) \subset \text{Aut}(G)$ and *rigid* if $\text{End}(G, g) = \{\text{id}_G\}$.

Theorem 7.6. *If G is a finitely generated abelian group, then (G, g) is weakly rigid if and only if (G, g) is isomorphic to one of the pointed groups from the list \mathcal{G} of Theorem 1.5.*

Proof. First we make a number of remarks.

(1) (G, g) is weakly rigid if and only if $(G, \alpha(g))$ is weakly rigid for some (or any) $\alpha \in \text{Aut}(G)$. Indeed if $\beta \in \text{End}(G, g)$ then $\alpha\beta\alpha^{-1} \in \text{End}(G, \alpha(g))$.

(2) By considering the zero endomorphism of G we see that if (G, g) is weakly rigid and $G \neq 0$ then $g \neq 0$.

(3) If $(G \oplus H, g \oplus h)$ is weakly rigid, then so are (G, g) and (H, h) .

(4) Let us observe that (\mathbb{Z}^2, g) is not weakly rigid for any g . Indeed, if $g = (a, b) \neq 0$, then the matrix $\begin{pmatrix} 1+b^2 & -ab \\ -ab & 1+a^2 \end{pmatrix}$ defines an endomorphism α of \mathbb{Z}^2 such that $\alpha(g) = g$, but α is not invertible since $\det(\alpha) = 1 + a^2 + b^2 > 1$.

(5) Let p be a prime and let $1 \leq e_1 \leq e_2, 0 \leq s_1 < e_1, 0 \leq s_2 < e_2$ be integers. If $(G, g) = (\mathbb{Z}/p^{e_1} \oplus \mathbb{Z}/p^{e_2}, p^{s_1} \oplus p^{s_2})$ is weakly rigid then $0 < s_2 - s_1 < e_2 - e_1$. Indeed if $s_1 \geq s_2$ then the matrix $\begin{pmatrix} 0 & p^{s_1-s_2} \\ 0 & 1 \end{pmatrix}$ induces a noninjective endomorphism of (G, g) . Also if $s_1 < s_2$ and $s_2 - s_1 \geq e_2 - e_1$ then $p^{e_1}\bar{b} = 0$ in \mathbb{Z}/p^{e_2} , where $b = p^{s_2-s_1}$ and so the matrix $\begin{pmatrix} 1 & 0 \\ b & 0 \end{pmatrix}$ induces a well-defined noninjective endomorphism of (G, g) .

(6) Let p be a prime and let $1 \leq k, 0 \leq s < e$ be integers. Suppose that $(\mathbb{Z} \oplus \mathbb{Z}/p^e, k \oplus p^s)$ is weakly rigid. Then k is divisible by p^{s+1} . Indeed, seeking a contradiction suppose that k can be written as $k = p^t c$ where $0 \leq t \leq s$ and c are integers such that c is not divisible by p . Let d be an integer such that $dc - 1$ is divisible by p^e . Then the matrix $\begin{pmatrix} 1 & 0 \\ dp^{s-t} & 0 \end{pmatrix}$ induces a noninjective endomorphism of $(\mathbb{Z} \oplus \mathbb{Z}/p^e, k \oplus p^s)$.

Suppose now that (G, g) is weakly rigid. We shall show that (G, g) is isomorphic to one of the pointed groups from the list \mathcal{G} . Since G is abelian and finitely generated it decomposes as a direct sum of its primary components

$$G \cong \mathbb{Z}^r \oplus G(p_1) \oplus \dots \oplus G(p_m) \tag{20}$$

where p_i are distinct prime numbers. Each primary component $G(p_i)$ is of the form

$$G(p_i) = \mathbb{Z}/p_i^{e_{i1}} \oplus \dots \oplus \mathbb{Z}/p_i^{e_{in(i)}} \tag{21}$$

where $1 \leq e_{i1} \leq \dots \leq e_{in(i)}$ are positive integers. Corresponding to the decomposition (20) we write the base point $g = g_0 \oplus g_1 \oplus \dots \oplus g_m$ with $g_0 \in \mathbb{Z}^r$ and $g_i \in G(p_i)$ for $i \geq 1$. If g_{ij} is the component of g_i in $\mathbb{Z}/p^{e_{ij}}$, then it follows from (1), (2) and (3) that we may assume that $g_{ij} = p^{s_{ij}}$ for some integer $0 \leq s_{ij} < e_{ij}$. Using (3) and (4) we deduce that $r = 1$ in (20) and that $g_0 = k \neq 0$ by (2). We may assume that $k \geq 1$ by (1). Then using (3) and (5) we deduce that for each $1 \leq i \leq m, 0 < s_{ij+1} - s_{ij} < e_{ij+1} - e_{ij}$ for $1 \leq j < n(i)$. Finally, from (3) and (6) we see that k is divisible by the product $p_1^{s_{1n(1)}} \dots p_m^{s_{mn(m)}}$. Therefore (G, g) is isomorphic to one of the pointed groups on the list \mathcal{G} .

Conversely, we shall prove that if (G, g) belongs to the list \mathcal{G} then (G, g) is weakly rigid. This is obvious if G is torsion free i.e. for $(\{0\}, 0)$ and (\mathbb{Z}, k) with $k \geq 1$.

Let us consider the case when G is a torsion group. Since

$$\text{End}(G(p_1) \oplus \cdots \oplus G(p_m), g_1 \oplus \cdots \oplus g_m) \cong \bigoplus_{i=1}^m \text{End}(G(p_i), g_i)$$

it suffices to assume that G is a p -group,

$$(G, g) = (\mathbb{Z}/p^{e_1} \oplus \cdots \oplus \mathbb{Z}/p^{e_n}, p^{s_1} \oplus \cdots \oplus p^{s_n})$$

with $0 \leq s_i < e_i$ for $i = 1, \dots, n$ and $0 < s_{i+1} - s_i < e_{i+1} - e_i$ for $1 \leq i < n$. For each $0 \leq i, j \leq n$ set $e_{ij} = \max\{e_i - e_j, 0\}$. It follows immediately that $s_i < e_{ij} + s_j$ for all $i \neq j$. Let $\alpha \in \text{End}(G, g)$. It is well known that α is induced by a square matrix $A = [a_{ij}] \in M_n(\mathbb{Z})$ with the property that each entry a_{ij} is divisible by $p^{e_{ij}}$ and so $a_{ij} = p^{e_{ij}} b_{ij}$ for some $b_{ij} \in \mathbb{Z}$, see [16]. Since $\alpha(g) = g$, we have $\sum_{j=1}^n \bar{b}_{ij} p^{e_{ij} + s_j} = p^{s_i}$ in \mathbb{Z}/p^{e_i} for all $0 \leq i \leq n$. Since $e_{ij} + s_j > s_i$ for $i \neq j$ and $e_i > s_i$ we see that $b_{ii} - 1$ must be divisible by p for all $1 \leq i \leq n$. Since $\det(A)$ is congruent to $b_{11} \cdots b_{nn}$ modulo p it follows that $\det(A)$ is not divisible by p and so $\alpha \in \text{Aut}(G)$ by [16].

Finally consider the case when $(G, g) = (\mathbb{Z} \oplus G(p_1) \oplus \cdots \oplus G(p_m), k \oplus g_1 \oplus \cdots \oplus g_m)$. If $\gamma \in \text{End}(G, g)$ then there exist $\alpha_i \in \text{End}(G(p_i), g_i)$ and $d_i \in G(p_i)$, $1 \leq i \leq n$, such that $\gamma(x_0 \oplus x_1 \oplus \cdots \oplus x_m) = x_0 \oplus (\alpha_1(x_1) + x_0 d_1) \oplus \cdots \oplus (\alpha_m(x_m) + x_0 d_m)$. Note that if each α_i is an automorphism then so is γ . Indeed, its inverse is $\gamma^{-1}(x_0 \oplus x_1 \oplus \cdots \oplus x_m) = x_0 \oplus (\alpha_1^{-1}(x_1) + x_0 c_1) \oplus \cdots \oplus (\alpha_m(x_m)^{-1} + x_0 c_m)$, where $c_i = -\alpha_i^{-1}(d_i)$. Therefore it suffices to consider the case $m = 1$, i.e.

$$(G, g) = (\mathbb{Z} \oplus \mathbb{Z}/p^{e_1} \oplus \cdots \oplus \mathbb{Z}/p^{e_n}, k \oplus p^{s_1} \oplus \cdots \oplus p^{s_n}),$$

and (G, g) is on the list \mathcal{G} (e). In particular $k = p^{s_n+1} \ell$ for some $\ell \in \mathbb{Z}$. Let $\gamma \in \text{End}(G, g)$. Then there exists $\alpha \in \text{End}(G(p))$ and $d \in G(p)$ such that $\gamma(x_0 \oplus x) = x_0 \oplus (\alpha(x) + x_0 d)$. Just as above, α is induced by a square matrix $A \in M_n(\mathbb{Z})$ of the form $A = [b_{ij} p^{e_{ij}}] \in M_n(\mathbb{Z})$ with $b_{ij} \in \mathbb{Z}$, $e_{ij} = \max\{e_i - e_j, 0\}$. Since $\gamma(g) = g$ we have that $p^{s_n+1} \ell d_i + \sum_{j=1}^n \bar{b}_{ij} p^{e_{ij} + s_j} = p^{s_i}$ in \mathbb{Z}/p^{e_i} for all $0 \leq i \leq n$, where the d_i are the components of d . By reasoning as in the case when G was a torsion group considered above, since $s_n + 1 > s_i$ for all $1 \leq i \leq n$, $e_{ij} + s_j > s_i$ for all $i \neq j$ and $e_i > s_i$, it follows again that each $b_{ii} - 1$ is divisible by p and that the endomorphism α of $G(p)$ induced by the matrix A is an automorphism. We conclude that γ is an automorphism. \square

7.5. Proof of Theorem 1.5

Proof. (ii) and (iii) Let D be a unital Kirchberg algebra such that D satisfies the UCT and $K_*(D)$ is finitely generated. Then D is KK -semiprojective by Proposition 3.14 and $KK(D, D)^{-1} = \{\alpha \in KK(D, D): K_*(\alpha) \text{ is bijective}\}$. In conjunction with Theorem 3.1, this shows that all unital $*$ -endomorphisms of D are KK -equivalences if and only if both $(K_0(D), [1_D])$ and $(K_1(D), 0)$ are weakly rigid. Equivalently, $K_1(D) = 0$ and $(K_0(D), [1_D])$ is weakly rigid. By Theorem 7.6 $(K_0(D), [1_D])$ is weakly rigid if and only if it is isomorphic to one pointed groups from the list \mathcal{G} of Theorem 1.5. We conclude the proof of (ii) and (iii) by applying Theorem 1.4.

(i) By Theorem 1.1 both \mathcal{O}_2 and \mathcal{O}_∞ have the automatic triviality property. Conversely, suppose that D has the automatic triviality property, where D is a unital Kirchberg algebra satisfying

the UCT and such that $K_*(D)$ is finitely generated. We shall prove that D is isomorphic to either \mathcal{O}_2 or \mathcal{O}_∞ .

Let Y be a finite connected CW-complex and let $\iota : D \rightarrow C(Y) \otimes D$ be the map $\iota(d) = 1 \otimes d$. Let $[D, C(Y) \otimes D]$ denote the homotopy classes of unital $*$ -homomorphisms from D to $C(Y) \otimes D$. By Theorem 3.1 the image of the map $\Delta : [D, C(Y) \otimes D] \rightarrow KK(D, C(Y) \otimes D)$ defined by $[\varphi] \mapsto KK(\varphi) - KK(\iota)$ coincides with the kernel of the restriction morphism $\rho : KK(D, C(Y) \otimes D) \rightarrow KK(\mathbb{C}1_D, C(Y) \otimes D)$.

We claim that $\ker \rho$ must vanish for all Y . Let $h \in \ker \rho$. Then there is a unital $*$ -homomorphism $\varphi : D \rightarrow C(Y) \otimes D$ such that $\Delta[\varphi] = h$. By Theorem 1.4, each unital endomorphism of D induces a KK -equivalence. Therefore, by Proposition 6.1 there is a $*$ -homomorphism $\Phi : D \rightarrow C(Y) \otimes D$ such that $\Phi_y \in \text{Aut}(D)$ for all $y \in Y$ and $KK(\Phi) = KK(\varphi)$. Therefore $\Delta[\Phi] = KK(\Phi) - KK(\iota) = h$. By hypothesis, the $\text{Aut}(D)$ -principal bundle constructed over the suspension of Y with characteristic map $y \mapsto \Phi_y$ is trivial. It follows then from [17, Thm. 8.2, p. 85] that this map is homotopic to the constant map $Y \rightarrow \text{Aut}(D)$ which shrinks Y to id_D . This implies that Φ is homotopic to ι and hence $h = 0$.

Let us now observe that $\ker \rho$ contains subgroups isomorphic to $\text{Hom}(K_1(D), K_1(D))$ and $\text{Ext}(K_0(D), K_0(D))$ if $Y = \mathbb{T}$, since D satisfies the UCT. It follows that both these groups must vanish and so $K_1(D) = 0$ and $K_0(D)$ is torsion free. On the other hand, $(K_0(D), [1_D])$ is weakly rigid by the first part of the proof. Since $K_0(D)$ is torsion free we deduce from Theorem 7.6 that either $K_0(D) = 0$ in which case $D \cong \mathcal{O}_2$ or that $(K_0(D), [1_D]) \cong (\mathbb{Z}, k)$, $k \geq 1$, in which case $D \cong M_k(\mathcal{O}_\infty)$ by the classification theorem of Kirchberg and Phillips.

To conclude the proof, it suffices to show that $\ker \rho \neq 0$ if $D = M_k(\mathcal{O}_\infty)$, $k \geq 2$ and Y is the two dimensional space obtained by attaching a disk to a circle by a degree- k map. Since $K_0(C(Y) \otimes \mathcal{O}_\infty) \cong \mathbb{Z} \oplus \mathbb{Z}/k$ we can identify the map ρ with the map $\mathbb{Z} \oplus \mathbb{Z}/k \rightarrow \mathbb{Z} \oplus \mathbb{Z}/k$, $x \mapsto kx$ and so $\ker \rho \cong \mathbb{Z}/k \neq 0$ if $k \geq 2$. \square

7.6. Added in proof

Some of the results from this paper are further developed in [9]. Theorem 1.2 was shown to hold for all stable Kirchberg algebras D . The assumption that X is finite dimensional is essential in Theorem 1.1. Theorem 1.5(ii) extends as follows: \mathcal{O}_2 , \mathcal{O}_∞ and $B \otimes \mathcal{O}_\infty$, where B is a unital UHF algebra of infinite type, are the only unital Kirchberg algebras which satisfy the UCT and have the automatic triviality property in the unital sense.

References

- [1] B. Blackadar, Shape theory for C^* -algebras, *Math. Scand.* 56 (2) (1985) 249–275.
- [2] B. Blackadar, *K-Theory for Operator Algebras*, 2nd ed., *Math. Sci. Res. Inst. Publ.*, vol. 5, Cambridge University Press, Cambridge, 1998.
- [3] É. Blanchard, Déformations de C^* -algèbres de Hopf, *Bull. Soc. Math. France* 124 (1) (1996) 141–215.
- [4] É. Blanchard, E. Kirchberg, Global Glimm halving for C^* -bundles, *J. Operator Theory* 52 (2004) 385–420.
- [5] É. Blanchard, E. Kirchberg, Non-simple purely infinite C^* -algebras: The Hausdorff case, *J. Funct. Anal.* 207 (2004) 461–513.
- [6] L.G. Brown, Stable isomorphism of hereditary subalgebras of C^* -algebras, *Pacific J. Math.* 71 (1977) 335–348.
- [7] J. Cuntz, On the homotopy groups of the space of endomorphisms of a C^* -algebra (with applications to topological Markov chains), in: *Operator Algebras and Group Representations*, vol. I, Neptun, 1980, in: *Monogr. Stud. Math.*, vol. 17, Pitman, Boston, MA, 1984, pp. 124–137.
- [8] J. Cuntz, W. Krieger, A class of C^* -algebras and topological Markov chains, *Invent. Math.* 56 (1980) 251–268.
- [9] M. Dadarlat, Fiberwise KK -equivalence of continuous fields of C^* -algebras, *K-Theory* 3 (2) (2009) 205–219.

- [10] M. Dadarlat, G.A. Elliott, One-parameter continuous fields of Kirchberg algebras, *Comm. Math. Phys.* 274 (3) (2007) 795–819.
- [11] M. Dadarlat, C. Pasnicu, Continuous fields of Kirchberg C^* -algebras, *J. Funct. Anal.* 226 (2005) 429–451.
- [12] J. Dixmier, C^* -algebras, North-Holland, Amsterdam, 1982.
- [13] S. Eilenberg, N. Steenrod, *Foundations of Algebraic Topology*, Princeton University Press, Princeton, NJ, 1952.
- [14] S. Eilers, T.A. Loring, Computing contingencies for stable relations, *Internat. J. Math.* 10 (3) (1999) 301–326.
- [15] J.M.G. Fell, The structure of algebras of operator fields, *Acta Math.* 106 (1961) 233–280.
- [16] C.J. Hillar, D.L. Rhea, Automorphisms of finite abelian groups, *Amer. Math. Monthly* 114 (10) (2007) 917–923.
- [17] D. Husemoller, *Fibre Bundles*, 3rd ed., Grad. Texts in Math., vol. 20, Springer-Verlag, New York, 1994, 1st ed., 1966.
- [18] G.G. Kasparov, Equivariant KK -theory and the Novikov conjecture, *Invent. Math.* 91 (1) (1988) 147–201.
- [19] E. Kirchberg, The classification of purely infinite C^* -algebras using Kasparov’s theory, preprint, 1994.
- [20] E. Kirchberg, Das nicht-kommutative Michael-auswahlprinzip und die klassifikation nicht-einfacher algebren, in: C^* -Algebras, Münster, 1999, Springer-Verlag, Berlin, 2000, pp. 92–141.
- [21] E. Kirchberg, M. Rørdam, Non-simple purely infinite C^* -algebras, *Amer. J. Math.* 122 (3) (2000) 637–666.
- [22] E. Kirchberg, M. Rørdam, Infinite non-simple C^* -algebras: Absorbing the Cuntz algebra O_∞ , *Adv. Math.* 167 (2) (2002) 195–264.
- [23] E. Kirchberg, S. Wassermann, Operations on continuous bundles of C^* -algebras, *Math. Ann.* 303 (4) (1995) 677–697.
- [24] H. Lin, Weak semiprojectivity in purely infinite C^* -algebras, *Canad. J. Math.* 59 (2) (2007) 343–371.
- [25] T. Loring, Lifting Solutions to Perturbing Problems in C^* -Algebras, in: *Fields Inst. Monogr.*, vol. 8, Amer. Math. Soc., Providence, RI, 1997.
- [26] B. Neubüser, Semiprojektivität und Realisierungen von rein unendlichen C^* -Algebren, PhD thesis, Universität Münster, 2000, Preprintreihe des SFB 478 — Geometrische Strukturen in der Mathematik; Heft 118.
- [27] M. Nilsen, C^* -bundles and $C_0(X)$ -algebras, *Indiana Univ. Math. J.* 45 (2) (1996) 463–477.
- [28] N.C. Phillips, A classification theorem for nuclear purely infinite simple C^* -algebras, *Doc. Math.* (5) (2000) 49–114.
- [29] M. Rørdam, Classification of Nuclear, Simple C^* -Algebras, *Encyclopaedia Math. Sci.*, vol. 126, Springer-Verlag, Berlin, 2002.
- [30] M. Rørdam, Stable C^* -algebras, in: *Operator Algebras and Applications*, in: *Adv. Stud. Pure Math.*, vol. 38, Math. Soc. Japan, Tokyo, 2004, pp. 177–199.
- [31] J. Rosenberg, C. Schochet, The Künneth theorem and the universal coefficient theorem for Kasparov’s generalized K -functor, *Duke Math. J.* 55 (2) (1987) 431–474.
- [32] J.S. Spielberg, Weak semiprojectivity for purely infinite C^* -algebras, *Canad. Math. Bull.* 50 (3) (2007) 460–468.
- [33] J.S. Spielberg, Semiprojectivity for certain purely infinite C^* -algebras, *Trans. Amer. Math. Soc.* 361 (6) (2009) 2805–2830.
- [34] W. Szymański, On semiprojectivity of C^* -algebras of directed graphs, *Proc. Amer. Math. Soc.* 130 (5) (2002) 1391–1399.
- [35] E. Vasselli, The C^* -algebra of a vector bundle of fields of Cuntz algebras, *J. Funct. Anal.* 222 (2) (2005) 491–502.
- [36] E. Vasselli, Bundles of C^* -algebras and the $KK(X; -, -)$ -bifunctor, in: C^* -Algebras and Elliptic Theory, in: *Trends Math.*, Birkhäuser, Basel, 2006, pp. 313–327.
- [37] S. Wassermann, private communication.
- [38] G.W. Whitehead, *Elements of Homotopy Theory*, Grad. Texts in Math., vol. 61, Springer-Verlag, New York, 1978.