

Lesson 17

pg 11

Non homogeneous Equations: Method of Undetermined Coefficients (3.5)

Given a nonhomogeneous equation

$y'' + p(t)y' + q(t)y = g(t)$, a function $y = \phi(t)$ is a solution if and only if $L[\phi] = \phi'' + p\phi' + q\phi = g(t)$.

Thm 3.5.1 If $Y_1(t)$ and $Y_2(t)$ are solutions to the nonhomogeneous equation above, then $Y_1(t) - Y_2(t)$ is a solution to the homogeneous equation $y'' + py' + qy = 0$.

Proof. $L[Y_1 - Y_2] = L[Y_1] - L[Y_2]$
 $= g(t) - g(t) = 0$.

This tells us that if $\{y_1, y_2\}$ is a fundamental set for the homogeneous equation, then

$$Y_1 - Y_2 = c_1 y_1 + c_2 y_2 \text{ for some } c_1 \text{ and } c_2$$

Thm 3.5.2 If $y_1(t)$ and $y_2(t)$ form a fundamental set for $y'' + py' + qy = 0$ and if $Y(t)$ is a specific solution to $y'' + py' + qy = g(t)$, then the general solution for $y'' + py' + qy = g(t)$ is $y = c_1 y_1(t) + c_2 y_2(t) + Y(t)$

We often call $y_c(t) = c_1 y_1(t) + c_2 y_2(t)$ the complementary solution and $Y(t)$ a particular solution

Lesson 17

pg. 2

Proof. $L[c_1 y_1 + c_2 y_2 + Y] = c_1 L[y_1] + c_2 L[y_2] + L[Y]$
 $= 0 + 0 + g(t)$

And if ϕ is a solution, then

$\phi - Y = c_1 y_1 + c_2 y_2$, so $\phi = c_1 y_1 + c_2 y_2 + Y$
for some coefficients c_1 and c_2 . \square

Hence, in order to solve a nonhomogeneous equation, we find the complementary solution, find a particular solution, and we add them.

How do we find a particular solution?

One way is using the Method of Undetermined Coefficients.

Given $ay'' + by' + cy = g(t)$, we look at each term of $g(t)$ individually and guess the form of a particular solution based on that.

- If $e^{\alpha t}$ is a term, guess $Ae^{\alpha t}$ is a term of $Y(t)$.
- If $\cos(\alpha t)$ or $\sin(\alpha t)$ is a term, guess $A \cos(\alpha t) + B \sin(\alpha t)$ is a term of $Y(t)$.
- If $a_n t^n + a_{n-1} t^{n-1} + \dots + a_1 t + a_0$ is a term, guess $A_n t^n + A_{n-1} t^{n-1} + \dots + A_1 t + A_0$ is a term of $Y(t)$.
- If a product of the above forms appears as a term, guess the product of the appropriate guesses is a term of $Y(t)$.

Lesson 17

pg. 3

Ex 1. If $g(t)$ is as given below, what is your initial guess for $Y(t)$?

(a) $g(t) = e^{2t} + \sin(3t)$

terms: e^{2t} $\sin(3t)$

guess: Ae^{2t} $B\sin(3t) + C\cos(3t)$

initial guess: $Y(t) = Ae^{2t} + B\sin(3t) + C\cos(3t)$

(b) $g(t) = 4t^3 + t^2\sin(2t)$

terms: $4t^3$ $t^2\sin(2t)$

guess: $A_3t^3 + A_2t^2 + A_1t + A_0$ $(B_2t^2 + B_1t + B_0)\cos(2t)$
 $+ (C_2t^2 + C_1t + C_0)\sin(2t)$

initial guess: $Y(t) = A_3t^3 + A_2t^2 + A_1t + A_0$
 $+ (B_2t^2 + B_1t + B_0)\cos(2t) + (C_2t^2 + C_1t + C_0)\sin(2t)$

(c) $g(t) = te^t + te^{3t}\cos t$

terms: te^t $te^{3t}\cos t$

guess: $(A_1t + A_0)e^t$ $(B_1t + B_0)e^{3t}\cos t + (C_1t + C_0)e^{3t}\sin t$

initial guess: $Y(t) = (A_1t + A_0)e^t + (B_1t + B_0)e^{3t}\cos t + (C_1t + C_0)e^{3t}\sin t$

Lesson 17

pg. 4

This technique is not good enough. What if one of the terms of $g(t)$ is part of the complementary solution? Taking inspiration from repeated roots, multiply that term's guess by t .

Ex 2. Given the complementary solution $y_c(t)$ and $g(t)$, what is your guess for $Y(t)$?

(a) $y_c(t) = c_1 e^t + c_2 e^{2t}$, $g(t) = e^t + \cos(3t)$

terms:	e^t	$\cos(3t)$
guess 1:	Ae^t	$B\cos(3t) + C\sin(3t)$
in y_c ?:	yes!	no!
guess 2:	Ate^t	Same as guess 1
	$Y(t) = Ate^t + B\cos(3t) + C\sin(3t)$	

(b) $y_c(t) = c_1 \cos t + c_2 \sin t$, $g(t) = \sin t + t^2 \cos 3t$

terms:	$\sin t$	$t^2 \cos 3t$
guess 1:	$A\cos t + B\sin t$	$(C_2 t^2 + C_1 t + C_0)\cos 3t + (D_2 t^2 + D_1 t + D_0)\sin 3t$
in y_c ?:	yes	no
guess 2:	$At\cos t + Bt\sin t$	Same as guess 1

$$Y(t) = At\cos t + Bt\sin t + (C_2 t^2 + C_1 t + C_0)\cos 3t + (D_2 t^2 + D_1 t + D_0)\sin 3t$$

Now that we can guess the form of $Y(t)$, we can determine the coefficients by plugging $Y(t)$ into the diff eq.

Method of Undetermined Coefficients

Given $ay'' + by' + cy = g(t)$

1. Find the complementary solution $y_c(t)$ by solving the homogeneous equation $ay'' + by' + cy = 0$.
2. Analyze the terms of $g(t)$ to guess the form of the particular solution $Y(t)$.
3. Plug $Y(t)$ into $ay'' + by' + cy = g(t)$, set up a system of equations, and solve for each coefficient.

Ex 3. Find the general solution to
 $y'' + y' - 2y = 3e^t + te^{-2t}$

$$y_c(t) = c_1 e^t + c_2 e^{-2t}$$

terms:	$3e^t$	te^{-2t}
guess 1:	Ae^t	$(Bt+C)e^{-2t}$
in y_c ?:	yes	yes
guess 2:	Ate^t	$(Bt^2 + Ct)e^{-2t}$

$$Y(t) = Ate^t + (Bt^2 + Ct)e^{-2t}$$

$$Y'(t) = Ate^t + Ae^t + (Bt^2 + Ct)(-2e^{-2t}) + (2Bt + C)e^{-2t}$$

$$= (At + A)e^t + (-2Bt^2 - 2Ct + 2Bt + C)e^{-2t}$$

$$Y''(t) = (At + A)e^t + Ae^t + (-2Bt^2 - 2Ct + 2Bt + C)(-2e^{-2t})$$

$$+ (-4Bt - 2C + 2B)e^{-2t}$$

$$= (At + 2A)e^t + (4Bt^2 + 4Ct - 4Bt - 2C - 4Bt - 2C + 2B)e^{-2t}$$

$$= (At + 2A)e^t + [4Bt^2 + (4C - 8B)t + (-4C + 2B)]e^{-2t}$$

Lesson 17

Pg. 6

Plug in to $y'' + y' - 2y = 3e^t + te^{-2t}$

$$(At + 2A)e^t + [4Bt^2 + (4C - 8B)t + (-4C + 2B)]e^{-2t} \\ + (At + A)e^t + [-2Bt^2 + (-2C + 2B)t + C]e^{-2t} \\ - 2[At e^t + (Bt^2 + Ct)e^{-2t}] = 3e^t + te^{-2t}$$

$$3Ae^t + [-6Bt + (-3C + 2B)]e^{-2t} = 3e^t + te^{-2t}$$

$$\left. \begin{array}{l} 3A = 3 \\ -6B = 1 \\ -3C + 2B = 0 \end{array} \right\} \Rightarrow \begin{array}{l} A = 1 \\ B = -\frac{1}{6} \\ 3C = -\frac{1}{3} \Rightarrow C = -\frac{1}{9} \end{array}$$

So $Y(t) = te^t + (-\frac{1}{6}t^2 - \frac{1}{9}t)e^{-2t}$

Thus, the general solution is

$$y(t) = c_1 e^t + c_2 e^{-2t} + te^t - \frac{1}{6}t^2 e^{-2t} - \frac{1}{9}te^{-2t}$$