



ON AN UNCONDITIONALLY STABLE SCHEME FOR THE UNSTEADY NAVIER-STOKES EQUATIONS

Author(s): Shen Jie

Source: *Journal of Computational Mathematics*, July 1990, Vol. 8, No. 3, PROCEEDINGS OF THE CHINA-US SEMINAR ON BOUNDARY INTEGRAL AND BOUNDARY METHODS IN PHYSICS AND ENGINEERING (July 1990), pp. 276-288

Published by: Institute of Computational Mathematics and Scientific/Engineering Computing

Stable URL: <https://www.jstor.org/stable/43692488>

REFERENCES

Linked references are available on JSTOR for this article:

https://www.jstor.org/stable/43692488?seq=1&cid=pdf-reference#references_tab_contents

You may need to log in to JSTOR to access the linked references.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

Institute of Computational Mathematics and Scientific/Engineering Computing is collaborating with JSTOR to digitize, preserve and extend access to *Journal of Computational Mathematics*

ON AN UNCONDITIONALLY STABLE SCHEME FOR THE UNSTEADY NAVIER-STOKES EQUATIONS*

Shen Jie

(*Department of Mathematics, Indiana University, Bloomington, IN 47405, U.S.A.*)

Abstract

Theoretical time step constraints of semi-implicit schemes are known to be more restrictive than should be in practice. We intend to alleviate the constraints with more smoothness assumptions on the solutions. By introducing a new scheme with modification on the treatment of the nonlinear term, we are able to prove that the scheme is unconditionally stable and convergent. Further more, we show that the modified scheme and the original semi-implicit one are equivalent under a weak condition on the time step and the number of space discretization points.

§1. Introduction

In numerical simulations of incompressible flow represented by the Navier-Stokes equations (1.1), one of the major difficulties is to construct a suitable time discretization scheme. The origin of such difficulty consists essentially of two parts:

(i) The pressure and the velocity in Navier-Stokes equations are coupled by the incompressibility constraint (1.1b) such that a direct inversion of the resulting discrete system is very expensive. A great number of fast Stokes solvers have been developed by using either an iterative method or a Green's function method (also called influence matrix method, see for instance [8]). Another remedy for removing this difficulty is to use the so called projection method initially proposed by A.J.Chorin and R.Temam (cf. [4], [11]) which separates the calculation of the pressure from that of the velocity. However, this kind of splitting schemes suffers from a large time splitting error which can only be removed by a sophisticated extrapolation process (cf. [10]).

(ii) The treatment of the nonlinear term: usually, explicit treatment of the nonlinear term leads to in some cases a restrictive theoretical time step constraint (see for instance [12]) while implicit treatment makes the resulting discrete system very difficult to be resolved.

In this paper, we concentrate on improving existing theoretical stability constraints for semi-implicit schemes in which the diffusion term is treated implicitly, leaving the convection term (i.e. nonlinear term) treated explicitly.

In many cases, one observes that a semi-implicit scheme gives stable results under a time step constraint which is much weaker than what the theoretical results predict, especially in cases where a smooth solution exists. A natural question one can ask is: can we improve the existing stability conditions by giving more smoothness assumptions on the solutions?

We will give a positive answer to this question by considering a concrete space discretization, namely, the Chebyshev-Galerkin approximation (we refer to [7] for a detailed presentation of this method). For other space discretizations, similar results could be obtained by

* Received January 11, 1988.

using the same technique. The idea used here can also be applied to other time-dependent elliptic nonlinear systems.

The unsteady Navier-Stokes equations in the primitive variable formulation are written as

$$\frac{\partial u}{\partial t} - \nu \Delta u + u \nabla u + \nabla p = f(x, t), \quad (x, t) \in Q = \Omega \times [0, T], \tag{1.1a}$$

$$\operatorname{div} u = 0, \quad \text{in } Q, \tag{1.1b}$$

$$u(x, 0) = u_0(x), \tag{1.1c}$$

$$u(x, t) = 0|_{\partial\Omega} \quad \forall t \in [0, T], \tag{1.1d}$$

where Ω is an open set in R^d ($d = 2$ or 3) with sufficiently smooth boundary, the unknowns are the vector function u (velocity) and the scalar function p (pressure). For the sake of simplicity, we assume that the velocity satisfies the homogeneous boundary condition.

We will restrict ourselves to the two dimensional case. More specifically, we consider $\Omega = (-1, 1) \times (-1, 1)$. The Chebyshev weight function defined in Ω is

$$\omega(x) = (1 - x_1^2)^{-\frac{1}{2}}(1 - x_2^2)^{-\frac{1}{2}} \quad \text{for } x = (x_1, x_2) \in \Omega.$$

The following functional spaces will be used in the sequel:

$$\mathcal{X} = \mathcal{H}_{0,\omega}^1(\Omega),$$

$$\mathcal{H}_\omega = \{u \in \mathcal{L}_\omega^2(\Omega) : \operatorname{div} u = 0, \quad u \cdot \vec{n} = 0\},$$

$$\mathcal{V}_\omega = \{u \in \mathcal{X} : \operatorname{div} u = 0\},$$

$$\tilde{\mathcal{V}}_\omega = \{u \in \mathcal{X} : \operatorname{div}(u \cdot \omega) = 0\},$$

where ω is the Chebyshev weight function, and $\mathcal{L}_\omega^2(\Omega)$ and $\mathcal{H}_{0,\omega}^1(\Omega)$ are weighted Sobolev spaces. To alleviate notations, we use calligraphic letters to denote vector function spaces, for instance, $\mathcal{L}_\omega^2 = (L_\omega^2)^2$.

With the help of these functional spaces, we can reformulate the problem (1.1) as

$$\begin{cases} \text{find } u(t) \in \mathcal{V}_\omega \text{ such that} \\ \frac{\partial}{\partial t} t(u, v)_\omega + \nu a_\omega(u, v) + (B(u), v)_\omega = \langle f, v \rangle_\omega, \quad \forall v \in \tilde{\mathcal{V}}_\omega, \\ u(0) = u_0 \end{cases} \tag{1.2}$$

where we have

$$(u, v)_\omega = (u, v\omega) = \int_\Omega uv\omega dx, \quad a_\omega(u, v) = (\nabla u, \nabla(v \cdot \omega)),$$

$$B(u) = \sum_{i=1}^d u_i \frac{\partial u}{\partial x_i} \text{ and } \langle \cdot, \cdot \rangle_\omega \text{ the duality relation between } \mathcal{X}' \text{ and } \mathcal{X}.$$

Due to the Chebyshev weight function involved here, the formulation (1.2) is not symmetric such that the existence of solutions for (1.2) is not covered by the conventional theory.

It is proved in [5] and [9] that the bilinear form $a_\omega(\cdot, \cdot)$ is continuous and coercive on $\mathcal{X} \times \mathcal{X}$. More precisely, we have

$$\begin{cases} a_\omega(u, v) \leq \alpha \|u\|_{1,\omega} \|v\|_{1,\omega}, \quad \forall u, v \in \mathcal{X}, \\ a_\omega(u, u) \geq \beta \|u\|_{1,\omega}^2, \quad \forall u \in \mathcal{X}. \end{cases} \tag{1.3}$$

Therefore, $\|\cdot\|_\omega = a_\omega(\cdot, \cdot)^{\frac{1}{2}}$ is a norm equivalent to $\|\cdot\|_{1,\omega}$ on \mathcal{X} . We will use in the sequel the norm $\|\cdot\|_\omega$ instead of $\|\cdot\|_{1,\omega}$ and we will denote the norm on $\mathcal{L}_\omega^2(\Omega)$ by $|\cdot|_\omega$.

In order to formulate a discrete Chebyshev-Galerkin approximation of (1.2), we define

S_N : The set of polynomials such that the order of each variable is less than or equal to N ;

$$\mathcal{X}_N = S_N \cap \mathcal{X}, \mathcal{V}_N = S_N \cap \mathcal{V}_\omega \text{ and } \tilde{\mathcal{V}}_N = S_N \cap \tilde{\mathcal{V}}_\omega.$$

Let us consider first the following fully discretized scheme which consists of the second order Crank-Nicolson and Adams-Bashforth scheme in time and the Chebyshev-Galerkin scheme in space:

$$\begin{cases} \text{Let } u_N^0 = \Pi_N u_0, \text{ find } u_N^n \in \mathcal{V}_N \text{ such that} \\ \frac{1}{k}(u_N^{n+1} - u_N^n, v)_\omega + \frac{\nu}{2} a_\omega(u_N^{n+1} + u_N^n, v) = \langle f^{n+\frac{1}{2}}, v \rangle_\omega \\ -1.5(B(u_N^n), v)_\omega + 0.5(B(u_N^{n-1}), v)_\omega, \quad \forall v \in \tilde{\mathcal{V}}_N \end{cases} \tag{1.4}$$

where $k = \frac{T}{K}$: time discretization step and $f^{n+\frac{1}{2}} = \frac{1}{k} \int_{nk}^{(n+1)k} f(x, t) dt$;

Π_N : a projection operator $\mathcal{X} \rightarrow \mathcal{X}_N$ such that

$$a_\omega(u - \Pi_N u, v) = 0, \quad \forall v \in \mathcal{X}_N, u \in \mathcal{X}. \tag{1.5}$$

Throughout the paper, we will use c and c_i to denote constants which can vary from one equation to another.

We infer from (1.5) and (1.3) that

$$a_\omega(u - \Pi_N u, u - \Pi_N u) = a_\omega(u - \Pi_N u, u - \phi) \leq c \|u - \Pi_N u\|_\omega \|u - \phi\|_\omega, \quad \forall u \in \mathcal{X}, \forall \phi \in \mathcal{X}_N. \tag{1.6}$$

Since the following is true (cf. [9]):

$$\inf_{\phi \in \mathcal{X}_N} \|u - \phi\|_\omega \leq c N^{1-\sigma} \|u\|_{\sigma,\omega}, \quad \forall u \in \mathcal{H}_\omega^\sigma(\Omega) \cap \mathcal{X}$$

we deduce from (1.3) and (1.6) that

$$\|u - \Pi_N u\|_\omega \leq c N^{1-\sigma} \|u\|_{\sigma,\omega}, \quad \forall u \in \mathcal{H}_\omega^\sigma(\Omega) \cap \mathcal{X}. \tag{1.7}$$

R.Temam analyzed in [12] this kind of scheme in a general space discretization form. He introduced two quantities $S(N)$ and $S_1(N)$ defined by

$$\begin{cases} \|u_N\| \leq S(N) |u_N|, \quad \forall u_N \in S_N, \\ |((u_N \cdot \nabla) v_N, w_N)| \leq S_1(N) |u_N| \cdot \|v_N\| \cdot |w_N|, \quad \forall u_N, v_N, w_N \in \mathcal{X}_N \end{cases} \tag{1.8}$$

and proved that the semi-implicit schemes are stable and convergent under the conditions:

$$kS^2(N) \leq C_1, \quad kS_1^2(N) \leq C_2. \tag{1.9}$$

These conditions are somewhat restrictive especially when a spectral method is adapted to the space discretization. For instance, we have $S(N) = O(N^2)$ for the Chebyshev or Legendre approximations (cf. [6]). It means that we should have a time step at least less than cN^{-4} to ensure the stability. This is evidently not a reasonable constraint in practice. It is then necessary to lighten this constraint by assuming more smoothness of the solution.

In the next section, we will introduce a modified scheme which differs from (1.4) in the treatment of the nonlinear term. We will prove the new scheme is unconditionally stable provided that the solution u is uniformly bounded in Q . Then, we will prove in Section 3 that the new scheme preserves the second order accuracy in time and the spectral accuracy in space. Finally, we will prove in Section 4 that the modified scheme and the original one are equivalent under a very weak condition.

§2. A Modified Scheme and Related Stability

We assume that u is uniformly bounded in Q , i.e.

(H1) There exists $M > 0$ such that $|u|_{L^\infty(Q)} \leq M$.

Following an idea of Bressan and Quarteroni (cf. [2]), we introduce the truncated function $H: R \rightarrow R$ defined as

$$H(y) = \begin{cases} y, & \text{if } |y| \leq 2M, \\ 2M, & \text{if } y \geq 2M, \\ -2M, & \text{if } y \leq -2M. \end{cases} \tag{2.1}$$

Let us set

$$N(u(x)) = \sum_{i=1}^2 \frac{\partial}{\partial x_i} (H(u_i(x))u(x)), \quad \forall u \in \mathcal{X}.$$

We observe that

$$\begin{cases} |u|_{L^\infty(Q)} \leq 2M \\ u \in \mathcal{V} \end{cases} \quad \text{implies} \quad N(u) = B(u).$$

We consider now the following modified scheme in which we replace $B(u)$ in (1.4) by $N(u)$:

$$\begin{cases} \text{Let } u_N^0 = \Pi_N u_0, \text{ find } u_N^n \in \mathcal{V}_N \text{ such that} \\ \frac{1}{k} (u_N^{n+1} - u_N^n, v)_\omega + \frac{\nu}{2} a_\omega (u_N^{n+1} + u_N^n, v) \\ = -(1.5N(u_N^n) - 0.5N(u_N^{n-1}), v)_\omega + \langle f^{n+\frac{1}{2}}, v \rangle_\omega, \quad \forall v \in \tilde{\mathcal{V}}_N. \end{cases} \tag{2.2}$$

To start (2.1), one needs to know also u_N^1 . We assume that u_N^1 is given such that the truncation error at time $t = k$ is $O(t^2)$. Such a u_N^1 may be obtained by for instance the Runge-Kutta scheme.

At each time step, the scheme (2.2) can be interpreted as a discrete Stokes type equation which admits a unique solution as proved in [3] by means of the following lemma which is essential for our analysis.

Lemma 1. $\forall u \in \mathcal{V}_N, \exists \phi \in H^2_{0,\omega} \cap S_N$ such that

$$u = \text{rot} \phi \text{ and } \omega^{-1} \text{curl}(\phi \omega) \in \tilde{\mathcal{V}}_N.$$

We define the operator $T: \mathcal{V}_N \rightarrow \tilde{\mathcal{V}}_N$ by

$$Tu = \omega^{-1} \text{curl}(\phi \omega). \tag{2.3}$$

Then

$$(u, Tu)_\omega = (\text{curl} \phi, \text{curl}(\phi \omega)) = a_\omega(\phi, \phi) = \|\phi\|_\omega^2 \geq \alpha \|u\|_\omega^2, \tag{2.4}$$

$$a_\omega(u, Tu) = (\Delta \phi, \Delta(\phi \omega)) \geq \beta \|\phi\|_{2,\omega}^2 \geq \beta \|u\|_\omega^2. \tag{2.5}$$

where $\text{curl} \phi = (\frac{\partial \phi}{\partial y}, \frac{\partial \phi}{\partial x})$.

Let us prove first a stability result.

Lemma 2. Let $u_0 \in \mathcal{V}_\omega$ and $f \in \mathcal{L}^2(0, T, \mathcal{X}')$. We assume moreover that the solution u of (1.1) satisfies (H1). Then the scheme (2.2) is unconditionally stable. More precisely, for $0 \leq m \leq K$, we have

$$|u_N^m|_\omega^2 + k\nu \sum_{n=0}^{m-1} \|u_N^{n+1} + u_N^n\|_\omega^2 \leq L_1(f, u_N^1, u_0), \tag{2.6}$$

$$\sum_{n=0}^{m-1} |u_N^{n+1} - u_N^n|_\omega^2 \leq L_1(f, u_N^1, u_0). \tag{2.7}$$

Proof. In order to alleviate notations, we will ignore the sub-index N whenever no confusion is possible.

Replacing v by $2k \cdot T(u^{n+1} + u^n)$ in (2.2) and using Lemma 1, we obtain

$$\begin{aligned} & 2(\|\phi^{n+1}\|_\omega^2 - \|\phi^n\|_\omega^2) + k\beta\nu \|\phi^{n+1} + \phi^n\|_{2,\omega}^2 \\ & = k < 2f^{n+\frac{1}{2}} - 3N(u^n) + N(u^{n-1}), T(u^{n+1} + u^n) >_\omega. \end{aligned} \tag{2.8}$$

where ϕ^n is the function associated to u_N^n (cf. Lemma 1).

We derive from the Schwarz inequality that

$$< 2f^{n+\frac{1}{2}}, v >_\omega \leq 2\|f^{n+\frac{1}{2}}\|_{\mathcal{X}'} \|v\|_\omega \leq \frac{\nu\beta}{8} \|v\|_\omega^2 + \frac{8}{\nu\beta} \|f^{n+\frac{1}{2}}\|_{\mathcal{X}'}^2. \tag{2.9}$$

The modified nonlinear term can be easily handled thanks to the definition of the function H . Actually, we derive from integration by parts and the Schwarz inequality

$$\begin{aligned} (N(u), v)_\omega &= \sum_{i,j=1}^2 \int_\Omega \frac{\partial}{\partial x_i} (H(u_i) \cdot u_j) v_j \omega dx = - \sum_{i,j=1}^2 \int_\Omega H(u_i) u_j \frac{\partial (v_j \omega_i)}{\partial x_i} \omega_j dx \\ &\leq 2M \sum_{i,j=1}^2 \int_{-1}^1 |u_j|_{\omega_j} \cdot \left| \frac{\partial (v_j \omega_i)}{\partial x_i} \right|_{\omega_j} dx_i. \end{aligned} \tag{2.10}$$

Then, by using the following result (cf. [5]):

$$\left| \omega_i^{-1} \frac{\partial(v_j \omega_i)}{\partial x_i} \right|_{\omega_i} \leq c \|v_j\|_{\omega_i}$$

we arrive at

$$(N(u), v)_\omega \leq c_1 M \|u\|_\omega \|v\|_\omega. \tag{2.11}$$

We recall that (cf. [9])

$$\int_\Omega \phi^2 \omega^5 dx \leq \alpha_1 \|\phi\|_\omega^2 \quad \text{and} \quad \int_\Omega \phi^2 \omega^9 dx \leq \beta_1 \|\phi\|_{2,\omega}^2, \quad \forall \phi \in \mathcal{X}. \tag{2.12}$$

One can then readily check that

$$\|T(u^{n+1} + u^n)\|_\omega \leq c_2 \|\phi^{n+1} + \phi^n\|_{2,\omega}.$$

We infer from (2.9), (2.10) and (2.12) that

$$\begin{aligned} & |k < 2 f^{n+\frac{1}{2}} - 3N(u^n) + N(u^{n-1}), T(u^{n+1} + u^n) >_\omega | \\ & \leq \frac{k\nu\beta}{2} \|\phi^{n+1} + \phi^n\|_{2,\omega}^2 + \frac{8k}{\nu\beta} \|f^{n+\frac{1}{2}}\|_{\mathcal{X}'}^2 \\ & \quad + \frac{c_2 M^2 k}{\nu} (\|\phi^{n-1}\|_\omega^2 + \|\phi^n\|_\omega^2). \end{aligned} \tag{2.13}$$

Now summing (2.13) for $n = 1, \dots, m - 1$ ($m \leq K$), we get

$$\begin{aligned} & 2\|\phi^m\|_\omega^2 + \frac{k\nu\beta}{2} \sum_{n=1}^{m-1} \|\phi^{n+1} + \phi^n\|_{2,\omega}^2 \\ & \leq \frac{8k}{\nu\beta} \sum_{n=1}^{m-1} \|f^{n+\frac{1}{2}}\|_{\mathcal{X}'}^2 + 2\|\phi^1\|_\omega^2 + \frac{2c_2 M^2 k}{\nu} \sum_{n=0}^{m-1} \|\phi^n\|_\omega^2. \end{aligned} \tag{2.14}$$

(2.6) can then be established by using the following simple remark:

Let a_m, b_m and c_m be three positive sequences with c_m increasing and such that

$$a_0 + b_0 \leq c_0, \quad a_m + b_m \leq c_m + \lambda \sum_{n=0}^{m-1} a_n, \quad \forall m \geq 1$$

with $\lambda > 0$, then

$$a_m + b_m \leq \exp(\lambda m) c_m, \quad \forall m \geq 1. \tag{2.15}$$

It is proved readily by induction that

$$a_m + b_m \leq (1 + \lambda)^m c_m,$$

and (2.15) follows then from $(1 + \lambda)^m \leq \exp(\lambda m)$.

We apply (2.15) with

$$a_m = 2\|\phi^m\|_\omega^2, \quad b_m = \frac{k\nu\beta}{2} \sum_{n=1}^{m-1} \|\phi^{n+1} + \phi^n\|_{2,\omega}^2$$

and

$$c_m = \frac{8k}{\nu\beta} \sum_{n=1}^{m-1} \|f^{n+\frac{1}{2}}\|_{X'}^2 + 2\|\phi^1\|_{\omega}^2.$$

By taking into account (2.4) and (2.5), we obtain

$$\begin{aligned} & 2\alpha|u^m|_{\omega}^2 + \frac{k\nu\beta}{2} \sum_{n=0}^{m-1} \|u^{n+1} + u^n\|_{\omega}^2 \\ & \leq \left(\frac{8}{\nu\beta}\|f\|_{L^2(0,T,X')}^2 + 2|u_N^1|_{\omega}^2 + \frac{c_2M^2k}{\nu}|u_N^0|_{\omega}^2\right) \exp\left(\frac{c_2M^2T}{\nu}\right), \quad \forall 1 \leq m \leq K. \end{aligned}$$

Similarly, by the aid of the last inequality, we can prove (2.7) if we replace ν by $2kT(u_N^{n+1} - u_N^n)$ in (2.2) and proceed exactly as we did for proving (2.6).

In virtue of the previous lemma, we can prove, by using a compactness argument exactly as in §3.5 of [12], the following convergence theorem.

Theorem 1. *Let $u_{k,N}(t)$ be the function from $[0, T]$ in \mathcal{V}_N defined by*

$$u_{k,N}(t) = u_N^m, \quad \forall t \in [mk, (m+1)k], \quad m = 0, 1, \dots, K-1.$$

Given $u_0 \in \mathcal{V}_{\omega}$, $f \in L^2(0, T, X')$ and we assume that the solution u satisfies (H1). Then

$$u_{k,N}(\cdot) \rightarrow u(\cdot), \quad \text{when } k, N^{-1} \rightarrow 0$$

in $L^2(0, T, \mathcal{V}_{\omega})$ weakly, $L^{\infty}(0, T, \mathcal{H}_{\omega})$ weak-star and $L^2(0, T, \mathcal{H}_{\omega})$ strongly.

The proof of this theorem is quite long and technical. We refer to [12] for a detailed presentation.

Remark 1. Theorem 1 also implies that

$$u \in L^2(0, T, \mathcal{V}_{\omega}) \cap L^{\infty}(0, T, \mathcal{H}_{\omega}) \tag{2.16}$$

where u is the solution of (1.1). Actually, the hypothesis (H1) can be removed if we work directly on the scheme (1.4); we get instead a conditional stability and convergence result, but (2.16) still holds.

§3. Error Estimates

In this section, we will derive first an error estimate by assuming that the solution u is sufficiently smooth. Namely, we assume

$$(H2) \quad u''(t) \in L^2(0, T, \mathcal{V}_{\omega}) \quad \text{and} \quad u'''(t) \in L^2(0, T, X').$$

We then prove that, by using this error estimate and an inequality which controls the L^{∞} norm by L^2_{ω} norm in the discrete space S_N , the schemes (1.4) and (2.2) are equivalent under a very weak condition which we will describe late on.

We establish first a preliminary result explaining the truncation error of the scheme (2.2).

Lemma 3. *We assume that the solution u of (1.1) satisfies (H1) and (H2). Then*

$$k \sum_{n=1}^{K-1} \|e_n^k\|_{X'}^2 \leq L_2(u, f)k^4 \tag{3.1}$$

where e_n^k is defined by

$$\begin{aligned} \langle e_n^k, v \rangle_\omega = & \frac{1}{k}(u(n+1) - u(n), v)_\omega + \frac{\nu}{2} a_\omega(u(n+1) + u(n), v) \\ & - (f^{n+\frac{1}{2}} - 1.5N(u(n)) + 0.5N(u(n-1)), v)_\omega, \quad \forall v \in \tilde{V}_\omega \end{aligned} \tag{3.2}$$

where we have set $u(n) = u(n \cdot k)$, $n = 0, 1, \dots, K$.

Proof. We develop $u(n-1)$, $u(n)$ and $u(n+1)$ at $t = k(n + \frac{1}{2})$ by using Taylor's formula with integral residue

$$\begin{aligned} u(n-1) &= u(n + \frac{1}{2}) - 1.5ku'(n + \frac{1}{2}) + \frac{9}{4}k \int_{k(n-1)}^{k(n+\frac{1}{2})} (t - k(n-1))u''(t)dt, \\ u(n) &= u(n + \frac{1}{2}) - 0.5ku'(n + \frac{1}{2}) + \frac{1}{4}k \int_{kn}^{k(n+\frac{1}{2})} (t - kn)u''(t)dt, \\ \langle u(n), v \rangle_\omega &= \langle u(n + \frac{1}{2}) - 0.5ku'(n + \frac{1}{2}) + \frac{1}{4}k^2u''(n + \frac{1}{2}), v \rangle_\omega \\ &\quad - \frac{1}{8}k^2 \langle \int_{kn}^{k(n+\frac{1}{2})} (t - k(n + \frac{1}{2}))u'''(t)dt, v \rangle_\omega, \\ \langle u(n+1), v \rangle_\omega &= \langle u(n + \frac{1}{2}) + 0.5ku'(n + \frac{1}{2}) + \frac{1}{4}k^2u''(n + \frac{1}{2}), v \rangle_\omega \\ &\quad + \frac{1}{8}k^2 \langle \int_{k(n+\frac{1}{2})}^{k(n+1)} (t - k(n + \frac{1}{2}))u'''(t)dt, v \rangle_\omega. \end{aligned}$$

A direct computation leads to

$$\begin{aligned} & |(1.5u(n) \cdot \nabla u(n) - 0.5u(n-1) \cdot \nabla u(n-1) - u(n + \frac{1}{2}) \cdot \nabla u(n + \frac{1}{2}), v)_\omega| \\ & \leq c_1 k \int_{k(n-1)}^{k(n+\frac{1}{2})} \|u''(t)\|_\omega dt \cdot \|v\|_\omega, \\ & \left| \left(\frac{1}{k}(u(n+1) - u(n)) - u'(n + \frac{1}{2}), v \right)_\omega \right| \leq c_2 k \int_{kn}^{k(n+1)} \|u''(t)\|_{X'} dt \cdot \|v\|_\omega, \\ & \left| a_\omega(0.5(u(n+1) + u(n)) - u(n + \frac{1}{2}), v)_\omega \right| \leq c_3 k \int_{kn}^{k(n+1)} \|u'''(t)\|_{X'} dt \cdot \|v\|_\omega. \end{aligned}$$

We then subtract (2.2) from (3.2). Using the above inequalities and the relation $N(u(n)) =$

$B(u(n))$, we can get

$$\|e_n^k\|_{X'} \leq c_4 k \int_{k(n-1)}^{k(n+1)} (\|u'''(t)\|_{X'} + \|u''(t)\|_{\omega}) dt.$$

It follows that

$$\|e_n^k\|_{X'}^2 \leq c_5 k^3 \int_{k(n-1)}^{k(n+1)} (\|u'''(t)\|_{X'}^2 + \|u''(t)\|_{\omega}^2) dt.$$

The summation of the last inequality for $n = 1, \dots, K - 1$ leads to

$$k \sum_{n=1}^{K-1} \|e_n^k\|_{X'}^2 \leq c_6 k^4 (\|u'''(t)\|_{L^2(0,T,X')}^2 + \|u''(t)\|_{L^2(0,T,\nu_{\omega})}^2).$$

We can now prove the following theorem.

Theorem 2. *Assuming that the solution u satisfies (H1), (H2) and*

$$u \in C([0, T], \mathcal{H}_{\omega}^s(\Omega)) \quad , \quad u' \in C([0, T], \mathcal{H}_{\omega}^{s-1}(\Omega)), \quad s \leq 2. \tag{3.3}$$

Then

$$\begin{aligned} & |u(m) - u_N^m|_{\omega}^2 + k\nu \sum_{n=0}^{m-1} \|u(n+1) + u(n) - u_N^n - u_N^{n+1}\|_{\omega}^2 \\ & \leq L_3(k^4 + |u_N^1 - \Pi_N u(1)|_{\omega}^2 + N^{2(1-s)}) \quad , \quad 1 \leq m \leq K. \end{aligned} \tag{3.4}$$

Proof. Let us set

$$e^n = \Pi_N u(n) - u_N^n \quad \text{and} \quad \bar{e}_n = u(n) - \Pi_N u(n) \quad \text{for} \quad n = 0, 1, \dots, K.$$

We can then reformulate (3.2) in the discrete form by the help of \bar{e}_n and Π_N :

$$\begin{aligned} & \frac{1}{k} (\Pi_N (u(n+1) - u(n)), v)_{\omega} + \frac{\nu}{2} a_{\omega} (\Pi_N (u(n+1) + u(n)), v) \\ & - (f^{n+\frac{1}{2}} - 1.5N(u(n)) + 0.5N(u(n-1))), v)_{\omega} \\ & = \langle e_n^k, v \rangle_{\omega} - \frac{1}{k} (\bar{e}_{n+1} - \bar{e}_n, v)_{\omega} \quad , \quad \forall v \in \tilde{V}_N. \end{aligned} \tag{3.5}$$

Using Taylor's formula as we did in the proof of Lemma 3, we can get

$$\begin{aligned} |(\bar{e}_{n+1} - \bar{e}_n, v)_{\omega}| & \leq \left\{ |u'(n + \frac{1}{2}) - \Pi_N u'(n + \frac{1}{2})|_{\omega} + c_1 k \int_{k(n-1)}^{k(n+1)} \|u'''(t)\|_{X'} dt \right\} \|v\|_{\omega} \\ & \leq |u'(n + \frac{1}{2}) - \Pi_N u'(n + \frac{1}{2})|_{\omega}^2 + 4c_1 k^3 \int_{k(n-1)}^{k(n+1)} \|u'''(t)\|_{X'} dt + \frac{1}{2} \|v\|_{\omega}^2. \end{aligned} \tag{3.6}$$

We subtract (3.5) from (2.2)

$$\begin{aligned} & \frac{1}{k} (e^{n+1} - e^n, v)_{\omega} + \frac{\nu}{2} a_{\omega} (e^{n+1} + e^n, v) = \langle e_n^k, v \rangle_{\omega} - \frac{1}{k} (\bar{e}_{n+1} - \bar{e}_n, v)_{\omega} \\ & + 1.5(N(u(n)) - N(u^n), v)_{\omega} - 0.5(N(u(n-1)) - N(u^{n-1}), v)_{\omega} \quad , \quad \forall v \in \tilde{V}_N. \end{aligned} \tag{3.7}$$

Let us majorize first the nonlinear terms. By definition

$$\begin{aligned}
 N(u(n)) - N(u^n) &= \sum_{i=1}^2 \frac{\partial}{\partial x_i} (H(u_i(n))u(n) - H(u_i^n)u^n) \\
 &= \sum_{i=1}^2 \frac{\partial}{\partial x_i} \{u(n)(H(u_i(n)) - H(u_i^n)) + H(u_i^n)(u(n) - u^n)\}.
 \end{aligned}$$

One can readily check from the definition of H that

$$|H(x) - H(y)| \leq |x - y|, \quad \forall x, y \in R.$$

which implies that

$$|H(u) - H(v)|_\omega \leq |u - v|_\omega, \quad \forall u, v \in \mathcal{L}^2_\omega(\Omega). \tag{3.8}$$

We then take the scalar product of (3.7) with v and integrate by parts. Using (3.8), (1.5) and (H1), we obtain:

$$\begin{aligned}
 |N(u(n)) - N(u^n), v|_\omega &\leq \{2M|u(n) - u^n|_\omega + |u(n)|_{\mathcal{L}^\infty(\Omega)}|u(n) - u^n|_\omega\} \|v\|_\omega \\
 &\leq c_2(|e^n|_\omega + |\bar{e}_n|_\omega) \|v\|_\omega \leq c_2(|e^n|_\omega^2 + |\bar{e}_n|_\omega^2) + \frac{1}{2}c_2 \|v\|_\omega^2.
 \end{aligned} \tag{3.9}$$

Replacing v by $k \cdot T(e^{n+1} + e^n)$ (where T is the operator defined in Lemma 1) in (3.7) and taking into account (3.6), (3.9) and Lemma 1, we find

$$\begin{aligned}
 |e^{n+1}|_\omega^2 - |e^n|_\omega^2 + \frac{k\nu}{4} \|e^{n+1} + e^n\|_\omega^2 &\leq c_3 k \{ |e^n|_\omega^2 + |e^{n+1}|_\omega^2 + |\bar{e}_n|_\omega^2 + |\bar{e}_{n-1}|_\omega^2 + \|\epsilon_n^k\|_{\mathcal{X}}^2, \\
 &+ |u'(n + \frac{1}{2}) - \Pi_N u'(n + \frac{1}{2})|_\omega^2 + k^3 \int_{k(n-1)}^{k(n+1)} \|u'''(t)\|_{\mathcal{X}}^2 dt \}.
 \end{aligned}$$

We then sum this inequality for $n = 1, \dots, m - 1$. Using (3.1) and (H2), we get

$$\begin{aligned}
 |e^m|_\omega^2 + \frac{k\nu}{4} \sum_{n=1}^{m-1} \|e^{n+1} + e^n\|_\omega^2 &\leq c_4 k^4 + |e^1|_\omega^2 + c_5 k \sum_{n=0}^{m-1} |e^n|_\omega^2 \\
 &+ c_5 k \sum_{n=1}^{m-1} \{ |u'(n + \frac{1}{2}) - \Pi_N u'(n + \frac{1}{2})|_\omega^2 + |\bar{e}_n|_\omega^2 \}.
 \end{aligned}$$

We can now apply (2.15) to the last inequality, which gives

$$\begin{aligned}
 |e^m|_\omega^2 + k\nu \sum_{n=1}^{m-1} \|e^{n+1} + e^n\|_\omega^2 &\leq c_6 k^4 + |e^1|_\omega^2 + c_6 k \sum_{n=0}^{m-1} |\bar{e}_n|_\omega^2 \\
 &+ c_6 k |e^0|_\omega^2 + c_6 k \sum_{n=0}^{m-1} \{ |u'(n + \frac{1}{2}) - \Pi_N u'(n + \frac{1}{2})|_\omega^2 \}.
 \end{aligned}$$

The proof is complete by combining (1.7), (3.3) and the relations

$$|u(n) - u^n|_\omega \leq |e^n|_\omega + |\bar{e}_n|_\omega,$$

$$|\bar{e}_n|_\omega \leq c \|\bar{e}_n\|_\omega \quad \text{and} \quad e^0 = 0$$

into the last inequality.

By using Theorem 2, we can now prove the following interesting result.

Theorem 3. *We suppose that all the conditions of Theorem 2 are satisfied. Then, there exists $N_0 > 0$ such that the schemes (1.4) and (2.2) are equivalent as long as $N > N_0$ and $kN^{\frac{1}{2}} \rightarrow 0$.*

Proof. It is sufficient to prove that

$$\max_{0 \leq n \leq K} |u_N^n - u(n)|_\infty \rightarrow 0 \quad (N \rightarrow \infty) \quad \text{provided} \quad kN^{\frac{1}{2}} \rightarrow 0.$$

Let us prove first that

$$|q|_\infty \leq N|q|_\omega, \quad \forall q \in S_N,$$

which is a kind of inverse inequality.

We recall the following orthogonality formula satisfied by the Chebyshev polynomials:

$$\int_{-1}^1 T_k(x)T_l(x)\omega(x)dx = \frac{\pi}{2}c_k\delta_{kl}$$

where $c_0 = 2$ and $c_k = 1$ for $k \geq 1$.

We infer from this formula that

$$\forall q = \sum_{k,l=0}^N q_{kl}T_k(x)T_l(y), \quad |q|_\omega^2 = \frac{\pi^2}{4} \sum_{k,l=0}^N c_k c_l q_{kl}^2.$$

Since $|T_k(x)| \leq 1, \forall x \in I$, we deduce that

$$|q|_\infty^2 = \left| \sum_{k,l=0}^N q_{kl}T_k(x)T_l(y) \right|_\infty^2 \leq \left\{ \sum_{k,l=0}^N |q_{kl}| \right\}^2 \leq (N+1)^2 \sum_{k,l=0}^N (q_{kl})^2 \leq \frac{4}{\pi^2} (N+1)^2 |q|_\omega^2.$$

We know from Theorem 2 that

$$\max_{m=0,1,\dots,K} |u_N^m - \Pi_N u(m)|_\omega^2 \leq c(k^4 + |u_N^1 - \Pi_N u(1)|_\omega^2 + N^{2(1-\sigma)}).$$

The last two inequalities imply that

$$\begin{aligned} |u_N^m - u(m)|_\infty^2 &\leq 2(|u_N^m - \Pi_N u(m)|_\infty^2 + |u(m) - \Pi_N u(m)|_\infty^2) \\ &\leq \frac{8(N+1)^2}{\pi^2} |u_N^m - \Pi_N u(m)|_\omega^2 + |u(m) - \Pi_N u(m)|_\infty^2 \\ &\leq cN^2(k^4 + |u_N^1 - \Pi_N u(1)|_\omega^2 + N^{2(1-\sigma)}) + |u(m) - \Pi_N u(m)|_\infty^2. \end{aligned} \tag{3.10}$$

The first term on the right-hand side of (3.10) tends to zero provided that

$$kN^{\frac{1}{2}} \rightarrow 0. \tag{3.11}$$

In virtue of the following Sobolev inequality (cf. for instance [1]):

$$|u|_\infty^2 \leq c \|u\|_2 \cdot |u|, \quad \forall u \in H^2(\Omega) \tag{3.12}$$

and by taking into account (1.7) and (3.3), the second term on the right-hand side of (3.10) also tends to zero. It means that

$$\max_{0 \leq n \leq K} |u_N^n - u(n)|_\infty \rightarrow 0, \quad N \rightarrow 0$$

under the condition (3.11).

Consequently, there exists $N_0 > 0$ such that

$$\max_{0 \leq n \leq K} |u_N^n - u(n)|_\infty \leq M, \quad \forall N \leq N_0.$$

Hence

$$\begin{aligned} \max_{m=0,1,\dots,K} |u_N^m|_\infty &\leq \max_{0 \leq n \leq K} |u_N^n - u(n)|_\infty + \max_{0 \leq n \leq K} |u(n)|_\infty \\ &\leq M + M = 2M \end{aligned}$$

which implies $N(u_N^m) = B(u_N^m)$.

Remark 2. By giving more smoothness assumptions on the solution u , we can reduce the very restrictive time step constraint (1.9) for the scheme (1.4) to the very weak condition (3.11) which is evidently not a constraint in practice since we should keep the time step k reasonably small to balance the spectral precision of the space discretization.

Acknowledgement. This work consists of part of my thesis done at the University of South-Paris under the direction of Professor Roger Temam. I would like to express my deep thanks to him for his continuous guidance and encouragement.

References

- [1] S. Agmon, Lectures on Elliptic Boundary Value Problems, New-York, 1965.
- [2] N. Bressan and A. Quarteroni, An implicit/explicit spectral method for the Burger's equations, *Calcolo*, **23**, 265-284.
- [3] C. Bernardi, C. Canuto and Y. Maday, Generalized inf-sup condition for Chebyshev approximation of the Navier-Stokes equations, *C.R. Acad. Sci., Paris* 303, serie I, 971-974.
- [4] A. J. Chorin, Numerical solution of the Navier-Stokes equations, *Math. Comp.*, **23** (1968), 341-354.
- [5] C. Canuto and A. Quarteroni, Spectral and pseudo-spectral methods for parabolic problems with non-periodic boundary conditions, *Calcolo*, vol. XVIII, fasc. III, 1981.
- [6] C. Canuto and A. Quarteroni, Approximation results for orthogonal polynomials in Sobolev spaces, *Math. Comp.*, **38** : 157 (1982), 67-86.
- [7] D. Gottlieb and S. A. Orszag, Numerical analysis of spectral methods, theory and application, CBMS Regional Conference Series in Applied Mathematics, SIAM, Philadelphia, 1977.
- [8] L. Kleiser and Schumann, Treatment of incompressibility and boundary conditions in 3-D numerical spectral simulation of plane channel flow, Proc. of the 3th GAMM conference on numer. methods in fluid mechanics, ed. by E. H. Hirshel (Viewig-Verlag Braunschweig), 1980, 165-173.

- [9] Y.Maday and B.Métivet, Chebyshev-spectral approximation of Navier-Stokes equations in a two dimensional domain, *Modélisation Mathématique et Analyse Numérique*, **21** : 1 (1987), 93-123.
- [10] S.A.Orszag, M.Israeli and M.Deville, Boundary condition for incompressible flow, *J. Sci. Comp.*, No. 2, 1987.
- [11] R.Temam, Sur l'approximation de la solution des équations de Navier-Stokes par la méthode des pas fractionnaires II, *Arch.Rat.Mech.Anal.*, **33** (1969), 377-385.
- [12] R.Temam, Navier-Stokes equations: Theory and Numerical Analysis, North Holland, 1979.