


6.2 Regions Between Curves

The area under $y = f(x)$ above the x -axis between $x = a$ and $x = b$

is $\int_a^b f(x) dx$


Area of a thin rectangle
height $f(x)$
width dx



$$\text{area} = \int_a^b f(x) dx$$

Sum of infinitely many $f(x) dx$ as x goes from a to b

this idea can be extended to find area between two curves



each rectangle has area $= [f(x) - g(x)] dx$

use integration to accumulate

$$\int_a^b [f(x) - g(x)] dx$$


top bottom

example Find the area of region bounded by

$$y = x^2 - 2x \quad \text{and} \quad y = 2x$$

parabola
opens up

line



$$y = x^2 - 2x$$

$$= x(x-2) \rightarrow x\text{-intercepts } (y=0) \text{ at } x=0, x=2$$

the left end of the region:
the right end of the region:

intersections of $y = x^2 - 2x$ and $y = 2x$

$$x^2 - 2x = 2x$$


$$x^2 - 4x = 0 \rightarrow x(x-4) = 0 \quad x=0, x=4$$

rectangle height: $2x - (x^2 - 2x) = 4x - x^2$

" width: dx

$$\text{area of region: } \int_0^4 (4x - x^2) dx = 4 \cdot \frac{x^2}{2} - \frac{x^3}{3} \Big|_0^4 = 2x^2 - \frac{1}{3}x^3 \Big|_0^4 = \left(32 - \frac{64}{3}\right) - 0 = \boxed{\frac{32}{3}}$$

Example Find area of region bound by $y = \cos x$ and $y = \sin 2x$ between $x = 0$ and $x = \frac{\pi}{2}$



$$\sin \frac{\pi}{2} = 1$$

$$\sin 2x = 1 \rightarrow 2x = \frac{\pi}{2} \text{ or } \frac{3\pi}{2}$$

$$x = \frac{\pi}{4} \text{ or } \frac{3\pi}{4}$$

①: $\cos x$ above
 $\sin 2x$ below

②: $\sin 2x$ above
 $\cos x$ below

Intersection: $\cos x = \sin 2x$

$$\cos x = 2 \sin x \cos x$$

$$\cos x - 2 \sin x \cos x = 0$$

$$\cos x (1 - 2 \sin x) = 0$$

$$\cos x = 0, \quad 1 - 2 \sin x = 0$$

$$\sin x = \frac{1}{2}$$

$$x = \frac{\pi}{2}$$

$$x = \frac{\pi}{6}$$

$$\underbrace{\int_0^{\pi/6} (\cos x - \sin 2x) dx}_{(1)} + \underbrace{\int_{\pi/6}^{\pi/2} (\sin 2x - \cos x) dx}_{(2)}$$

$$\int \cos x dx = \sin x + C$$


$$\int \sin x dx = -\cos x + C$$

$$= \left(\sin x + \frac{1}{2} \cos 2x \right) \Big|_0^{\pi/6} + \left(-\frac{1}{2} \cos 2x - \sin x \right) \Big|_{\pi/6}^{\pi/2}$$

$$= \left(\sin \frac{\pi}{6} + \frac{1}{2} \cos \frac{\pi}{3} \right) - \left(\sin 0 + \frac{1}{2} \cos 0 \right) \\ + \left(-\frac{1}{2} \cos \pi - \sin \frac{\pi}{2} \right) - \left(-\frac{1}{2} \cos \frac{\pi}{3} - \sin \frac{\pi}{6} \right)$$

$$= \left(\frac{1}{2} + \frac{1}{4} \right) - \left(0 + \frac{1}{2} \right) + \left(\frac{1}{2} - 1 \right) - \left(-\frac{1}{4} - \frac{1}{2} \right) = \boxed{\frac{1}{2}}$$

if we can integrate in terms of x , we can integrate in terms of y




$$\text{area} = \int_c^d [f(y) - g(y)] dy$$

example

$$x = y^2 - 6, \quad x = y$$

parabola
opening RIGHT

↳ line




$$x = y^2 - 6 \quad \text{find } y\text{-ints (} x=0 \text{)}$$
$$0 = y^2 - 6 \quad y = \pm \sqrt{6}$$

if we use vertical rectangles,
then the roles of "top" and
"bottom" switch/change
at some point

→ ok, but maybe there is
something easier

horizontal rectangles are a little easier



this time, "right" is always $x = y$
"left" is always $x = y^2 - 6$
no switching!

intersections: $y^2 - 6 = y$
 $y^2 - y - 6 = 0$
 $(y - 3)(y + 2) = 0$
 $y = -2, y = 3$

$$\int_{-2}^3 \left[\underset{\text{right}}{y} - \underset{\text{left}}{(y^2 - 6)} \right] dy = \int_{-2}^3 (y - y^2 + 6) dy$$
$$= \left. \frac{y^2}{2} - \frac{y^3}{3} + 6y \right|_{-2}^3 = \dots = \boxed{\frac{125}{6}}$$