

EXAMPLES OF SECTIONS 6.1

Question 1. T is a linear transformation from \mathbb{P}_2 to \mathbb{P}_2 , \mathbb{P}_2 is the space of all polynomials of degree no more than 2, and

$$T(x^2 - 1) = x^2 + x - 3, \quad T(2x) = 4x, \quad T(3x + 2) = 2x + 6.$$

Find $T(1)$, $T(x)$, and $T(x^2)$.

Solutions.

1. We identify T as a linear transformation from \mathbb{R}^3 to \mathbb{R}^3 by the map

$$ax^2 + bx + c \mapsto \begin{bmatrix} a \\ b \\ c \end{bmatrix}.$$

By the given conditions, we have

$$T(1, 0, -1) = (1, 1, -3), \quad T(0, 2, 0) = (0, 4, 0), \quad T(0, 3, 2) = (0, 2, 6).$$

We immediately have

$$T(0, 1, 0) = \frac{1}{2}T(0, 2, 0) = (0, 2, 0).$$

Let

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 3 \\ -1 & 0 & 2 \end{bmatrix}.$$

Solving

$$A\vec{x} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix},$$

we get $\vec{x} = \begin{bmatrix} 0 \\ -\frac{3}{4} \\ \frac{1}{2} \end{bmatrix}$. Therefore,

$$T(0, 0, 1) = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 4 & 2 \\ -3 & 0 & 6 \end{bmatrix} \begin{bmatrix} 0 \\ -\frac{3}{4} \\ \frac{1}{2} \end{bmatrix} = \begin{bmatrix} 0 \\ -2 \\ 3 \end{bmatrix}.$$

Finally,

$$T(1, 0, 0) = T(1, 0, -1) + T(0, 0, 1) = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}.$$

Now we restore this result back to the space \mathbb{P}_2 and obtain

$$T(1) = -2x + 3, \quad T(x) = 2x, \quad T(x^2) = x^2 - x.$$