

Smooth Density for Some Nilpotent Rough Differential Equations

Yaozhong Hu · Samy Tindel

Received: 11 April 2011 / Revised: 3 August 2011 / Published online: 13 October 2011
© Springer Science+Business Media, LLC 2011

Abstract In this note, we provide a nontrivial example of a differential equation driven by a fractional Brownian motion with Hurst parameter $1/3 < H < 1/2$ whose solution admits a smooth density with respect to Lebesgue measure. The result is obtained through the use of an explicit representation of the solution when the vector fields of the equation are nilpotent, plus a Norris-type lemma in the rough paths context.

Keywords Fractional Brownian motion · Rough paths · Malliavin calculus · Nilpotent · Hypocoellipticity

Mathematics Subject Classification (2000) 60H07 · 60H10 · 65C30

1 Introduction

Let $B = (B^1, \dots, B^d)$ be a d -dimensional fractional Brownian motion with Hurst parameter $1/3 < H < 1/2$, defined on a complete probability space $(\Omega, \mathcal{F}, \mathbf{P})$. Recall that this means that all the components B^i of B are independent centered Gaussian processes with covariance

$$R_H(t, s) := \mathbf{E}[B_t^i B_s^i] = \frac{1}{2}(s^{2H} + t^{2H} - |t - s|^{2H}). \quad (1)$$

Y. Hu (✉)
Department of Mathematics, University of Kansas, Lawrence, KS 66045, USA
e-mail: hu@math.ku.edu

S. Tindel
Institut Élie Cartan Nancy, Université de Nancy 1, B.P. 239, 54506 Vandœuvre-lès-Nancy Cedex,
France
e-mail: tindel@iecn.u-nancy.fr

In particular, the paths of B are γ -Hölder continuous for all $\gamma \in (0, H)$. This paper is concerned with a class of \mathbb{R}^m -valued stochastic differential equations driven by B , of the form

$$dy_t = \sum_{i=1}^d V_i(y_t) dB_t^i, \quad t \in [0, T], \quad y_0 = a, \tag{2}$$

where $T > 0$ is a fixed time horizon, $a \in \mathbb{R}^m$ stands for a given initial condition, and (V_1, \dots, V_d) is a family of smooth vector fields of \mathbb{R}^m .

Stochastic differential systems driven by fractional Brownian motion have been the object of intensive studies during the past decade, both for their theoretical interest and for the wide range of applications they open, covering, for instance, finance [13, 30] or biophysics [20, 28] situations. The first aim in the theory has thus been to settle some reasonable tools allowing one to solve equations of type (2). This has been achieved, when the Hurst parameter H of the underlying fBm is $> 1/2$, thanks to methods of fractional integration [26, 32], or simply by means of Young-type integration (see, e.g., [14]). When one moves to more irregular cases, namely $H < 1/2$, the standard method by now in order to solve equations like (2) relies on rough paths considerations, as explained, for instance, in [12, 14, 21].

A second natural step in the study of fractional differential systems consists in establishing some properties about their probability law. Some substitute for the semi-group property governing $\mathcal{L}(y_t)$ in the Markovian case (namely where $H = 1/2$) has been given in [2, 23], in terms of asymptotic expansions in a neighborhood of $t = 0$. Some considerable efforts have also been made in order to analyze the density of $\mathcal{L}(y_t)$ with respect to Lebesgue measure. To that respect, in the regular case $H > 1/2$ the situation is rather clear: the existence of a density is shown in [27] under some standard nondegeneracy conditions, the smoothness of the density is established in [19] under elliptic conditions on the coefficients, and this result is extended to the hypoelliptic case in [3]. In all, this set of results replicates what has been obtained for the usual Brownian motion, at the price of huge technical complications.

In the irregular case $H < 1/2$, the picture is far from being so complete. Indeed, the existence part of the density results have been thoroughly studied under elliptic and Hörmander conditions (see [6, 12] for a complete review). However, when one wishes to establish the smoothness of the density, some strong moment assumptions on the inverse of the Malliavin derivative of y_t are usually required. These moment estimates are still an important open question in the field, as well as the smoothness of density for random variables like y_t .

The current paper proposes to make a step in this direction, and we wish to prove that $\mathcal{L}(y_t)$ can be decomposed as $p_t(z) dz$ for a smooth function p_t in some special nontrivial examples of (2). Namely, we will handle in the sequel the case of nilpotent vector fields V_1, \dots, V_d (see Hypothesis 4.1 for a precise description), and in this context we shall derive the following density result:

Theorem 1.1 *Suppose that the vector fields $V_i, 1 = 1, 2, \dots, d$, are smooth with all derivatives bounded and that they are n -nilpotent in the sense that their Lie brackets of order n vanish for some positive integer n . We also assume that V_1, \dots, V_d satisfy Hörmander’s hypoelliptic condition (their Lie brackets generate \mathbb{R}^m at any point*

$x \in \mathbb{R}^m$) and that all the Lie brackets of order greater than or equal to 2 are constant. Then for all $t > 0$, the probability law of the random variable y_t defined by (2) admits a smooth density with respect to Lebesgue measure.

Notice that the hypoelliptic assumption is quite natural in our context. Indeed, it would certainly be too restrictive to consider a family of vector fields V_1, \dots, V_d being nilpotent and elliptic at the same time. Moreover, some interesting examples of equations satisfying our standing assumptions will be given below. It should be stressed however that the basic aim of this article is to prove that smoothness of density results can be obtained for rough differential equations driven by a fractional Brownian motion in some specific situations, even if the general hypoelliptic case is still an important open problem. We refer to [4] for another case, based on skew-symmetric properties, where a similar theorem holds.

In order to prove Theorem 1.1, two main ingredients have to be highlighted:

- (i) Working under the nilpotent assumptions described above enables us to use a Strichartz-type representation for the solution to our equation, given in terms of a finite chaos expansion. This allows us to derive some bounds for the moments of both y_t and its Malliavin derivative, which is the main missing tool on the way to smoothness-of-density results for rough differential equations in the general case.
- (ii) With the integrability of Malliavin derivative in hand, we shall follow the standard probabilistic way to prove the smoothness of density under Hörmander's conditions, for which we refer to [16, 22, 24]. To this purpose, the second main ingredient is a Norris-type lemma, which has to be extended (in the rough path context) to *controlled processes*. It should be mentioned at this point that a similar result has been proven recently (and independently) in [17].

These two ingredients will be developed in the remainder of the article.

Here is how our article is structured: Some preliminaries on rough differential equations and fractional Brownian motion are given in Sect. 2. Section 3 is devoted to the proof of our Norris-type lemma for controlled processes in the sense of [14]. Finally, Malliavin calculus tools and their application to density results for the random variable y_t are presented at Sect. 4.

Notation In the remainder of the article, c, c_1, c_2 will stand for generic positive constants which may change from line to line. We also write $a \lesssim b$ (resp. $a \asymp b$) when $a \leq cb$ (resp. $a = cb$) for a universal constant c .

2 Rough Differential Equations and Fractional Brownian Motion

Generalized integrals will be needed in the sequel in order to define and solve equations of the form (2) and also to get an equivalent of Norris lemma in our context. Though all those elements might be obtained within the landmark of usual rough paths setting [12, 21], we have chosen here to work with the algebraic integration framework, which (from our point of view) is more amenable to handy calculations.

In this section, we recall thus the main concepts of algebraic integration. Namely, we state the definition of the spaces of increments of the operator δ and its inverse called Λ (or sewing map). We also recall some elementary but useful algebraic relations on the spaces of increments. The interested reader is sent to [14] for a complete account on the topic or to [8, 15] for a more detailed summary.

2.1 Increments

The extended integral we deal with is based on the notion of increments, together with an elementary operator δ acting on them.

The notion of increment can be introduced in the following way: for two arbitrary real numbers $\ell_2 > \ell_1 \geq 0$, a vector space V , and an integer $k \geq 1$, we denote by $\mathcal{C}_k([\ell_1, \ell_2]; V)$ the set of continuous functions $g : [\ell_1, \ell_2]^k \rightarrow V$ such that $g_{t_1 \dots t_k} = 0$ whenever $t_i = t_{i+1}$ for some $i \in \{0, \dots, k - 1\}$. Such a function will be called a $(k - 1)$ -*increment*, and we will set $\mathcal{C}_*([\ell_1, \ell_2]; V) = \bigcup_{k \geq 1} \mathcal{C}_k([\ell_1, \ell_2]; V)$. To simplify the notation, we will write $\mathcal{C}_k(V)$ if there is no ambiguity about $[\ell_1, \ell_2]$.

The operator δ is an operator acting on k -increments and is defined as follows on $\mathcal{C}_k(V)$:

$$\delta : \mathcal{C}_k(V) \rightarrow \mathcal{C}_{k+1}(V), \quad (\delta g)_{t_1 \dots t_{k+1}} = \sum_{i=1}^{k+1} (-1)^i g_{t_1 \dots \hat{t}_i \dots t_{k+1}}, \tag{3}$$

where \hat{t}_i means that this particular argument is omitted. Then a fundamental property of δ , which is easily verified, is that $\delta\delta = 0$, where $\delta\delta$ is considered as an operator from $\mathcal{C}_k(V)$ to $\mathcal{C}_{k+2}(V)$. We will denote $\mathcal{ZC}_k(V) = \mathcal{C}_k(V) \cap \text{Ker } \delta$ and $\mathcal{BC}_k(V) = \mathcal{C}_k(V) \cap \text{Im } \delta$.

Some simple examples of actions of δ , which will be the ones we will really use throughout the article, are obtained by letting $g \in \mathcal{C}_1(V)$ and $h \in \mathcal{C}_2(V)$. Then, for any $t, u, s \in [\ell_1, \ell_2]$, we have

$$(\delta g)_{st} = g_t - g_s \quad \text{and} \quad (\delta h)_{sut} = h_{st} - h_{su} - h_{ut}. \tag{4}$$

Our future discussions will mainly rely on k -increments with $k = 2$ or $k = 3$, for which we will use some analytical assumptions. Namely, we measure the size of these increments by Hölder norms defined in the following way: for $f \in \mathcal{C}_2(V)$, let

$$\|f\|_\mu = \sup_{s,t \in [\ell_1, \ell_2]} \frac{|f_{st}|}{|t - s|^\mu} \quad \text{and} \quad \mathcal{C}_2^\mu(V) = \{f \in \mathcal{C}_2(V); \|f\|_\mu < \infty\}. \tag{5}$$

Using this notation, we define in a natural way $\mathcal{C}_1^\mu(V) = \{f \in \mathcal{C}_1(V); \|\delta f\|_\mu < \infty\}$. In the sequel, we also handle norms, including supremums, of the form

$$\|f\|_{\mu, \infty} = \|f\|_\mu + \|f\|_\infty, \quad \text{and} \quad \mathcal{C}_1^{\mu, 0}(V) = \{f \in \mathcal{C}_1(V); \|f\|_{\mu, \infty} < \infty\}. \tag{6}$$

In the same way, for $h \in \mathcal{C}_3(V)$, we set

$$\|h\|_{\gamma, \rho} = \sup_{s, u, t \in [\ell_1, \ell_2]} \frac{|h_{sut}|}{|u - s|^\gamma |t - u|^\rho}, \tag{7}$$

$$\|h\|_\mu = \inf \left\{ \sum_i \|h_i\|_{\rho_i, \mu - \rho_i}; h = \sum_i h_i, 0 < \rho_i < \mu \right\},$$

where the last infimum is taken over all sequences $\{h_i, i \in \mathbb{N}\} \subset \mathcal{C}_3(V)$ such that $h = \sum_i h_i$ and over all choices of the numbers $\rho_i \in (0, \mu)$. Then $\|\cdot\|_\mu$ is easily seen to be a norm on $\mathcal{C}_3(V)$, and we define

$$\mathcal{C}_3^\mu(V) := \{h \in \mathcal{C}_3(V); \|h\|_\mu < \infty\}.$$

Eventually, let $\mathcal{C}_3^{1+}(V) = \bigcup_{\mu > 1} \mathcal{C}_3^\mu(V)$, and note that the same kind of norms can be considered on the spaces $\mathcal{ZC}_3(V)$, leading to the definition of the spaces $\mathcal{ZC}_3^\mu(V)$ and $\mathcal{ZC}_3^{1+}(V)$. In order to avoid ambiguities, we sometimes denote in the following by $\mathcal{N}[\cdot; \mathcal{C}_j^\kappa]$ the κ -Hölder norm on the space \mathcal{C}_j for $j = 1, 2, 3$. For $\zeta \in \mathcal{C}_j(V)$, we also set $\mathcal{N}[\zeta; \mathcal{C}_j^0(V)] = \sup_{s \in [\ell_1; \ell_2]^j} \|\zeta_s\|_V$.

The invertibility of δ under Hölder regularity conditions is an essential tool for the construction of our generalized integrals and can be summarized as follows:

Theorem 2.1 (The sewing map) *Let $\mu > 1$. For any $h \in \mathcal{ZC}_3^\mu(V)$, there exists a unique $\Delta h \in \mathcal{C}_2^\mu(V)$ such that $\delta(\Delta h) = h$. Furthermore,*

$$\|\Delta h\|_\mu \leq \frac{1}{2 - 2\mu} \mathcal{N}[h; \mathcal{C}_3^\mu(V)]. \tag{8}$$

This gives rise to a continuous linear map $\Lambda : \mathcal{ZC}_3^\mu(V) \rightarrow \mathcal{C}_2^\mu(V)$ such that $\delta\Lambda = id_{\mathcal{ZC}_3^\mu(V)}$.

Proof The original proof of this result can be found in [14]. We refer to [8, 15] for two simplified versions. □

The sewing map creates a first link between the structures we just introduced and the problem of integration of irregular functions:

Corollary 2.2 (Integration of small increments) *For any 1-increment $g \in \mathcal{C}_2(V)$ such that $\delta g \in \mathcal{C}_3^{1+}$, set $h = (id - \Lambda\delta)g$. Then, there exists $f \in \mathcal{C}_1(V)$ such that $h = \delta f$ and*

$$\delta f_{st} = \lim_{|\Pi_{st}| \rightarrow 0} \sum_{i=0}^n g_{t_i t_{i+1}},$$

where the limit is over any partition $\Pi_{st} = \{t_0 = s, \dots, t_n = t\}$ of $[s, t]$ whose mesh tends to zero. The 1-increment δf is the indefinite integral of the 1-increment g .

We also need some product rules for the operator δ . For this, recall the following convention: for $g \in C_n([\ell_1, \ell_2]; \mathbb{R}^{l,d})$ and $h \in C_m([\ell_1, \ell_2]; \mathbb{R}^{d,p})$, let gh be the element of $C_{n+m-1}([\ell_1, \ell_2]; \mathbb{R}^{l,p})$ defined by

$$(gh)_{t_1, \dots, t_{m+n-1}} = g_{t_1, \dots, t_n} h_{t_n, \dots, t_{m+n-1}} \tag{9}$$

for $t_1, \dots, t_{m+n-1} \in [\ell_1, \ell_2]$. With this notation, the following elementary rule holds:

Proposition 2.3 *Let $g \in C_2([\ell_1, \ell_2]; \mathbb{R}^{l,d})$ and $h \in C_1([\ell_1, \ell_2]; \mathbb{R}^d)$. Then gh is an element of $C_2([\ell_1, \ell_2]; \mathbb{R}^l)$ and $\delta(gh) = \delta gh - g\delta h$.*

2.2 Random Differential Equations

One of the main appeals of the algebraic integration theory is that differential equations driven by a γ -Hölder signal x can be defined and solved rather quickly in this setting. In the case of an Hölder exponent $\gamma > 1/3$, the required structures are just the notion of *controlled processes* and the Lévy area based on x .

Indeed, recall that we wish to consider an equation of the form

$$dy_t = \sum_{i=1}^d V_i(y_t) dx_t^i, \quad t \in [0, T], \quad y_0 = a, \tag{10}$$

where a is a given initial condition in \mathbb{R}^m , x is an element of $C_1^\gamma([0, T]; \mathbb{R}^d)$, and (V_1, \dots, V_d) is a family of smooth vector fields of \mathbb{R}^m . Then it is natural that the increments of a candidate for a solution to (10) should be controlled by the increments of x in the following way:

Definition 2.4 Let z be a path in $C_1^\kappa(\mathbb{R}^m)$ with $1/3 < \kappa \leq \gamma$, and set $\delta x := \mathbf{x}^1$. We say that z is a weakly controlled path based on x if $z_0 = a$ with $a \in \mathbb{R}^m$ and if $\delta z \in C_2^\kappa(\mathbb{R}^m)$ has a decomposition $\delta z = \zeta \mathbf{x}^1 + r$, that is, for any $s, t \in [0, T]$,

$$\delta z_{st} = \zeta_s^j \mathbf{x}_{st}^{1,j} + r_{st}, \tag{11}$$

where we have used the summation over repeated indices convention, and with $\zeta^1, \dots, \zeta^d \in C_1^\kappa(\mathbb{R}^m)$, as well as $r \in C_2^{2\kappa}(\mathbb{R}^m)$.

The space of weakly controlled paths will be denoted by $\mathcal{Q}_{\kappa,a}^x(\mathbb{R}^m)$, and a process $z \in \mathcal{Q}_{\kappa,a}^x(\mathbb{R}^m)$ can be considered in fact as a couple (z, ζ) . The space $\mathcal{Q}_{\kappa,a}^x(\mathbb{R}^m)$ is endowed with a natural seminorm given by

$$\mathcal{N}[z; \mathcal{Q}_{\kappa,a}^x(\mathbb{R}^m)] = \mathcal{N}[z; C_1^\kappa(\mathbb{R}^m)] + \sum_{j=1}^d \mathcal{N}[\zeta^j; C_1^{\kappa,0}(\mathbb{R}^m)] + \mathcal{N}[r; C_2^{2\kappa}(\mathbb{R}^m)], \tag{12}$$

where the quantities $\mathcal{N}[g; C_1^\kappa]$ have been defined in Sect. 2.1. For the Lévy area associated to x , we assume the following structure:

Hypothesis 2.5 *The path $x : [0, T] \rightarrow \mathbb{R}^d$ is γ -Hölder continuous with $\frac{1}{3} < \gamma \leq 1$ and admits a so-called Lévy area, that is, a process $\mathbf{x}^2 \in \mathcal{C}_2^{2\gamma}(\mathbb{R}^{d,d})$, which satisfies $\delta \mathbf{x}^2 = \mathbf{x}^1 \otimes \mathbf{x}^1$, namely*

$$\delta \mathbf{x}_{sut}^{2,ij} = [\mathbf{x}^{1,i}]_{su} [\mathbf{x}^{1,j}]_{ut}$$

for any $s, u, t \in [0, T]$ and $i, j \in \{1, \dots, d\}$.

To illustrate the idea behind the construction of the generalized integral, assume that the paths x and z are smooth and also for simplicity that $d = m = 1$. Then the Riemann–Stieltjes integral of z with respect to x is well defined, and we have

$$\int_s^t z_u dx_u = z_s(x_t - x_s) + \int_s^t (z_u - z_s) dx_u = z_s \mathbf{x}_{st}^1 + \int_s^t (\delta z)_{su} dx_u$$

for $\ell_1 \leq s \leq t \leq \ell_2$. If z admits the decomposition (11), we obtain

$$\int_s^t (\delta z)_{su} dx_u = \int_s^t (\zeta_s \mathbf{x}_{su}^1 + \rho_{su}) dx_u = \zeta_s \int_s^t \mathbf{x}_{su}^1 dx_u + \int_s^t \rho_{su} dx_u. \tag{13}$$

Moreover, if we set

$$\mathbf{x}_{st}^2 := \int_s^t \mathbf{x}_{su}^1 dx_u, \quad \ell_1 \leq s \leq t \leq \ell_2,$$

then it is quickly verified that \mathbf{x}^2 is the Lévy area associated to x . Hence we can write

$$\int_s^t z_u dx_u = z_s \mathbf{x}_{sz}^1 + \zeta_s \mathbf{x}_{st}^2 + \int_s^t \rho_{su} dx_u.$$

Now recast this equation as

$$\int_s^t \rho_{su} dx_u = \int_s^t z_u dx_u - z_s \mathbf{x}_{st}^1 - \zeta_s \mathbf{x}_{st}^2 \tag{14}$$

and apply the increment operator δ to both sides of this equation. For smooth paths z and x , we have

$$\delta \left(\int z dx \right) = 0, \quad \delta(z \mathbf{x}^1) = -\delta z \mathbf{x}^1,$$

by Proposition 2.3 (recall also our convention (9) on products of increments). Hence, applying these relations to the right-hand side of (14) and using decomposition (11), the properties of the Lévy area, and again Proposition 2.3, we obtain

$$\begin{aligned} \left[\delta \left(\int \rho dx \right) \right]_{sut} &= \delta z_{su} \mathbf{x}_{ut}^1 + \delta \zeta_{su} \mathbf{x}_{ut}^2 - \zeta_s \delta \mathbf{x}_{ut}^2 \\ &= \zeta_s \mathbf{x}_{su}^1 \mathbf{x}_{ut}^1 + \rho_{su} \mathbf{x}_{ut}^1 + \delta \zeta_{su} \mathbf{x}_{ut}^2 - \zeta_s \mathbf{x}_{su}^1 \mathbf{x}_{ut}^1 \\ &= \rho_{su} \mathbf{x}_{ut}^1 + \delta \zeta_{su} \mathbf{x}_{ut}^2. \end{aligned}$$

Summarizing, we have derived the representation

$$\left[\delta \left(\int \rho \, dx \right) \right]_{sut} = \rho_{su} \mathbf{x}_{ut}^1 + \delta \zeta_{su} \mathbf{x}_{ut}^2.$$

As we are dealing with smooth paths, we have that $\delta(\int \rho \, dx)$ lies in the space \mathcal{ZC}_3^{1+} and thus belongs to the domain of Λ due to Proposition 2.1. (Recall that $\delta\delta = 0$.) Hence, it follows that

$$\int_s^t \rho_{su} \, dx_u = \Lambda_{st}(\rho \mathbf{x}^1 + \delta \zeta \mathbf{x}^2),$$

and inserting this identity into (13), we end up with

$$\int_s^t z_u \, dx_u = z_s \mathbf{x}_{st}^1 + \zeta_s \mathbf{x}_{st}^2 + \Lambda_{st}(\rho \mathbf{x}^1 + \delta \zeta \mathbf{x}^2).$$

Since in addition

$$\rho \mathbf{x}^1 + \delta \zeta \mathbf{x}^2 = -\delta(z \mathbf{x}^1 + \zeta \mathbf{x}^2),$$

we can also write this as

$$\int z_u \, dx_u = (\text{id} - \Lambda\delta)(z \mathbf{x}^1 + \zeta \mathbf{x}^2).$$

Thus we have expressed the Riemann–Stieltjes integral of z with respect to x in terms of the sewing map Λ , the couple $(\mathbf{x}^1, \mathbf{x}^2)$, and of increments of z . This can now be generalized to the nonsmooth case. Note that Corollary 2.2 justifies the use of the notion of integral.

Proposition 2.6 *For fixed $\frac{1}{3} < \kappa \leq \gamma$, let x be a path satisfying Hypothesis 2.5 on an arbitrary interval $[0, T]$. Furthermore, let $z \in \mathcal{Q}_{\kappa, \alpha}^x([\ell_1, \ell_2]; \mathbb{R}^d)$ be such that the increments of z are given by (11). Define \hat{z} by $\hat{z}_{\ell_1} = \hat{\alpha}$ with $\hat{\alpha} \in \mathbb{R}$ and*

$$\delta \hat{z}_{st} = [(\text{id} - \Lambda\delta)(z^i \mathbf{x}^{1,i} + \zeta^{ji} \mathbf{x}^{2,ij})]_{st} \tag{15}$$

for $\ell_1 \leq s \leq t \leq \ell_2$. Then $\mathcal{J}(z^ \, dx) := \hat{z}$ is a well-defined element of $\mathcal{Q}_{\kappa, \hat{\alpha}}^x([\ell_1, \ell_2]; \mathbb{R})$ and coincides with the usual Riemann integral whenever z and x are smooth functions.*

Moreover, the Hölder norm of $\mathcal{J}(z^* \, dx)$ can be estimated in terms of the Hölder norm of the integrator z . (For this and also for a proof of the above proposition, see, e.g., [14].) This allows us to use a fixed-point argument to obtain the existence of a unique solution for rough differential equations.

Theorem 2.7 *For fixed $\frac{1}{3} < \kappa < \gamma$, let x be a path satisfying Hypothesis 2.5 on an arbitrary interval $[0, T]$. Consider a given initial condition a in \mathbb{R}^m and a family (V_1, \dots, V_d) of C^3 vector fields of \mathbb{R}^m , bounded with bounded derivatives. Let $\|f\|_{\mu, \infty} = \|f\|_{\infty} + \|\delta f\|_{\mu}$ be the usual Hölder norm of a path $f \in \mathcal{C}_1([0, T]; \mathbb{R}^l)$. Then we have:*

- (1) Equation (10) admits a unique solution y in $\mathcal{Q}_{\kappa,a}^x([0, T]; \mathbb{R}^m)$ for any $T > 0$, and there exists a polynomial $P_T : \mathbb{R}^2 \rightarrow \mathbb{R}^+$ such that

$$\mathcal{N}[y; \mathcal{Q}_{\kappa,a}^x([0, T]; \mathbb{R}^m)] \leq P_T(\|\mathbf{x}^1\|_\gamma, \|\mathbf{x}^2\|_{2\gamma}). \tag{16}$$

- (2) Let $F : \mathbb{R}^m \times \mathcal{C}_1^\gamma([0, T]; \mathbb{R}^d) \times \mathcal{C}_2^{2\gamma}([0, T]; \mathbb{R}^{m,m}) \rightarrow \mathcal{C}_1^\gamma([0, T]; \mathbb{R}^m)$ be the mapping defined by

$$F(a, \mathbf{x}^1, \mathbf{x}^2) = y,$$

where y is the unique solution of (10). This mapping is locally Lipschitz continuous in the following sense: Let \tilde{x} be another driving rough path with corresponding Lévy area $\tilde{\mathbf{x}}^2$, and \tilde{a} be another initial condition. Moreover denote by \tilde{y} the unique solution of the corresponding differential equation. Then, there exists an increasing function $K_T : \mathbb{R}^4 \rightarrow \mathbb{R}^+$ such that

$$\begin{aligned} \|y - \tilde{y}\|_{\gamma, \infty, T} &\leq K_T(\|\mathbf{x}^1\|_\gamma, \|\tilde{\mathbf{x}}^1\|_\gamma, \|\mathbf{x}^2\|_{2\gamma}, \|\tilde{\mathbf{x}}^2\|_{2\gamma}) \\ &\times (|a - \tilde{a}| + \|\mathbf{x}^1 - \tilde{\mathbf{x}}^1\|_\gamma + \|\mathbf{x}^2 - \tilde{\mathbf{x}}^2\|_{2\gamma}). \end{aligned} \tag{17}$$

The theorem above is borrowed from [12, 14, 21], and we send the reader to these references for more details on the topic.

2.3 Fractional Brownian Motion

We shall recall here how the abstract Theorem 2.7 applies to fractional Brownian motion. We will also give some basic notions on stochastic analysis with respect to fBm, mainly borrowed from [25], which will turn out to be useful in the sequel.

As already mentioned in the introduction, on a finite interval $[0, T]$ and for some fixed $H \in (1/3, 1/2)$, we consider the canonical probability space $(\Omega, \mathcal{F}, \mathbf{P})$ associated with fractional Brownian motion with Hurst parameter H . That is, $\Omega = \mathcal{C}_0([0, T]; \mathbb{R}^d)$ is the Banach space of continuous functions vanishing at 0 equipped with the supremum norm, \mathcal{F} is the Borel sigma-algebra, and \mathbf{P} is the unique probability measure on Ω such that the canonical process $B = \{B_t, t \in [0, T]\}$ is a d -dimensional fractional Brownian motion with Hurst parameter H . Specifically, B has d independent coordinates, each one being a centered Gaussian process with covariance given by (1).

2.3.1 Functional Spaces

Let \mathcal{E} be the set of the space of d -dimensional elementary functions on $[0, T]$:

$$\begin{aligned} \mathcal{E} = \left\{ f = (f_1, \dots, f_d); f_j = \sum_{i=0}^{n_j-1} a_i^j \mathbf{1}_{[t_i^j, t_{i+1}^j)}, \right. \\ \left. 0 = t_0 < t_1^j < \dots < t_{n_j-1}^j < t_{n_j}^j = T, \text{ for } j = 1, \dots, d \right\}. \end{aligned} \tag{18}$$

We call \mathcal{H} the completion of \mathcal{E} with respect to the semiinner product

$$\langle f, g \rangle_{\mathcal{H}} = \sum_{i=1}^d \langle f_i, g_i \rangle_{\mathcal{H}_i}, \quad \text{where } \langle \mathbf{1}_{[0,t]}, \mathbf{1}_{[0,s]} \rangle_{\mathcal{H}_i} := R(s, t), \quad s, t \in [0, T].$$

Then, one constructs an isometry $K_H^* : \mathcal{H} \rightarrow L^2([0, 1]; \mathbb{R}^d)$ such that

$$K_H^*(\mathbf{1}_{[0,t_1]}, \dots, \mathbf{1}_{[0,t_d]}) = (\mathbf{1}_{[0,t_1]} K_H(t_1, \cdot), \dots, \mathbf{1}_{[0,t_d]} K_H(t_d, \cdot)),$$

where the kernel K_H is given by

$$K_H(t, u) = c_H \left[\left(\frac{u}{t} \right)^{\frac{1}{2}-H} (t-u)^{H-\frac{1}{2}} + \left(\frac{1}{2} - H \right) u^{\frac{1}{2}-H} \int_u^t v^{H-\frac{3}{2}} (v-u)^{H-\frac{1}{2}} dv \right] \mathbf{1}_{\{0 < u < t\}}, \quad (19)$$

with a strictly positive constant c_H , whose exact value is irrelevant for our purpose. Notice that this kernel verifies $R_H(t, s) = \int_0^{s \wedge t} K_H(t, r) K_H(s, r) dr$. Moreover, observe that K_H^* can be represented in the following form: for $\varphi = (\varphi_1, \dots, \varphi_d) \in \mathcal{H}$, we have

$$K_H^* \varphi = (K_H^* \varphi^1, \dots, K_H^* \varphi^d),$$

where $[K_H^* \varphi^i]_t = d_H t^{1/2-H} [D_{T^-}^{1/2-H} (u^{-(1/2-H)} \varphi^i)]_t$

for a strictly positive constant d_H . In particular, each \mathcal{H}_i is a fractional integral space of the form $\mathcal{I}_{T^-}^{1/2-H} (L^2([0, T]))$ and $C_1^{1/2-H}([0, T]) \subset \mathcal{H}_i$.

2.3.2 Malliavin Derivatives

Let us start by defining the Wiener integral with respect to B : for any element f in \mathcal{E} whose expression is given as in (18), we define the Wiener integral of f with respect to B as

$$B(f) := \sum_{j=1}^d \sum_{i=0}^{n_j-1} a_i^j (B_{t_{i+1}}^j - B_{t_i}^j).$$

We also denote this integral as $\int_0^T f(t) dB_t$, since it coincides with a pathwise integral with respect to B .

For $\theta : \mathbb{R} \rightarrow \mathbb{R}$ and $j \in \{1, \dots, d\}$, denote by $\theta^{[j]}$ the function with values in \mathbb{R}^d having all the coordinates equal to zero except the j th coordinate that equals to θ . It is readily seen that

$$\mathbf{E}[B(\mathbf{1}_{[0,s]}^{[j]}) B(\mathbf{1}_{[0,t]}^{[k]})] = \delta_{j,k} R_{s,t}.$$

This definition can be extended by linearity and closure to elements of \mathcal{H} , and we obtain the relation

$$\mathbf{E}[B(f) B(g)] = \langle f, g \rangle_{\mathcal{H}},$$

valid for any couple $(f, g) \in \mathcal{H}^2$. In particular, $B(\cdot)$ defines an isometric map from \mathcal{H} into a subspace of $L^2(\Omega)$. It should be pointed out that $(\Omega, \mathcal{H}, \mathbf{P})$ defines an abstract Wiener space, on which chaos decompositions can be settled. We do not develop this aspect of the theory for sake of conciseness, but we will use later the fact that all L^p norms are equivalent on finite chaos.

We can now proceed to the definition of Malliavin derivatives, for which we need an additional notation:

Notation 2.8 For $n, p \geq 1$, a function $f \in C^p(\mathbb{R}^n; \mathbb{R})$ and any tuple $(i_1, \dots, i_p) \in \{1, \dots, d\}^p$, we set $\partial_{i_1 \dots i_p} f$ for $\frac{\partial^p f}{\partial x_{i_1} \dots \partial x_{i_p}}$.

With this notation in hand, let \mathcal{S} be the family of smooth functionals F of the form

$$F = f(B(h_1), \dots, B(h_n)), \quad (20)$$

where $h_1, \dots, h_n \in \mathcal{H}$, $n \geq 1$, and f is a smooth function with polynomial growth, together with all its derivatives. Then, the Malliavin derivative of such a functional F is the \mathcal{H} -valued random variable defined by

$$DF = \sum_{i=1}^n \partial_i f(B(h_1), \dots, B(h_n)) h_i.$$

For all $p > 1$, it is known that the operator D is closable from $L^p(\Omega)$ into $L^p(\Omega; \mathcal{H})$ (see, e.g., [25, Chap. 1]). We will still denote by D the closure of this operator, whose domain is usually denoted by $\mathbb{D}^{1,p}$ and is defined as the completion of \mathcal{S} with respect to the norm

$$\|F\|_{1,p} := (E(|F|^p) + E(\|DF\|_{\mathcal{H}}^p))^{\frac{1}{p}}.$$

It should also be noticed that partial Malliavin derivatives with respect to each component B^j of B will be invoked: they are defined, for a functional F of the form (20) and $j = 1, \dots, d$, as

$$D^j F = \sum_{i=1}^n \partial_i f(B(h_1), \dots, B(h_n)) h_i^{[j]}$$

and then extended by closure arguments again. We refer to [25, Chap. 1] again for the definition of higher derivatives and Sobolev spaces $\mathbb{D}^{k,p}$ for $k > 1$.

2.3.3 Levy Area of fBm

There are many ways to define the Levy area \mathbf{B}^2 associated to fBm, and the reader is referred to [12, Chap. 15] for a complete review of these. The recent paper [11] is however of special interest for us, since it enables a direct definition of \mathbf{B}^2 by Wiener chaos techniques. It can be summarized in the following way:

Proposition 2.9 *Let $1/3 < H < 1/2$ be a fixed Hurst parameter. Then the fBm B belongs almost surely to any space C_1^γ for $\gamma < H$, and it gives rise to an increment $\mathbf{B}^2 \in C_2^{2\gamma}$ which satisfies Hypothesis 2.5. Furthermore, for any $0 \leq s < t \leq T$, \mathbf{B}_{st}^2 is an element of the second chaos associated to B , and*

$$\mathbf{E}[|\mathbf{B}_{st}^2|^p] \leq c_p (t - s)^{2Hp}, \quad p \geq 1.$$

Moreover, the iterated integrals of B can be obtained as limits of Riemann-type integrals. Indeed, for $k \geq 1$ and $0 \leq s < t \leq T$, consider the simplex

$$S_k([s, t]) = \{(u_1, \dots, u_k); s \leq u_1 < \dots < u_k \leq t\}. \tag{21}$$

For a given partition Π of $[0, T]$, we also denote by B^Π the linearization of B based on Π . Combining the results of [11, 12], the following proposition holds:

Proposition 2.10 *Let $k \geq 1$, and for a sequence of partitions $(\Pi_n)_{n \geq 1}$, set $B^n := B^{\Pi_n}$. For $0 \leq s < t \leq T$ and $(i_1, \dots, i_k) \in \{1, \dots, d\}^k$, we consider then*

$$\mathbf{B}_{st}^{k,n,i_1,\dots,i_k} = \int_{S_k([s,t])} dB_{u_1}^{n,i_1} \dots dB_{u_k}^{n,i_k},$$

understood in the Riemann sense. Then there exists a sequence of partitions $(\Pi_n)_{n \geq 1}$ such that $\mathbf{B}^{k,n,i_1,\dots,i_k}$ converges almost surely and in L^2 , as an element of $C_2^{k\gamma}$ for any $\gamma < H$, to an element called $\mathbf{B}^{k,i_1,\dots,i_k}$. When $k = 1$, we obtain the increment δB of our fBm. When $k = 2$, the limit corresponds to the increment \mathbf{B}^2 of Proposition 2.9.

As a corollary of the previous considerations, we have the following:

Proposition 2.11 *Assume that $1/3 < H < 1/2$. Then Theorem 2.7 applies almost surely to the fBm paths, enhanced with the Levy area \mathbf{B}^2 . We are thus able to solve the equation*

$$dy_t = \sum_{i=1}^d V_i(y_t) dB_t^i, \quad t \in [0, T], \quad y_0 = a, \tag{22}$$

under the conditions of Theorem 2.7.

3 A Norris-Type Lemma

Norris’ lemma [24] is one of the basic ingredients in order to obtain the smoothness of densities for solutions to stochastic differential equations under hypoelliptic conditions and was already extended to fBm with Hurst parameter $H > 1/2$ in [3]. We shall extend in the current section this lemma to the rough paths context. A preliminary step along this direction consists of proving the following elementary lemma:

Lemma 3.1 *Let $0 < \alpha < \rho < 1$, and consider $b \in \mathcal{C}([0, T])$. Then for any $0 < \eta < 1$, we have*

$$\|b\|_{\alpha, \infty} \leq C_{\alpha, \rho} [\eta \|b\|_{\rho, \infty} + \eta^{-1/(\rho-\alpha)} \|b\|_{L_1}], \tag{23}$$

where we recall that $\|b\|_{\alpha, \infty}$ has been defined by (6).

Proof Recall that $\|b\|_{\alpha}$ has been defined by (5). Thus, for $0 < \alpha < \rho < 1$, we have

$$\|b\|_{\alpha} = \sup_{s,t} \left[\left(\frac{|\delta b_{st}|}{|t-s|^{\rho}} \right)^{\frac{\alpha}{\rho}} |\delta b_{st}|^{1-\frac{\alpha}{\rho}} \right] \leq 2^{1-\frac{\alpha}{\rho}} \|b\|_{\infty}^{1-\frac{\alpha}{\rho}} \|b\|_{\rho}^{\frac{\alpha}{\rho}}.$$

Thus, for an arbitrary constant $\eta > 0$, we have

$$\|b\|_{\alpha} \leq C_{\alpha, \rho} (\eta \|b\|_{\rho} + \eta^{-\frac{\alpha}{\rho-\alpha}} \|b\|_{\infty}),$$

thanks to Young’s inequality. Invoke now the interpolation inequality [3, formula (3.17)] in order to get

$$\begin{aligned} \|b\|_{\alpha} &\leq C_{\alpha, \rho} (\eta \|b\|_{\rho} + \eta^{-\frac{\alpha}{\rho-\alpha}} [\gamma \|b\|_{\rho} + \gamma^{-\frac{1}{1+\alpha}} \|b\|_{L_1}]) \\ &= C_{\alpha, \rho} [\eta \|b\|_{\rho} + \eta^{-1/(\rho-\alpha)} \|b\|_{L_1}], \end{aligned}$$

where we have chosen $\gamma = \eta^{\frac{\alpha}{\rho-\alpha}}$. Invoking again [3, formula (3.17)] in order to go from $\|\cdot\|_{\alpha}$ to $\|\cdot\|_{\alpha, \infty}$ norms, and this finishes our proof. \square

We can now turn to the announced Norris-type lemma, whose proof is an adaptation of [3] to the case of controlled processes.

Proposition 3.2 *Assume that B is a fractional Brownian motion with $H > 1/3$. Let z be a controlled path in $\mathcal{Q}_{\gamma}^B(\mathbb{R}^m)$, with decomposition*

$$\delta z_{st}^i = \zeta_s^{i,j} \mathbf{B}_{st}^{1,j} + r_{st}^i. \tag{24}$$

We assume that $1/3 < \alpha < \gamma < H$ and that the quantity $\mathbf{E}[\mathcal{N}^p[z; \mathcal{Q}_{\gamma}^B(\mathbb{R}^m)]]$ is finite for all $p \geq 1$. Set $\delta y_{st} = \mathcal{J}(z^ dB)$ according to Proposition 2.6. Then there exists $q > 0$ such that, for every $p > 0$, we can find a strictly positive constant c_p such that*

$$\mathbf{P}(\|y\|_{\gamma, \infty} < \varepsilon, \text{ and } \|z\|_{\alpha, \infty} > \varepsilon^q) < c_p \varepsilon^p.$$

Proof In order to avoid cumbersome indices, we shall prove our result in the case of one-dimensional processes. Generalization to the multidimensional setting is a matter of trivial considerations. We also work on the interval $[0, 1]$ instead of $[0, T]$ for the sake of notational simplicity. As a last preliminary observation, note that if z admits the decomposition (24), then according to (15), we have

$$\delta y = z \mathbf{B}^1 + \zeta \mathbf{B}^2 + y^{\sharp}, \quad \text{where } y^{\sharp} = \Lambda(r \mathbf{B}^1 + \delta \zeta \mathbf{B}^2).$$

Similarly to what is done in [3], we consider two time scales $\delta \ll \Delta \ll 1$. We assume moreover that $\Delta/\delta = r$ with $r \in \mathbb{N}$. We use a partition $\{t_n; n \leq 1/\delta\}$ of $[0, 1]$ with $t_n = n\delta$, so that $t_{Nr} = N\Delta$. Some increments below will then be frozen on the time scale Δ , in order to take advantage of some averaging properties of the process B .

Step 1: Coarse Graining on Increments Consider then n such that $(N - 1)r \leq n \leq Nr - 1$, so that $(N - 1)\Delta \leq t_n \leq N\Delta - \delta$. According to (15), we have

$$\begin{aligned} \delta y_{t_n t_{n+1}} &= z_{t_n} \mathbf{B}^1_{t_n t_{n+1}} + \zeta_{t_n} \mathbf{B}^2_{t_n t_{n+1}} + y^\sharp_{t_n t_{n+1}} \\ &= z_{N\Delta} \mathbf{B}^1_{t_n t_{n+1}} - \delta z_{t_n, N\Delta} \mathbf{B}^1_{t_n t_{n+1}} + \zeta_{t_n} \mathbf{B}^2_{t_n t_{n+1}} + y^\sharp_{t_n t_{n+1}}, \end{aligned}$$

where y^\sharp is an increment in $C^{3\gamma}_2$. Thus,

$$z_{N\Delta} \mathbf{B}^1_{t_n t_{n+1}} = \delta y_{t_n t_{n+1}} + \delta z_{t_n, N\Delta} \mathbf{B}^1_{t_n t_{n+1}} - \zeta_{t_n} \mathbf{B}^2_{t_n t_{n+1}} - y^\sharp_{t_n t_{n+1}}. \tag{25}$$

Set now $X_N = \sum_{n=(N-1)r}^{Nr-1} \|\mathbf{B}^1_{t_n t_{n+1}}\|^4$ and $Y_N = X_N^{1/4}$. Then

$$|z_{N\Delta}|^4 |X_N| = \sum_{n=(N-1)r}^{Nr-1} |z_{N\Delta} \mathbf{B}^1_{t_n t_{n+1}}|^4.$$

Furthermore, invoking relation (25), we get

$$|z_{N\Delta} \mathbf{B}^1_{t_n t_{n+1}}| \leq \|y\|_\gamma \delta^\gamma + \|z\|_\gamma \|B\|_\gamma \delta^\gamma \Delta^\gamma + \|\zeta\|_\infty \|\mathbf{B}^2\|_{2\gamma} \delta^{2\gamma} + \|y^\sharp\|_{3\gamma} \delta^{3\gamma}.$$

Raising this inequality to power 4 and summing over $(N - 1)r \leq n \leq Nr - 1$, we obtain

$$|z_{N\Delta}^4 X_N| \leq \delta^{4\gamma-1} \Delta (\|y\|_\gamma + \|z\|_\gamma \|B\|_\gamma \Delta^\gamma + \|\zeta\|_\infty \|\mathbf{B}^2\|_{2\gamma} \delta^\gamma + \|y^\sharp\|_{3\gamma} \delta^{2\gamma})^4,$$

and therefore

$$|z_{N\Delta}| Y_N \leq \delta^{\gamma-1/4} \Delta^{1/4} (\|y\|_\gamma + \|z\|_\gamma \|B\|_\gamma \Delta^\gamma + \|\zeta\|_\infty \|\mathbf{B}^2\|_{2\gamma} \delta^\gamma + \|y^\sharp\|_{3\gamma} \delta^{2\gamma}).$$

Sum now over N (recall that $1 \leq N \leq 1/\Delta$) in order to get

$$\begin{aligned} &\sum_{N=1}^{1/\Delta} |z_{N\Delta}| Y_N \\ &\leq \delta^{\gamma-1/4} \Delta^{-3/4} (\|y\|_\gamma + \|z\|_\gamma \|B\|_\gamma \Delta^\gamma + \|\zeta\|_\infty \|\mathbf{B}^2\|_{2\gamma} \delta^\gamma + \|y^\sharp\|_{3\gamma} \delta^{2\gamma}). \tag{26} \end{aligned}$$

Step 2: Behavior of a 4th-Order Variation Throughout the proof, we shall use the notations \lesssim, \asymp given in the introduction. For $K \geq 1$, set $\tilde{X}_K = \sum_{n=1}^K |\mathbf{B}^1_{t_n t_{n+1}}|^4$. We shall prove that

$$\mathbf{E}[\tilde{X}_K] \asymp K \delta^{4H}, \quad \text{and} \quad \mathbf{Var}(\tilde{X}_K) \lesssim K \delta^{8H}. \tag{27}$$

Indeed, a simple scaling argument shows that $\tilde{X}_K \stackrel{(\mathcal{L})}{=} \delta^{4H} \hat{X}_K$ with $\hat{X}_K = \sum_{n=1}^K |\mathbf{B}_{n,n+1}^1|^4$. Introduce now the k th Hermite polynomial H_k (see [25] for a definition and properties of these objects) and notice in particular that $H_2(x) = x^2 - 1$ and $H_4(x) = x^4 - 6x^2 + 3$. This enables to decompose \hat{X}_K as

$$\hat{X}_K = \sum_{n=1}^K [H_4(\mathbf{B}_{n,n+1}^1) + 6H_2(\mathbf{B}_{n,n+1}^1)] + 3K. \tag{28}$$

Recall now that for a centered Gaussian vector (U, V) in \mathbb{R}^2 such that $\mathbf{E}[U^2] = \mathbf{E}[V^2] = 1$, we have

$$\mathbf{E}[H_k(U)] = 0, \quad \text{and} \quad \mathbf{E}[H_k(U) H_l(V)] = k! (\mathbf{E}[U V])^k \mathbf{1}_{(k=l)}. \tag{29}$$

Plugging this identity into (28), this immediately yields $\mathbf{E}[\hat{X}_K] = 3K$, which is our first assertion in (27). In addition, the second part of (29) entails

$$\mathbf{Var}(\hat{X}_K) = 2 \sum_{n_1, n_2=1}^K (12\alpha_{n_1, n_2}^4 + \alpha_{n_1, n_2}^2) := 2S_K,$$

where we have set

$$\begin{aligned} \alpha_{n_1, n_2} &= \mathbf{E}[\mathbf{B}_{n_1, n_1+1}^1 \mathbf{B}_{n_2, n_2+1}^1] \\ &= \frac{1}{2} [|n_2 - n_1 + 1|^{2H} + |n_2 - n_1 - 1|^{2H} - 2|n_2 - n_1|^{2H}]. \end{aligned}$$

Summarizing, we have obtained that

$$\mathbf{Var}(\tilde{X}_K) = \delta^{8H} \mathbf{Var}(\hat{X}_K) = 2\delta^{8H} S_K. \tag{30}$$

We will now prove that $S_K \lesssim K$. Indeed, write first

$$S_K = \sum_{1 \leq n_1 \leq K} (12\alpha_{n_1, n_1}^4 + \alpha_{n_1, n_1}^2) + 2 \sum_{1 \leq n_1 < n_2 \leq K} (12\alpha_{n_1, n_2}^4 + \alpha_{n_1, n_2}^2) := S_K^1 + 2S_K^2.$$

Then, since $\alpha_{n_1, n_1} = 1$, it is readily checked that $S_K^1 \lesssim K$. Moreover, the term S_K^2 can be decomposed into

$$\begin{aligned} S_K^2 &= \sum_{1 \leq n_1 \leq K-1} (12\alpha_{n_1, n_1+1}^4 + \alpha_{n_1, n_1+1}^2) \\ &+ \sum_{1 \leq n_1, n_2 \leq K, n_2 - n_1 \geq 2} (12\alpha_{n_1, n_2}^4 + \alpha_{n_1, n_2}^2) := S_K^{21} + S_K^{22}. \end{aligned}$$

Notice now that $\alpha_{n_1, n_1+1} = -[1 - 2^{-2(H-1/2)}]$, which easily yields $S_K^{21} \leq c_H K$.

As far as S_K^{22} is concerned, write, for $n_2 - n_1 \geq 2$,

$$\alpha_{n_1, n_2} = H(2H - 1) \int_0^1 dr \int_{-r}^r |(n_2 - n_1) + u|^{2H-2} du,$$

which immediately yields $\alpha_{n_1, n_2} \lesssim |(n_2 - n_1) - 1|^{2H-2}$. Thus,

$$\begin{aligned} S_K^{22} &\lesssim \sum_{n_1=1}^{K-2} \sum_{n_2=n_1+2}^K |(n_2 - n_1) - 1|^{8H-8} + 3|(n_2 - n_1) - 1|^{4H-4} \\ &\lesssim \sum_{n_1=1}^{K-2} \sum_{m=1}^{K-n_1+1} m^{4H-4} \leq K \sum_{m=1}^{\infty} m^{4H-4} \lesssim K, \end{aligned}$$

since $\sum_{m=1}^{\infty} m^{4H-4}$ is finite whenever $H < 3/4$.

Gathering our bounds on S_K^1 , S_K^2 , and S_K^{22} , we obtain $S_K \lesssim K$, and plugging this bound into (30), we end up with $\mathbf{Var}(\tilde{X}_K) \lesssim K \delta^{8H}$, which is our claim. This finishes the proof of (27).

Step 3: Concentration Inequalities for Y_N Let us recall that X_N is in the 4th chaos of the fBm B . Hence, a result by Borell [5] entails

$$\mathbf{P}\left(\frac{|X_N - \mathbf{E}[X_N]|}{[\mathbf{Var}(X_N)]^{1/2}} \geq u\right) \leq c_1 e^{-c_2 u^{1/2}}, \quad u \geq 0,$$

for two universal constant $c_1, c_2 > 0$. With (27) in hand, this yields

$$\mathbf{P}\left(|X_N - 3\Delta\delta^{4H-1}| \geq \Delta^{1/2}\delta^{4H-1/2}u\right) \leq c_1 e^{-c_2 u^{1/2}}, \quad u \geq 0. \tag{31}$$

We now wish to produce a concentration inequality for $Y_N = X_N^{1/4}$. Since X_N is a small random quantity of order $\Delta\delta^{4H-1}$, let us use the inequality

$$|b^{1/4} - a^{1/4}| \lesssim \xi^{-3/4}|b - a|, \quad \text{where } \xi \in (a \wedge b, a \vee b).$$

Apply this with $\xi = \frac{3}{2}\Delta\delta^{4H-1}$ in order to get

$$\mathbf{P}\left(|Y_N - 3^{1/4}\Delta^{1/4}\delta^{H-1/4}| \geq c\Delta^{-3/4}\delta^{-3(4H-1)/4}\Delta^{1/2}\delta^{4H-1/2}u\right) \leq A_1 + A_2 \tag{32}$$

with

$$\begin{aligned} A_1 &= \mathbf{P}\left(|X_N - 3\Delta\delta^{4H-1}| \geq c\Delta^{1/2}\delta^{4H-1/2}u, X_N \geq \frac{3}{2}\Delta\delta^{4H-1}\right), \\ A_2 &= \mathbf{P}\left(X_N \leq \frac{3}{2}\Delta\delta^{4H-1}\right). \end{aligned}$$

Furthermore, a straightforward application of (31) gives

$$A_1 \leq c_1 e^{-c_2 u^{1/2}} \quad \text{and} \quad A_2 \leq c_1 e^{-c_2(\Delta/\delta)^{1/4}}.$$

Plugging these inequalities into (32), we end up with the following concentration inequality for Y_N :

$$\mathbf{P}\left(|Y_N - 3^{1/4}\Delta^{1/4}\delta^{H-1/4}| \geq c\Delta^{-1/4}\delta^{H+1/4}u\right) \leq c_1 e^{-c_2 u^{1/2}} + c_1 e^{-c_2(\Delta/\delta)^{1/4}}. \tag{33}$$

We shall thus retain the fact that Y_N is a random quantity of order $3^{1/4} \Delta^{1/4} \delta^{H-1/4}$, with fluctuations of order $\Delta^{-1/4} \delta^{H+1/4}$:

$$|Y_N - 3^{1/4} \Delta^{1/4} \delta^{H-1/4}| \approx \delta^{H+1/4} \Delta^{-1/4}. \tag{34}$$

Step 4: Use of the Interpolation Inequality Start again from (26). One would like to have an approximation of the L^1 norm of z appearing on the left-hand side of this inequality, that is, one would like to replace Y_N by Δ . To this purpose, replace first Y_N by its approximation $\Delta^{1/4} \delta^{H-1/4}$ from the last step. This yields an inequality of the form

$$\begin{aligned} \Delta^{1/4} \delta^{H-1/4} \sum_{N=1}^{1/\Delta} |z_{N\Delta}| &\leq \|z\|_\infty \sum_{N=1}^{1/\Delta} |Y_N - 3^{1/4} \Delta^{1/4} \delta^{H-1/4}| \\ &\quad + \delta^{\gamma-1/4} \Delta^{-3/4} (\|y\|_\gamma + \|z\|_\gamma \|B\|_\gamma \Delta^\gamma \\ &\quad + \|\zeta\|_\infty \|B^2\|_{2\gamma} \delta^\gamma + \|y^\sharp\|_{3\gamma} \delta^{2\gamma}). \end{aligned}$$

Rescale this inequality in order to get Δ multiplying on the left-hand side, which gives:

$$\begin{aligned} \Delta \sum_{N=1}^{1/\Delta} |z_{N\Delta}| &\leq \|z\|_\infty R_{\delta,\Delta} + \delta^{-(H-\gamma)} (\|y\|_\gamma + \|z\|_\gamma \|B\|_\gamma \Delta^\gamma \\ &\quad + \|\zeta\|_\infty \|B^2\|_{2\gamma} \delta^\gamma + \|y^\sharp\|_{3\gamma} \delta^{2\gamma}), \end{aligned} \tag{35}$$

where

$$R_{\delta,\Delta} := \delta^{-(H-1/4)} \Delta^{3/4} \sum_{N=1}^{1/\Delta} |Y_N - 3^{1/4} \Delta^{1/4} \delta^{H-1/4}|. \tag{36}$$

Furthermore, it is well known that $|\Delta \sum_{N=1}^{1/\Delta} |z_{N\Delta}| - \|z\|_{L^1}| \leq \|z\|_\gamma \Delta^\gamma$, so that we can recast (35) into

$$\begin{aligned} \|z\|_{L^1} &\leq \|z\|_\gamma \Delta^\gamma + \|z\|_\infty R_{\delta,\Delta} \\ &\quad + \delta^{-(H-\gamma)} (\|y\|_\gamma + \|z\|_\gamma \|B\|_\gamma \Delta^\gamma + \|\zeta\|_\infty \|B^2\|_{2\gamma} \delta^\gamma + \|y^\sharp\|_{3\gamma} \delta^{2\gamma}). \end{aligned} \tag{37}$$

Recall that we take $0 < \alpha < \gamma < H$ and set $\nu_H := 1/(\gamma - \alpha)$. According to (23), we have

$$\|z\|_{\alpha,\infty} \lesssim \eta \|z\|_{\gamma,\infty} + \eta^{-\nu_H} \|z\|_{L^1}$$

for any (small enough) constant η , and plugging (37) into this last relation, we obtain

$$\begin{aligned} \|z\|_{\alpha,\infty} &\lesssim \eta^{-\nu_H} \Delta^\gamma \|z\|_{\gamma,\infty} + \eta^{-\nu_H} \|z\|_\infty R_{\delta,\Delta} + \eta \|z\|_{\gamma,\infty} \\ &\quad + \eta^{-\nu_H} (\|y\|_{\gamma,\infty} \delta^{-(H-\gamma)} + \|z\|_{\gamma,\infty} \|B\|_\gamma \delta^{-(H-\gamma)} \Delta^\gamma) \end{aligned}$$

$$+ \|\zeta\|_\infty \left\| \mathbf{B}^2 \right\|_{2\gamma} \delta^{2\gamma-H} + \|y^\sharp\|_{3\gamma} \delta^{3\gamma-H}.$$

Defining $\hat{\epsilon}$ as $\hat{\epsilon} = H - \gamma$, we get

$$\begin{aligned} \|z\|_{\alpha,\infty} &\lesssim \eta^{-\nu H} \Delta^\gamma \|z\|_{\gamma,\infty} + \eta^{-\nu H} \|z\|_\infty R_{\delta,\Delta} + \eta \|z\|_{\gamma,\infty} \\ &\quad + \eta^{-\nu H} (\|y\|_{\gamma,\infty} \delta^{-\hat{\epsilon}} + \|z\|_{\gamma,\infty} \|B\|_\gamma \delta^{-\hat{\epsilon}} \Delta^\gamma \\ &\quad + \|\zeta\|_\infty \left\| \mathbf{B}^2 \right\|_{2\gamma} \delta^{\gamma-\hat{\epsilon}} + \|y^\sharp\|_{3\gamma} \delta^{2\gamma-\hat{\epsilon}}). \end{aligned} \tag{38}$$

Step 5: Tuning the Parameters Recall that we have chosen $0 \ll \delta \ll \Delta \ll 1$. We express this fact in terms of powers of ϵ by taking $\delta = \epsilon^\mu$ and $\Delta = \epsilon^\lambda$ with $\mu > \lambda > 0$. We also choose η of the form $\eta = \epsilon^{\tau/\nu H}$. We shall now see how to choose λ, μ, τ conveniently: write (38) as

$$\begin{aligned} \|z\|_{\alpha,\infty} &\lesssim \epsilon^{-\tau+\lambda\gamma} \|z\|_{\gamma,\infty} + \epsilon^{-\tau} \|z\|_\infty R_{\delta,\Delta} + \epsilon^{\tau/\nu H} \|z\|_{\gamma,\infty} \\ &\quad + \|y\|_{\gamma,\infty} \epsilon^{-\tau-\hat{\epsilon}\mu} + \|z\|_{\gamma,\infty} \|B\|_\gamma \epsilon^{-\tau-\hat{\epsilon}\mu+\lambda\gamma} \\ &\quad + \|\zeta\|_\infty \left\| \mathbf{B}^2 \right\|_{2\gamma} \epsilon^{-\tau+\mu(\gamma-\hat{\epsilon})} + \|y^\sharp\|_{3\gamma} \epsilon^{-\tau+\mu(2\gamma-\hat{\epsilon})}. \end{aligned} \tag{39}$$

In order to be able to bound z when y is assumed to satisfy $\|y\|_{\gamma,\infty} \leq \epsilon$, the coefficients in the right-hand side of (39) should fulfill the following conditions:

- The coefficient in front of $\|y\|_{\gamma,\infty}$ should be smaller than ϵ^{-1} .
- The other coefficients should be $\ll 1$.

Looking at the exponents in (39), assuming that $\hat{\epsilon}$ is arbitrarily small, and letting for the moment $R_{\delta,\Delta}$ apart, this imposes the following relations:

$$\lambda\gamma > \tau \quad \text{and} \quad 0 < \tau < 1. \tag{40}$$

Let us go back now to the evaluation of $R_{\delta,\Delta}$, given by expression (36), with the order of magnitude of $|Y_N - \Delta^{1/4} \delta^{H-1/4}|$ given by (34). Therefore,

$$R_{\delta,\Delta} \approx \delta^{-(H-1/4)} \Delta^{3/4} \Delta^{-1} \delta^{H+1/8} \Delta^{-1/4} = \delta^{1/2} \Delta^{-1/2}.$$

Expressing this in terms of powers of ϵ , we end up with

$$\eta^{-\nu H} R_{\delta,\Delta} \approx \epsilon^\kappa \quad \text{with} \quad \kappa = \frac{\mu - \lambda}{2} - \tau.$$

If we wish this remainder term to be small, this adds the condition

$$\mu - \lambda > 2\tau, \tag{41}$$

which can be fulfilled easily. From now on, we shall assume that both (40) and (41) are satisfied.

Step 6: Conclusion Recall that we wish to study the probability $\mathbf{P}(\|y\|_{\gamma,\infty} < \varepsilon, \text{ and } \|z\|_{\alpha,\infty} > \varepsilon^q)$. This quantity is obviously bounded by $B_1 + B_2$, where

$$B_1 = \mathbf{P}(\|y\|_{\gamma,\infty} < \varepsilon, \|z\|_{\alpha,\infty} > \varepsilon^q, |R_{\delta,\Delta}| \leq c \Delta^{-1/4} \delta^{H+1/4} (\Delta/\delta)^{\tilde{\varepsilon}}),$$

$$B_2 = \mathbf{P}(|R_{\delta,\Delta}| \geq c \Delta^{-1/4} \delta^{H+1/4} (\Delta/\delta)^{\tilde{\varepsilon}}),$$

where $\tilde{\varepsilon}$ is an arbitrary small positive constant. Furthermore, inequalities (33) and (36) yield, for any $p \geq 1$,

$$B_2 \leq c_1 e^{-c_2(\Delta/\delta)^{\tilde{\varepsilon}/2}} + c_1 e^{-c_2(\Delta/\delta)^{1/4}} \leq c_{p,\lambda,\mu,\tilde{\varepsilon}} \varepsilon^p,$$

where we have used the fact that $\delta/\Delta = e^{\mu-\lambda}$.

We can now bound B_1 : notice that according to (39), if we are working on

$$(|R_{\delta,\Delta}| \leq c \Delta^{-1/4} \delta^{H+1/4} (\Delta/\delta)^{\tilde{\varepsilon}}) \cap (\|y\|_{\gamma,\infty} < \varepsilon),$$

then there exists a $\rho > 0$ such that

$$\|z\|_{\alpha,\infty} \lesssim \varepsilon^\rho [1 + \mathcal{N}^2[z; \mathcal{Q}_\gamma^B(\mathbb{R}^m)] + \|B\|_\gamma^2 + \|\mathbf{B}^2\|_{2\gamma}^2].$$

Moreover, recall that $\mathcal{N}[z; \mathcal{Q}_\gamma^B(\mathbb{R}^m)]$ is assumed to be an L^r random variable for all $r \geq 1$, while $\|B\|_\gamma$ and $\|\mathbf{B}^2\|_{2\gamma}$ are also elements of L^r , since they can be bounded by a finite chaos random variable. Thus, the Chebyshev inequality can be applied here, which entails

$$B_1 \lesssim (1 + \mathbf{E}[\mathcal{N}^{2l}[z; \mathcal{Q}_\gamma^B(\mathbb{R}^m)] + \|B\|_\gamma^{2l} + \|\mathbf{B}^2\|_{2\gamma}^{2l}]) \varepsilon^{l(\rho-q)}$$

for an arbitrary $l \geq 1$. It is now sufficient to choose $q < \rho$ and l large enough so that $l(\rho - q) = p$ to conclude the proof by putting together our bounds on B_1 and B_2 . □

4 Malliavin Calculus for Solutions to Fractional SDEs

This section is the core of our paper, where we derive the smoothness of density for the solution to (22). We first recall some classical notions on representations of solutions to SDEs and then move to Malliavin calculus considerations.

4.1 Representation of Solutions to SDEs

The first representation results for solutions to SDEs in terms of the driving vector fields can be traced back to the seminal work of Chen [7]. They have then been deeply analyzed in [18, 29] and also lie at the basis of the rough path theory [21]. We have chosen here to present these formulas according to [1], which is a recent and didactically useful account on the topic.

Recall that we are considering a d -dimensional fBm B with $1/3 < H < 1/2$. According to Sect. 2.3, this allows us to construct some increments \mathbf{B}^k out of B which

can be seen as limits of Riemann iterated integrals over the simplex $\mathcal{S}_k([s, t])$, as recalled in Proposition 2.10. Furthermore, one can solve (22) under the conditions of Theorem 2.7.

Let us introduce some additional notation: let \mathcal{V} be the space of smooth bounded vector fields over \mathbb{R}^m . Given $V \in \mathcal{V}$, we use $\exp(V)$ to denote the exponential map, which maps a vector to another vector and is defined by the relation $[\exp(V)](\xi) = \Psi_1(\xi)$, where $\{\Psi_t(\xi); t \geq 0\}$ is a solution to the ordinary differential equation

$$\partial_t \Psi_t(\xi) = V(\Psi_t(\xi)), \quad \Psi_0(\xi) = \xi. \tag{42}$$

The aim of the Chen–Strichartz formula is to express the solution y_t to (22), for an arbitrary $t \in [0, T]$, as $y_t = [\exp(Z_t)](a)$ for a certain $Z_t \in \mathcal{V}$.

To this purpose, let us give some more classical notations on vector fields: if $V, W \in \mathcal{V}$, then the vector field $[V, W] \in \mathcal{V}$ (called Lie bracket of V and W) is defined by

$$[V, W]^i = V^l \partial_{x_l} W^i - W^l \partial_{x_l} V^i.$$

Notice that this notion is usually introduced through the interpretation of \mathcal{V} as a set of first-order differential operators. A Lie bracket of order k can also be defined inductively for $k \geq 2$ by setting

$$[U_1 \cdots U_k] = [[U_1 \cdots U_{k-1}], U_k]$$

for $U_1 \dots U_k \in \mathcal{V}$. With this notation in hand, our main assumption on the vector fields V_1, \dots, V_d governing (22) is the following:

Hypothesis 4.1 *The vector fields V_1, \dots, V_d are n -nilpotent for some given positive integer n . Namely, for any $(i_1, \dots, i_n) \in \{1, \dots, d\}^n$, we have $[V_{i_1} \cdots V_{i_n}] = 0$.*

We are now ready to state our formulation of Strichartz’ identity, for which we need two last notations: for $k \geq 1$, we call \mathfrak{S}_k the set of permutations of $\{1, \dots, k\}$. Moreover, for $\sigma \in \mathfrak{S}_k$, write $e(\sigma)$ for the quantity $\text{Card}(\{j \in \{1, \dots, k-1\}; \sigma(j) > \sigma(j+1)\})$. Then the following formula is proven, e.g., in [1, 18, 29]:

Proposition 4.2 *Under the hypothesis of Theorem 2.7, let y be the solution to (22). Assume Hypothesis 4.1 holds true, and consider $t \in [0, T]$. Then $y_t = [\exp(Z_t)](a)$, where Z_t can be expressed as follows:*

$$Z_t = \sum_{k=1}^{n-1} \sum_{i_1, \dots, i_k=1}^d \mathbf{V}_{i_1, \dots, i_k} \psi_t^{i_1, \dots, i_k}, \quad \text{with } \psi_t^{i_1, \dots, i_k} = \sum_{\sigma \in \mathfrak{S}_k} \frac{(-1)^{e(\sigma)}}{k^2 \binom{k-1}{e(\sigma)}} \mathbf{B}_{0t}^{\mathbf{k}, i_{\tau(1)}, \dots, i_{\tau(k)}},$$

where we have set $\tau = \sigma^{-1}$ and $\mathbf{V}_{i_1, \dots, i_k} = [V_{i_1} \cdots V_{i_k}]$ in the formula above.

As a warmup to the computations below, we prove now that one can extend our inequality (16) thanks to the Strichartz representation, covering the case of unbounded vector fields with bounded derivatives:

Proposition 4.3 *Suppose that Hypothesis 4.1 holds and that the smooth vector fields $V_i, i = 1, 2, \dots, d$, have bounded derivative. Assume moreover that all the Lie brackets of order greater than or equal to 2 are bounded vector fields. Then the solution y_t of (22) admits moments of any order. Namely, for any $m > 1$ and any $T \in (0, \infty)$,*

$$\mathbf{E}\left[\sup_{0 \leq t \leq T} |y_t|^m\right] < \infty. \tag{43}$$

Proof One can restate Proposition 4.2 as follows: for any $t \leq T$, the random variable y_t can be expressed as $y_t = \phi_t^1$, where $\phi_s^i := \phi_s : \mathbb{R}^d \rightarrow \mathbb{R}^d$ satisfies (for t fixed)

$$\partial_s \phi_s = \sum_{k=1}^{n-1} \sum_{i_1, \dots, i_k=1}^d \psi_t^{i_1, \dots, i_k} \mathbf{V}_{i_1, \dots, i_k}(\phi_s), \quad 0 \leq s \leq 1, \quad \phi_0 = a.$$

Let us separate the first-order integrals in this equation, in order to get

$$\partial_s \phi_s = \sum_{i=1}^d V_i(\phi_s) B_t^i + \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d \psi_t^{i_1, \dots, i_k} \mathbf{V}_{i_1, \dots, i_k}(\phi_s), \quad \phi_0 = a. \tag{44}$$

Since $V_i, i = 1, 2, \dots, d$, have bounded derivatives and since all the Lie brackets of order greater than or equal to 2 are bounded, we see that

$$\begin{aligned} |\partial_s \phi_s| &\leq \sum_{i=1}^d |V_i(\phi_s)| |B_t^i| + \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d |\psi_t^{i_1, \dots, i_k}| |\mathbf{V}_{i_1, \dots, i_k}(\phi_s)| \\ &\leq c_1 |\phi_s| \sum_{i=1}^d \sup_{0 \leq t \leq T} |B_t^i| + c_2 \sum_{k=1}^{n-1} \sum_{i_1, \dots, i_k=1}^d \sup_{0 \leq t \leq T} |\psi_t^{i_1, \dots, i_k}|. \end{aligned}$$

Thus by Gronwall’s lemma, we have

$$|\phi_s| \leq c_2 \left(\sum_{k=1}^{n-1} \sum_{i_1, \dots, i_k=1}^d \sup_{0 \leq t \leq T} |\psi_t^{i_1, \dots, i_k}| \right) \exp \left\{ c_1 \sum_{i=1}^d \sup_{0 \leq t \leq T} |B_t^i| \right\}.$$

This inequality holds for all $0 \leq s \leq 1$. Thus,

$$\sup_{0 \leq t \leq T} |y_t| \leq c_2 \left(\sum_{k=1}^{n-1} \sum_{i_1, \dots, i_k=1}^d \sup_{0 \leq t \leq T} |\psi_t^{i_1, \dots, i_k}| \right) \exp \left\{ c_1 \sum_{i=1}^d \sup_{0 \leq t \leq T} |B_t^i| \right\},$$

which implies (43) by the theorem of Fernique [10]. □

4.2 Malliavin Derivative

This subsection is devoted to enhancing our Proposition 4.3, and proving that the Malliavin derivative of y_t has also bounded moments of any order in our particular

nilpotent situation. Notice once again that the boundedness of moments of the Malliavin derivative is still an open problem for rough differential equations in the general case. We refer to Sect. 2.3.2 for notations on Malliavin calculus.

Theorem 4.4 *Let the vector fields $V_i, 1 = 1, 2, \dots, d$, be smooth with all derivatives bounded, satisfying Hypothesis 4.1. Assume that all the Lie brackets of order greater than or equal to 2 are constants. Then the Malliavin derivative of y_t has moments of any order. More precisely, for any $q > 1$ and $T \in (0, \infty)$,*

$$\mathbf{E}\left[\sup_{0 \leq u \leq t \leq T} |D_u y_t|^q\right] < \infty. \tag{45}$$

Proof Go back to our representation (44), which can easily be differentiated in the Malliavin calculus sense in order to obtain

$$\begin{aligned} \partial_s D_u \phi_s &= \sum_{i=1}^d \nabla V_i(\phi_s) B_t^i D_u \phi_s + \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d \psi_t^{i_1, \dots, i_k} \nabla \mathbf{V}_{i_1, \dots, i_k}(\phi_s) D_u \phi_s \\ &+ \sum_{i=1}^d V_i(\phi_s) \mathbf{1}_{[0,t]}^{[i]}(u) + \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d D_u \psi_t^{i_1, \dots, i_k} \mathbf{V}_{i_1, \dots, i_k}(\phi_s), \end{aligned}$$

where we have set $\nabla \mathbf{V}_{i_1, \dots, i_k}$ for the (matrix-valued) gradient of $\mathbf{V}_{i_1, \dots, i_k}$, and where we recall that the notation $\mathbf{1}_{[0,t]}^{[i]}$ has been introduced in Sect. 2.3.2.

Since we assume that all the Lie brackets of order greater than or equal to 2 of the vector fields V_i are constant vector fields, it is easily checked that

$$\begin{aligned} \partial_s D_u \phi_s &= \sum_{i=1}^d \nabla V_i(\phi_s) B_t^i D_u \phi_s + \sum_{i=1}^d V_i(\phi_s) \mathbf{1}_{[0,t]}^{[i]}(u) \\ &+ \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d D_u \psi_t^{i_1, \dots, i_k} \mathbf{V}_{i_1, \dots, i_k}(\phi_s). \end{aligned} \tag{46}$$

Therefore there exist two positive constants c_1, c_2 such that

$$\begin{aligned} |\partial_s D_u \phi_s| &\leq c_1 \sum_{i=1}^d |B_t^i| |D_u \phi_s| \\ &+ c_2 \sum_{i=1}^d [1 + |\phi_s|] \mathbf{1}_{[0,t]}^{[i]}(u) + \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d |D_u \psi_t^{i_1, \dots, i_k}| |\mathbf{V}_{i_1, \dots, i_k}(\phi_s)|. \end{aligned}$$

By Gronwall’s lemma we obtain

$$\sup_{0 \leq s \leq 1, 0 \leq u \leq t \leq T} |D_u \phi_s|$$

$$\begin{aligned} &\leq c_2 \exp \left\{ c_1 \sum_{i=1}^d \sup_{0 \leq t \leq T} |B_t^i| \right\} \times \left\{ \sum_{i=1}^d [1 + \sup_{0 \leq s \leq 1} |\phi_s|] \right. \\ &\quad \left. \times \left(1 + \sup_{0 \leq t \leq T} \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d \sup_{0 \leq s \leq 1, 0 \leq u \leq t \leq T} |D_u \psi_t^{i_1, \dots, i_k}| |\mathbf{V}_{i_1, \dots, i_k}(\phi_s)| \right) \right\}. \end{aligned}$$

Thus,

$$\begin{aligned} &\sup_{0 \leq u \leq t \leq T} |D_u y_t| \\ &\leq c_2 \exp \left\{ c_1 \sum_{i=1}^d \sup_{0 \leq t \leq T} |B_t^i| \right\} \\ &\quad \times \left\{ \sum_{i=1}^d [1 + \sup_{0 \leq t \leq T} |y_t|] \left(1 + \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d \sup_{0 \leq s \leq 1, 0 \leq u \leq t \leq T} |D_u \psi_t^{i_1, \dots, i_k}| \right) \right\}, \end{aligned}$$

which ends the proof easily by the boundedness of moments for y_t , $D_u \psi_t^{i_1, \dots, i_k}$ and B_t^i . □

Example 4.5 A classical example of nilpotent vector fields in \mathbb{R}^3 is due to Yamato [31]. Let us check that this example fulfils our standing assumptions. Indeed, the example provided in [31] is the following:

$$A_1 = 0, \quad A_2 = \frac{\partial}{\partial x_1} + 2x_2 \frac{\partial}{\partial x_3}, \quad \text{and} \quad A_3 = \frac{\partial}{\partial x_2} - 2x_1 \frac{\partial}{\partial x_3}.$$

Then

$$[A_2, A_3] = -4 \frac{\partial}{\partial x_3}, \quad [[A_2, A_3], A_2] = [[A_2, A_3], A_3] = 0.$$

It is thus readily checked that the conditions of Theorem 4.4 are met for these vector fields. Moreover, in this particular case the solution to (22) is explicit, and we have

$$y_1^1 = y_1 + B_t^2, \quad y_t^2 = y_2 + B_t^3, \quad y_t^3 = y_3 + 2(\mathbf{B}_{0t}^{2,32} - \mathbf{B}_{0t}^{2,23})$$

if the solution starts from the initial condition (y_1, y_2, y_3) . Interestingly enough, though the solution is explicit here, the smoothness of the density of y_t is not immediate, and we recover here the results of [9].

4.3 Stochastic Flows

The probabilistic proof of the smoothness of density for diffusion processes originally given by Malliavin [22] heavily relies on stochastic flows methods and their relationship with stochastic derivatives. We now establish those relations for SDEs driven by a fractional Brownian motion.

To this aim, denote by $y^{s,a}$ the solution to (22) starting from the initial condition $y_s = a$ at time s :

$$dy_t^{s,a} = \sum_{i=1}^d V_i(y_t^{s,a}) dB_t^i, \quad t \in [s, T], \quad y_s^{s,a} = a. \tag{47}$$

The above equation gives rise to a family of smooth nonlinear mappings $\Phi_{s,t} : \mathbb{R}^m \rightarrow \mathbb{R}^m$, $0 \leq s \leq t \leq T$, determined by $\Phi_{s,t}(a) := y_t^{s,a}$, and the family $\{\Phi_{s,t}; 0 \leq s \leq t \leq T\}$ has the following flow property (we refer, e.g., to [12] for the properties of flows driven by rough paths quoted below):

$$\Phi_{s,t} = \Phi_{u,t} \circ \Phi_{s,u}, \quad 0 \leq s \leq u \leq t \leq T.$$

Let $J_{s,t}$ denote the gradient of the nonlinear mapping $\Phi_{s,t}$ with respect to the initial condition. Then the family $\{J_{s,t}; s \leq t \leq T\}$ also satisfies the relation $J_{s,t} = J_{u,t} J_{s,u}$ for $0 \leq s \leq u \leq t \leq T$. In addition, the map $J_{s,t}$ is invertible, and we have $J_{s,t} = J_{0,t} J_{0,s}^{-1}$. The equation followed by $J_{0,t}$ is obtained by differentiating formally (47) with respect to the initial value a , which yields

$$dJ_{0,t} = \sum_{i=1}^d \nabla V_i(y_t) J_{0,t} dB_t^i, \quad J_{0,0} = I.$$

By applying the rules of differential calculus for rough paths, we also get that $J_{0,t}^{-1}$ is solution to the following equation:

$$dJ_{0,t}^{-1} = - \sum_{i=1}^d \nabla V_i(y_t) J_{0,t}^{-1} dB_t^i, \quad J_{0,0}^{-1} = I. \tag{48}$$

We have thus ended up with two linear equations for the derivatives of the flow. In our nilpotent case, we are thus able to bound these derivatives along the same lines as for Theorem 4.4:

Theorem 4.6 *Let the vector fields V_i , $1 = 1, 2, \dots, d$, be smooth with all derivatives bounded and satisfy Hypothesis 4.1. Assume that all the Lie brackets of order greater than or equal to 2 are constant. Then the Jacobian $J_{0,t}$ and its inverse $J_{0,t}^{-1}$ have moments of any order: for any $q \geq 1$ and $T \in (0, \infty)$,*

$$\mathbf{E} \left[\sup_{0 \leq t \leq T} |J_{0,t}|^q \right] < \infty, \quad \text{and} \quad \mathbf{E} \left[\sup_{0 \leq t \leq T} |J_{0,t}^{-1}|^q \right] < \infty. \tag{49}$$

Proof As mentioned in the proof of Proposition 4.3, one can write $\Phi_{0,t}(a) = \exp(Z_t)(a) = \phi_1(a)$, where $\phi_1(a)$ satisfies (44). Thus, if we introduce $\tilde{J}_s = \nabla \phi_s$, then $J_{0,t} = \tilde{J}_1$ where \tilde{J}_s satisfies

$$\partial_s \tilde{J}_s = \sum_{i=1}^d B_t^i \nabla V_i(\phi_s) \tilde{J}_s + \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d \psi_t^{i_1, \dots, i_k} \nabla \mathbf{V}_{i_1, \dots, i_k}(\phi_s) \tilde{J}_s, \quad \tilde{J}_0 = I.$$

The first part of (49) is thus proved following the steps of Proposition 4.3.

As far as the second part of (49) is concerned, observe that $J_{0,t}^{-1} = \bar{J}_1$, where \bar{J}_s is inverse of \tilde{J}_s . It is clear that \bar{J}_s satisfies

$$\partial_s \bar{J}_s = - \sum_{i=1}^d \nabla V_i(\phi_s) \bar{J}_s B_t^i - \sum_{k=2}^{n-1} \sum_{i_1, \dots, i_k=1}^d \psi_t^{i_1, \dots, i_k} \nabla V_{i_1, \dots, i_k}(\phi_s) \bar{J}_s, \quad \bar{J}_0 = I.$$

Once again, the methodology of Proposition 4.3 easily yields our claim. □

Corollary 4.7 *Under the same assumptions as in Theorem 4.6, the following holds:*

(i) *For any $0 < \gamma < H$ and $q \geq 1$, we have*

$$\mathbf{E}[\|Dy_t\|_{\gamma, \infty}^q] + \mathbf{E}[\|J_{0,\cdot}\|_{\gamma, \infty}^q] + \mathbf{E}[\|J_{0,\cdot}^{-1}\|_{\gamma, \infty}^q] < c_{T,q} \tag{50}$$

for a finite constant $c_{T,q}$.

- (ii) *As a consequence, inequality (50) also holds when the $\|\cdot\|_{\gamma, \infty}$ norms are replaced by norms in \mathcal{H} , where \mathcal{H} has been defined in Sect. 2.3.1.*
- (iii) *For any smooth bounded vector field U on \mathbb{R}^m and $t \in [0, T]$, set $Z_t^U := \langle J_{0,t}^{-1}U(y_t), \eta \rangle$. Then Z^U is a controlled process and satisfies the inequality $\mathbf{E}[\mathcal{N}^q[Z^U; \mathcal{Q}^\gamma(\mathbb{R}^m)]] \leq c_{T,q}$ for any $q \geq 1$ and a finite constant $c_{T,q}$.*

Proof Going back to (46), it is readily checked that all the terms $u \mapsto D_u \psi_t^{i_1, \dots, i_k}$ are C_1^γ -Hölder continuous on $[0, t]$ for any $\gamma < H$, since the elements $\psi_t^{i_1, \dots, i_k}$ are nice multiple integrals with respect to B . Moreover, we have

$$\mathbf{E}[\|D\psi_t^{i_1, \dots, i_k}\|_{\gamma, \infty}^q] < \infty$$

for any $m \geq 1$. This easily yields $\mathbf{E}[\|Dy_t^{i_1, \dots, i_k}\|_{\gamma, \infty}^q] < \infty$ by a standard application of Gronwall’s lemma, as in the proof of Theorem 4.4.

Our second assertion stems from the fact that one can choose $1/2 - H < \gamma < H$, since $H > 1/3$. For such γ , we have $C_1^\gamma \subset \mathcal{H}$, which ends the proof.

Finally, our claim (iii) derives from the fact that the equation governing Z^U is of the following form:

$$Z_t^U = \langle \eta, U(a) \rangle + \sum_{j=1}^d \int_0^t Z_s^{[U, V_j]} dB_s^j. \tag{51}$$

The process Z^U can thus be decomposed as a controlled process as in Sect. 2.2, and since we already have estimates for $J_{0,t}^{-1}$ and y_t , the bound on $\mathbf{E}[\mathcal{N}^q[Z^U; \mathcal{Q}^\gamma(\mathbb{R}^m)]]$ follows easily. □

4.4 Proof of Theorem 1.1

As mentioned in the introduction, once we have shown the integrability of the Malliavin derivative and proved a Norris-type lemma, the proof of our main theorem goes

along classical lines. We have chosen to follow here the exposition of [16], to which we refer for further details.

Step 1: Reduction to a Lower Bound on Hölder Norms Recall that the process Z^U has been defined for any smooth vector field U in Corollary 4.7. For any $p \geq 1$, our first goal is to reduce our problem to the existence of a constant c_p such that

$$\mathbf{P}(\|Z^{V_k}\|_{\alpha,\infty} \leq \varepsilon) \leq c_p \varepsilon^p \tag{52}$$

for $1 \leq k \leq d$, a given $\alpha \in (1/3, 1/2)$, and all $\varepsilon \in (0, 1)$, and where we observe that all the norms below are understood as norms on $[0, T]$.

Indeed, according to [16, Relation (4.9)], the smoothness of density can be obtained from the estimate

$$\mathbf{P}(\|Z^{V_k}\|_{\mathcal{H}} \leq \varepsilon) \leq c_p \varepsilon^p,$$

where we recall that \mathcal{H} has been defined in Sect. 2.3.1. Furthermore, we have

$$\mathbf{P}(\|Z^{V_k}\|_{\mathcal{H}} \leq \varepsilon) \leq \mathbf{P}(\|Z^{V_k}\|_{L^2} \leq \varepsilon) \leq \mathbf{P}(\|Z^{V_k}\|_{L^1} \leq \varepsilon).$$

It is thus sufficient for our purposes to check that

$$\mathbf{P}(\|Z^{V_k}\|_{L^1} \leq \varepsilon) \leq c_p \varepsilon^p. \tag{53}$$

In order to go from (53) to (52), let us use our interpolation bound (23) in the following form: for any $0 < \eta < 1$ and $0 < \alpha < \rho < H$, we have

$$\|b\|_{L^1} \geq \eta^{1/(\rho-\alpha)} (\|b\|_{\alpha,\infty} - C_{\alpha,\rho} \eta \|b\|_{\rho,\infty}).$$

Take now $\delta \in (0, 1)$ to be fixed later on and $\eta^{1/(\rho-\alpha)} = \varepsilon^{1-\delta}$, that is, $\eta = \varepsilon^{(\rho-\alpha)(1-\delta)}$. Then

$$\mathbf{P}(\|Z^{V_k}\|_{L^1} \leq \varepsilon) \leq \mathbf{P}(\|Z^{V_k}\|_{\alpha,\infty} \leq 2\varepsilon^\delta) + R, \quad \text{where } R = \mathbf{P}\left(\|Z^{V_k}\|_{\rho,\infty} \geq \frac{1}{4c} \varepsilon^\nu\right), \tag{54}$$

with $\nu = \rho - \alpha - (1 + (\rho - \alpha))\delta$. Choose now δ small enough, so that $\nu > 0$. Since $\|Z^{V_k}\|_{\rho,\infty}$ admits moments of any order according to Corollary 4.7, it is easily checked that R can be made smaller than any quantity of the form $c_q \varepsilon^q$. It is thus sufficient to prove (52) in order to get the smoothness of density for y_t .

Step 2: An Iterative Procedure For $l \geq 1$ and $x \in \mathbb{R}^m$, let $\mathcal{V}_l(x)$ be the vector space generated by the Lie brackets of order l of our vector fields V_1, \dots, V_d at point x :

$$\mathcal{V}_l(x) = \text{Span}\{[V_{k_1} \cdots V_{k_j}](x); j \leq l, 1 \leq k_1, \dots, k_j \leq d\}.$$

We assume that the vector fields are ℓ -hypoelliptic for a given $\ell > 0$, which can be read as $\mathcal{V}_\ell(x) = \mathbb{R}^m$ for any $x \in \mathbb{R}^m$. In order to start our induction procedure, we set $\alpha_1 = \alpha$, so that we have to prove that $\mathbf{P}(\|Z^{V_k}\|_{\alpha_1,\infty} \leq \varepsilon) \leq c_p \varepsilon^p$.

Recall that Z^{V_k} satisfies the relation

$$Z_t^{V_k} = \langle \eta, V_k(a) \rangle + \sum_{j=1}^d \int_0^t Z_s^{[V_k, V_j]} d\mathbf{B}_s^j.$$

Thus Proposition 3.2 asserts that for any $1/3 < \alpha_2 < \alpha_1 < H$, there exists $q_2 > 0$ such that

$$\mathbf{P}\left(\left(\|Z^{V_k}\|_{\alpha_1, \infty} \leq \varepsilon\right) \cap \left(\bigcup_{j=1}^d \left(\|Z^{[V_k, V_{k_2}]}\|_{\alpha_2, \infty} > \varepsilon^{q_2}\right)\right)\right) \leq c_p \varepsilon^p.$$

Relation (52) is thus implied by

$$\mathbf{P}\left(\left(\|Z^{V_k}\|_{\alpha_1, \infty} \leq \varepsilon\right) \cap \left(\bigcap_{j=1}^d \left(\|Z^{[V_{k_1}, V_{k_2}]}\|_{\alpha_2, \infty} \leq \varepsilon^{q_2}\right)\right)\right) \leq c_p \varepsilon^p.$$

Iterating this procedure, we end up with the following claim to prove: $\mathcal{B}_\ell(\varepsilon) \leq c_p \varepsilon^p$ for all $\varepsilon \in (0, 1)$, with

$$\mathcal{B}_\ell(\varepsilon) = \mathbf{P}\left(\|Z^{V_k}\|_{\alpha_1, \infty} \leq \varepsilon, \|Z^{[V_{k_1}, V_{k_2}]}\|_{\alpha_2, \infty} \leq \varepsilon^{q_2}, \dots, \|Z^{[V_{k_1} \cdots V_{k_\ell}]}\|_{\alpha_\ell, \infty} \leq \varepsilon^{q_\ell}\right),$$

where the intersection above extends to all possible combinations $1 \leq k_1, \dots, k_\ell \leq d$, and where $1/3 < \alpha_\ell < \dots < \alpha_1 < H$.

Going back now to the very definition of Z^U as $Z_t^U = \langle J_{0,t}^{-1}U(y_t), \eta \rangle$, it is readily checked that

$$\mathcal{B}_\ell(\varepsilon) \leq \mathbf{P}\left(\langle \eta, V_{k_1}(a) \rangle \leq \varepsilon, \dots, \langle \eta, [V_{k_1} \cdots V_{k_\ell}](a) \rangle \leq \varepsilon^{q_\ell}\right).$$

Owing to the fact that $\mathcal{V}_\ell(a) = \mathbb{R}^m$, we thus have $\mathcal{B}_\ell(\varepsilon) = 0$ for ε small enough, which ends the proof.

Acknowledgements Y. Hu is partially supported by a grant from the Simons Foundation #209206. S. Tindel is partially supported by the ANR grant ECRU. We would like to thank Ivan Nourdin for pointing us out the use of Hermite polynomials in the computation of 4th-order variations (see the proof of Proposition 3.2, Step 2).

References

1. Baudoin, F.: An Introduction to the Geometry of Stochastic Flows. Imperial College Press, London (2005)
2. Baudoin, F., Coutin, L.: Operators associated with a stochastic differential equation driven by fractional Brownian motions. *Stoch. Process. Appl.* **117**(5), 550–574 (2007)
3. Baudoin, F., Hairer, M.: A version of Hörmander’s theorem for the fractional Brownian motion. *Probab. Theory Relat. Fields* **139**(3–4), 373–395 (2007)
4. Baudoin, F., Ouyang, C., Tindel, S.: Gaussian bounds for the density of solutions of stochastic differential equations driven by fractional Brownian motions. In preparation
5. Borell, C.: Tail probabilities in Gauss space. In: *Vector Space Measures and Applications II* (Proc. Conf., Univ. Dublin), Dublin, 1977. Lecture Notes in Math., vol. 77, pp. 73–82. Springer, Berlin (1978)

6. Cass, T., Friz, P.: Densities for rough differential equations under Hörmander's condition. *Ann. Math.* **171**(3), 2115–2141 (2010)
7. Chen, K.: Integration of paths, Geometric invariants and a generalized Baker–Hausdorff formula. *Ann. Math.* **65**(1) (1957)
8. Deya, A., Tindel, S.: Rough Volterra equations 2: convolutional generalized integrals. Arxiv preprint (2008)
9. Driscoll, P.: Smoothness of density for the area process of fractional Brownian motion. Arxiv preprint (2010)
10. Fernique, X.: Fonctions aléatoires gaussiennes, vecteurs aléatoires gaussiens. Centre de Recherches Mathématiques, Montreal (1997)
11. Ferreira, A., Utzet, F.: Lévy area for Gaussian processes: A double Wiener–Itô integral approach. Arxiv preprint (2010)
12. Friz, P., Victoir, N.: *Multidimensional Stochastic Processes as Rough Paths: Theory and Applications*. Cambridge University Press, Cambridge (2010)
13. Guasoni, P.: No arbitrage under transaction costs, with fractional Brownian motion and beyond. *Math. Finance* **16**(3), 569–582 (2006)
14. Gubinelli, M.: Controlling rough paths. *J. Funct. Anal.* **216**, 86–140 (2004)
15. Gubinelli, M., Tindel, S.: Rough evolution equations. *Ann. Probab.* **38**(1), 1–75 (2010)
16. Hairer, M.: On Malliavin's proof of Hörmander's theorem. Arxiv preprint (2011)
17. Hairer, M., Pillai, N.: Regularity of laws and ergodicity of hypoelliptic SDEs driven by rough paths. Arxiv preprint (2011)
18. Hu, Y.: Calculs formels sur les EDS de Stratonovitch. In: *Séminaire de Probabilités, XXIV, 1988/89*. Lecture Notes in Math., vol. 1426, pp. 453–460. Springer, Berlin (1990)
19. Hu, Y., Nualart, D.: Differential equations driven by Hölder continuous functions of order greater than $1/2$. In: *Abel Symp.*, vol. 2, pp. 349–413 (2007)
20. Kou, S., Sunney-Xie, X.: Generalized Langevin equation with fractional Gaussian noise: subdiffusion within a single protein molecule. *Phys. Rev. Lett.* **93**(18) (2004)
21. Lyons, T., Qian, Z.: *System Control and Rough Paths*. Oxford University Press, London (2002)
22. Malliavin, P.: Stochastic calculus of variation and hypoelliptic operators. In: *Proceedings of the International Symposium on Stochastic Differential Equations (Res. Inst. Math. Sci., Kyoto Univ.)*, Kyoto, 1976, pp. 195–263. Wiley, New York (1978)
23. Neuenkirch, A., Nourdin, I., Rößler, A., Tindel, S.: Trees and asymptotic developments for fractional diffusion processes. *Ann. Inst. Henri Poincaré Probab. Stat.* **45**(1), 157–174 (2009)
24. Norris, J.: Simplified Malliavin calculus. In: *Séminaire de Probabilités 20 (1984/85)*. Lecture Notes in Math, vol. 1204 pp. 101–130. Springer, Berlin (1986)
25. Nualart, D.: *Malliavin Calculus and Related Topics*. Springer, Berlin (2006)
26. Nualart, D., Răşcanu, A.: Differential equations driven by fractional Brownian motion. *Collect. Math.* **53**(1), 55–81 (2002)
27. Nualart, D., Saussereau, B.: Malliavin calculus for stochastic differential equations driven by a fractional Brownian motion. *Stoch. Process. Appl.* **119**(2), 391–409 (2009)
28. Odde, D., Tanaka, E., Hawkins, S., Buettner, H.: Stochastic dynamics of the nerve growth cone and its microtubules during neurite outgrowth. *Biotechnol. Bioeng.* **50**(4), 452–461 (1996)
29. Strichartz, R.: The Campbell–Baker–Hausdorff–Dynkin formula and solutions of differential equations. *J. Funct. Anal.* **72**, 320–345 (1987)
30. Willinger, W., Taqqu, M.S., Teverovsky, V.: Stock market prices and long-range dependence. *Finance Stoch.* **3**(1), 1–13 (1999)
31. Yamato, Y.: Stochastic differential equations and nilpotent Lie algebras. *Z. Wahrscheinlichkeitstheor. Verw. Geb.* **47**(2), 213–229 (1979)
32. Zähle, M.: Integration with respect to fractal functions and stochastic calculus I. *Probab. Theory Relat. Fields* **111**, 333–374 (1998)