

## MA 261 PRACTICE PROBLEMS

1. If the line  $\ell$  has symmetric equations

$$\frac{x-1}{2} = \frac{y}{-3} = \frac{z+2}{7},$$

find a vector equation for the line  $\ell'$  that contains the point  $(2, 1, -3)$  and is parallel to  $\ell$ .

- A.  $\vec{r} = (1 + 2t)\vec{i} - 3t\vec{j} + (-2 + 7t)\vec{k}$
- B.  $\vec{r} = (2 + t)\vec{i} - 3\vec{j} + (7 - 2t)\vec{k}$
- C.  $\vec{r} = (2 + 2t)\vec{i} + (1 - 3t)\vec{j} + (-3 + 7t)\vec{k}$
- D.  $\vec{r} = (2 + 2t)\vec{i} + (-3 + t)\vec{j} + (7 - 3t)\vec{k}$
- E.  $\vec{r} = (2 + t)\vec{i} + \vec{j} + (7 - 3t)\vec{k}$

2. Find parametric equations of the line containing the points  $(1, -1, 0)$  and  $(-2, 3, 5)$ .

- A.  $x = 1 - 3t, y = -1 + 4t, z = 5t$
- B.  $x = t, y = -t, z = 0$
- C.  $x = 1 - 2t, y = -1 + 3t, z = 5t$
- D.  $x = -2t, y = 3t, z = 5t$
- E.  $x = -1 + t, y = 2 - t, z = 5$

3. Find an equation of the plane that contains the point  $(1, -1, -1)$  and has normal vector  $\frac{1}{2}\vec{i} + 2\vec{j} + 3\vec{k}$ .

- A.  $x - y - z + \frac{9}{2} = 0$
- B.  $x + 4y + 6z + 9 = 0$
- C.  $\frac{x-1}{\frac{1}{2}} = \frac{y+1}{2} = \frac{z+1}{3}$
- D.  $x - y - z = 0$
- E.  $\frac{1}{2}x + 2y + 3z = 1$

4. Find an equation of the plane that contains the points  $(1, 0, -1)$ ,  $(-5, 3, 2)$ , and  $(2, -1, 4)$ .

- A.  $6x - 11y + z = 5$
- B.  $6x + 11y + z = 5$
- C.  $11x - 6y + z = 0$
- D.  $\vec{r} = 18\vec{i} - 33\vec{j} + 3\vec{k}$
- E.  $x - 6y - 11z = 12$

5. Find parametric equations of the line tangent to the curve  $\vec{r}(t) = t\vec{i} + t^2\vec{j} + t^3\vec{k}$  at the point  $(2, 4, 8)$

- A.  $x = 2 + t, y = 4 + 4t, z = 8 + 12t$
- B.  $x = 1 + 2t, y = 4 + 4t, z = 12 + 8t$
- C.  $x = 2t, y = 4t, z = 8t$
- D.  $x = t, y = 4t, z = 12t$
- E.  $x = 2 + t, y = 4 + 2t, z = 8 + 3t$

6. The position function of an object is

$$\vec{r}(t) = \cos t\vec{i} + 3 \sin t\vec{j} - t^2\vec{k}$$

Find the velocity, acceleration, and speed of the object when  $t = \pi$ .

|    | Velocity                           | Acceleration              | Speed                |
|----|------------------------------------|---------------------------|----------------------|
| A. | $-\vec{i} - \pi^2\vec{k}$          | $-3\vec{j} - 2\pi\vec{k}$ | $\sqrt{1 + \pi^4}$   |
| B. | $\vec{i} - 3\vec{j} + 2\pi\vec{k}$ | $-\vec{i} - 2\vec{k}$     | $\sqrt{10 + 4\pi^2}$ |
| C. | $3\vec{j} - 2\pi\vec{k}$           | $-\vec{i} - 2\vec{k}$     | $\sqrt{9 + 4\pi^2}$  |
| D. | $-3\vec{j} - 2\pi\vec{k}$          | $\vec{i} - 2\vec{k}$      | $\sqrt{9 + 4\pi^2}$  |
| E. | $\vec{i} - 2\vec{k}$               | $-3\vec{j} - 2\pi\vec{k}$ | $\sqrt{5}$           |

7. A smooth parametrization of the semicircle which passes through the points  $(1, 0, 5)$ ,  $(0, 1, 5)$  and  $(-1, 0, 5)$  is

- A.  $\vec{r}(t) = \sin t\vec{i} + \cos t\vec{j} + 5\vec{k}, 0 \leq t \leq \pi$
- B.  $\vec{r}(t) = \cos t\vec{i} + \sin t\vec{j} + 5\vec{k}, 0 \leq t \leq \pi$
- C.  $\vec{r}(t) = \cos t\vec{i} + \sin t\vec{j} + 5\vec{k}, \frac{\pi}{2} \leq t \leq \frac{3\pi}{2}$
- D.  $\vec{r}(t) = \cos t\vec{i} + \sin t\vec{j} + 5\vec{k}, 0 \leq t \leq \frac{\pi}{2}$
- E.  $\vec{r}(t) = \sin t + \cos t\vec{j} + 5\vec{k}, \frac{\pi}{2} \leq t \leq \frac{3\pi}{2}$

8. The length of the curve  $\vec{r}(t) = \frac{2}{3}(1+t)^{\frac{3}{2}}\vec{i} + \frac{2}{3}(1-t)^{\frac{3}{2}}\vec{j} + t\vec{k}$ ,  $-1 \leq t \leq 1$  is

- A.  $\sqrt{3}$
- B.  $\sqrt{2}$
- C.  $\frac{1}{2}\sqrt{3}$
- D.  $2\sqrt{3}$
- E.  $\sqrt{2}$

9. The level curves of the function  $f(x, y) = \sqrt{1 - x^2 - 2y^2}$  are

- A. circles
- B. lines
- C. parabolas
- D. hyperbolas
- E. ellipses

10. The level surface of the function  $f(x, y, z) = z - x^2 - y^2$  that passes through the point  $(1, 2, -3)$  intersects the  $(x, z)$ -plane ( $y = 0$ ) along the curve

- A.  $z = x^2 + 8$
- B.  $z = x^2 - 8$
- C.  $z = x^2 + 5$
- D.  $z = -x^2 - 8$
- E. does not intersect the  $(x, z)$ -plane

11. Match the graphs of the equations with their names:

- |                             |                 |
|-----------------------------|-----------------|
| (1) $x^2 + y^2 + z^2 = 4$   | (a) paraboloid  |
| (2) $x^2 + z^2 = 4$         | (b) sphere      |
| (3) $x^2 + y^2 = z^2$       | (c) cylinder    |
| (4) $x^2 + y^2 = z$         | (d) double cone |
| (5) $x^2 + 2y^2 + 3z^2 = 1$ | (e) ellipsoid   |

- A. 1b, 2c, 3d, 4a, 5e
- B. 1b, 2c, 3a, 4d, 5e
- C. 1e, 2c, 3d, 4a, 5b
- D. 1b, 2d, 3a, 4c, 5e
- E. 1d, 2a, 3b, 4e, 5c

12. Suppose that  $w = u^2/v$  where  $u = g_1(t)$  and  $v = g_2(t)$  are differentiable functions of  $t$ . If  $g_1(1) = 3$ ,  $g_2(1) = 2$ ,  $g'_1(1) = 5$  and  $g'_2(1) = -4$ , find  $\frac{dw}{dt}$  when  $t = 1$ .

- A. 6
- B.  $33/2$
- C.  $-24$
- D. 33
- E. 24

13. If  $w = e^{uv}$  and  $u = r + s$ ,  $v = rs$ , find  $\frac{\partial w}{\partial r}$ .

- A.  $e^{(r+s)rs}(2rs + r^2)$
- B.  $e^{(r+s)rs}(2rs + s^2)$
- C.  $e^{(r+s)rs}(2rs + r^2)$
- D.  $e^{(r+s)rs}(1 + s)$
- E.  $e^{(r+s)rs}(r + s^2)$ .

14. If  $f(x, y) = \cos(xy)$ ,  $\frac{\partial^2 f}{\partial x \partial y} =$
- A.  $-xy \cos(xy)$       B.  $-xy \cos(xy) - \sin(xy)$       C.  $-\sin(xy)$   
 D.  $xy \cos(xy) + \sin(xy)$       E.  $-\cos(xy)$
15. Assuming that the equation  $xy^2 + 3z = \cos(z^2)$  defines  $z$  implicitly as a function of  $x$  and  $y$ , find  $\frac{\partial z}{\partial x}$ .
- A.  $\frac{y^2}{3-\sin(z^2)}$       B.  $\frac{-y^2}{3+\sin(z^2)}$       C.  $\frac{y^2}{3+2z \sin(z^2)}$       D.  $\frac{-y^2}{3+2z \sin(z^2)}$       E.  $\frac{-y^2}{3-2z \sin(z^2)}$
16. If  $f(x, y) = xy^2$ , then  $\nabla f(2, 3) =$
- A.  $12\vec{i} + 9\vec{j}$       B.  $18\vec{i} + 18\vec{j}$       C.  $9\vec{i} + 12\vec{j}$       D.  $21$       E.  $\sqrt{2}$ .
17. Find the directional derivative of  $f(x, y) = 5 - 4x^2 - 3y$  at  $(x, y)$  towards the origin
- A.  $-8x - 3$       B.  $\frac{-8x^2 - 3y}{\sqrt{x^2 + y^2}}$       C.  $\frac{-8x - 3}{\sqrt{64x^2 + 9}}$       D.  $8x^2 + 3y$       E.  $\frac{8x^2 + 3y}{\sqrt{x^2 + y^2}}$ .
18. For the function  $f(x, y) = x^2y$ , find a unit vector  $\vec{u}$  for which the directional derivative  $D_{\vec{u}}f(2, 3)$  is zero.
- A.  $\vec{i} + 3\vec{j}$       B.  $\frac{i+3j}{\sqrt{10}}$       C.  $\vec{i} - 3\vec{j}$       D.  $\frac{i-3j}{\sqrt{10}}$       E.  $\frac{3i-j}{\sqrt{10}}$ .
19. Find a vector pointing in the direction in which  $f(x, y, z) = 3xy - 9xz^2 + y$  increases most rapidly at the point  $(1, 1, 0)$ .
- A.  $3\vec{i} + 4\vec{j}$       B.  $\vec{i} + \vec{j}$       C.  $4\vec{i} - 3\vec{j}$       D.  $2\vec{i} + \vec{k}$       E.  $-\vec{i} + \vec{j}$ .
20. Find a vector that is normal to the graph of the equation  $2 \cos(\pi xy) = 1$  at the point  $(\frac{1}{6}, 2)$ .
- A.  $6\vec{i} + \vec{j}$       B.  $-\sqrt{3}\vec{i} - \vec{j}$       C.  $12\vec{i} + \vec{j}$       D.  $\vec{j}$       E.  $12\vec{i} - \vec{j}$ .
21. Find an equation of the tangent plane to the surface  $x^2 + 2y^2 + 3z^2 = 6$  at the point  $(1, 1, -1)$ .
- A.  $-x + 2y + 3z = 2$       B.  $2x + 4y - 6z = 6$       C.  $x - 2y + 3z = -4$   
 D.  $2x + 4y - 6z = 0$       E.  $x + 2y - 3z = 6$ .
22. Find an equation of the plane tangent to the graph of  $f(x, y) = \pi + \sin(\pi x^2 + 2y)$  when  $(x, y) = (2, \pi)$ .
- A.  $4\pi x + 2y - z = 9\pi$       B.  $4x + 2\pi y - z = 10\pi$       C.  $4\pi x + 2\pi y + z = 10\pi$   
 D.  $4x + 2\pi y - z = 9\pi$       E.  $4\pi x + 2y + z = 9\pi$ .

23. The differential  $df$  of the function  $f(x, y, z) = xe^{y^2-z^2}$  is

- A.  $df = xe^{y^2-z^2}dx + xe^{y^2-z^2}dy + xe^{y^2-z^2}dz$
- B.  $df = xe^{y^2-z^2}dx dy dz$
- C.  $df = e^{y^2-z^2}dx - 2xye^{y^2-z^2}dy + 2xze^{y^2-z^2}dz$
- D.  $df = e^{y^2-z^2}dx + 2xye^{y^2-z^2}dy - 2xze^{y^2-z^2}dz$
- E.  $df = e^{y^2-z^2}(1 + 2xy - 2xz)$

24. The function  $f(x, y) = 2x^3 - 6xy - 3y^2$  has

- A. a relative minimum and a saddle point
- B. a relative maximum and a saddle point
- C. a relative minimum and a relative maximum
- D. two saddle points
- E. two relative minima.

25. Consider the problem of finding the minimum value of the function  $f(x, y) = 4x^2 + y^2$  on the curve  $xy = 1$ . In using the method of Lagrange multipliers, the value of  $\lambda$  (even though it is not needed) will be

- A. 2
- B. -2
- C.  $\sqrt{2}$
- D.  $\frac{1}{\sqrt{2}}$
- E. 4.

26. Evaluate the iterated integral  $\int_1^3 \int_0^x \frac{1}{x} dy dx$ .

- A.  $-\frac{8}{9}$
- B. 2
- C.  $\ln 3$
- D. 0
- E.  $\ln 2$ .

27. Consider the double integral,  $\iint_R f(x, y)dA$ , where  $R$  is the portion of the disk  $x^2 + y^2 \leq 1$ , in the upper half-plane,  $y \geq 0$ . Express the integral as an iterated integral.

- A.  $\int_{-1}^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} f(x, y) dy dx$
- B.  $\int_{-1}^0 \int_0^{\sqrt{1-x^2}} f(x, y) dy dx$
- C.  $\int_{-1}^1 \int_0^{\sqrt{1-x^2}} f(x, y) dy dx$
- D.  $\int_0^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} f(x, y) dy dx$
- E.  $\int_0^1 \int_0^{\sqrt{1-x^2}} f(x, y) dy dx$ .

28. Find  $a$  and  $b$  for the correct interchange of order of integration:

$$\int_0^2 \int_{x^2}^{2x} f(x, y) dy dx = \int_0^4 \int_a^b f(x, y) dx dy.$$

- A.  $a = y^2, b = 2y$
- B.  $a = \frac{y}{2}, b = \sqrt{y}$
- C.  $a = \frac{y}{2}, b = y$
- D.  $a = \sqrt{y}, b = \frac{y}{2}$
- E. cannot be done without explicit knowledge of  $f(x, y)$ .

29. Evaluate the double integral  $\iint_R y dA$ , where  $R$  is the region of the  $(x, y)$ -plane inside the triangle with vertices  $(0, 0)$ ,  $(2, 0)$  and  $(2, 1)$ .

- A. 2
- B.  $\frac{8}{3}$
- C.  $\frac{2}{3}$
- D. 1
- E.  $\frac{1}{3}$ .

30. The volume of the solid region in the first octant bounded above by the parabolic sheet  $z = 1 - x^2$ , below by the  $xy$  plane, and on the sides by the planes  $y = 0$  and  $y = x$  is given by the double integral

- A.  $\int_0^1 \int_0^x (1 - x^2) dy dx$
- B.  $\int_0^1 \int_0^{1-x^2} x dy dx$
- C.  $\int_{-1}^1 \int_{-x}^x (1 - x^2) dy dx$
- D.  $\int_0^1 \int_x^0 (1 - x^2) dy dx$
- E.  $\int_0^1 \int_x^{1-x^2} dy dx$ .

31. The area of one leaf of the three-leaved rose bounded by the graph of  $r = 5 \sin 3\theta$  is

A.  $\frac{5\pi}{6}$       B.  $\frac{25\pi}{12}$       C.  $\frac{25\pi}{6}$       D.  $\frac{5\pi}{3}$       E.  $\frac{25\pi}{3}$ .

32. Find the area of the portion of the plane  $x + 3y + 2z = 6$  that lies in the first octant.

A.  $3\sqrt{11}$       B.  $6\sqrt{7}$       C.  $6\sqrt{14}$       D.  $3\sqrt{14}$       E.  $6\sqrt{11}$ .

33. A solid region in the first octant is bounded by the surfaces  $z = y^2$ ,  $y = x$ ,  $y = 0$ ,  $z = 0$  and  $x = 4$ . The volume of the region is

A. 64      B.  $\frac{64}{3}$       C.  $\frac{32}{3}$       D. 32      E.  $\frac{16}{3}$ .

34. An object occupies the region bounded above by the sphere  $x^2 + y^2 + z^2 = 32$  and below by the upper nappe of the cone  $z^2 = x^2 + y^2$ . The mass density at any point of the object is equal to its distance from the  $xy$  plane. Set up a triple integral in rectangular coordinates for the total mass  $m$  of the object.

A.  $\int_{-4}^4 \int_{-\sqrt{16-x^2}}^{\sqrt{16-x^2}} \int_{-\sqrt{x^2+y^2}}^{\sqrt{32-x^2-y^2}} z dz dy dx$   
 B.  $\int_{-4}^4 \int_{-\sqrt{16-x^2}}^{\sqrt{16-x^2}} \int_{\sqrt{x^2+y^2}}^{\sqrt{32-x^2-y^2}} z dz dy dx$   
 C.  $\int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{-\sqrt{x^2+y^2}}^{\sqrt{32-x^2-y^2}} z dz dy dx$   
 D.  $\int_0^4 \int_0^{\sqrt{16-x^2}} \int_{\sqrt{x^2+y^2}}^{\sqrt{32-x^2-y^2}} z dz dy dx$   
 E.  $\int_{-4}^4 \int_{-\sqrt{16-x^2}}^{\sqrt{16-x^2}} \int_{\sqrt{x^2+y^2}}^{\sqrt{32-x^2-y^2}} xy dz dy dx.$

35. Do problem 34 in spherical coordinates.

A.  $\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^{\sqrt{32}} \rho^3 \cos \varphi \sin \varphi d\rho d\varphi d\theta$   
 B.  $\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^{\sqrt{32}} \rho \cos \varphi \sin \varphi d\rho d\varphi d\theta$   
 C.  $\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^{\sqrt{32}} \rho^3 \sin^2 \varphi d\rho d\varphi d\theta$   
 D.  $\int_0^{2\pi} \int_0^{\frac{\pi}{2}} \int_0^{\sqrt{32}} \rho^3 \cos \varphi \sin \varphi d\rho d\varphi d\theta$   
 E.  $\int_0^{2\pi} \int_0^{\frac{\pi}{4}} \int_0^{\sqrt{32}} \rho \cos \varphi d\rho d\varphi d\theta.$

36. The double integral  $\int_0^1 \int_0^{\sqrt{1-x^2}} y^2(x^2 + y^2)^3 dy dx$  when converted to polar coordinates becomes

A.  $\int_0^\pi \int_0^1 r^9 \sin^2 \theta dr d\theta$       B.  $\int_0^{\frac{\pi}{2}} \int_0^1 r^8 \sin^2 \theta dr d\theta$       C.  $\int_0^\pi \int_0^1 r^8 \sin \theta dr d\theta$   
 D.  $\int_0^{\frac{\pi}{2}} \int_0^1 r^8 \sin \theta dr d\theta$       E.  $\int_0^{\frac{\pi}{2}} \int_0^1 r^9 \sin^2 \theta dr d\theta.$

37. Which of the triple integrals converts

$$\int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} \int_{\sqrt{x^2+y^2}}^2 dz dy dx$$

from rectangular to cylindrical coordinates?

A.  $\int_0^\pi \int_0^2 \int_r^2 r dz dr d\theta$       B.  $\int_0^{2\pi} \int_0^2 \int_r^2 r dz dr d\theta$       C.  $\int_0^{2\pi} \int_{-2}^2 \int_r^2 r dz dr d\theta$   
 D.  $\int_0^\pi \int_0^2 \int_r^2 r dz dr d\theta$       E.  $\int_0^{\frac{2\pi}{2}} \int_{-2}^2 \int_r^2 r dz dr d\theta.$

38. If  $D$  is the solid region above the  $xy$ -plane that is between  $z = \sqrt{4 - x^2 - y^2}$  and  $z = \sqrt{1 - x^2 - y^2}$ , then  $\iiint_D \sqrt{x^2 + y^2 + z^2} dV =$

A.  $\frac{14\pi}{3}$       B.  $\frac{16\pi}{3}$       C.  $\frac{15\pi}{2}$       D.  $8\pi$       E.  $15\pi.$

39. Determine which of the vector fields below are conservative, i. e.  $\vec{F} = \text{grad } f$  for some function  $f$ .

1.  $\vec{F}(x, y) = (xy^2 + x)\vec{i} + (x^2y - y^2)\vec{j}$ .
2.  $\vec{F}(x, y) = \frac{x}{y}\vec{i} + \frac{y}{x}\vec{j}$ .
3.  $\vec{F}(x, y, z) = ye^z\vec{i} + (xe^z + e^y)\vec{j} + (xy + 1)e^z\vec{k}$ .

A. 1 and 2      B. 1 and 3      C. 2 and 3      D. 1 only      E. all three

40. Let  $\vec{F}$  be any vector field whose components have continuous partial derivatives up to second order, let  $f$  be any real valued function with continuous partial derivatives up to second order, and let  $\nabla = \vec{i} \frac{\partial}{\partial x} + \vec{j} \frac{\partial}{\partial y} + \vec{k} \frac{\partial}{\partial z}$ . Find the incorrect statement.

- A.  $\text{curl}(\text{grad } f) = \vec{0}$       B.  $\text{div}(\text{curl } \vec{F}) = 0$       C.  $\text{grad}(\text{div } \vec{F}) = 0$   
 D.  $\text{curl } \vec{F} = \nabla \times \vec{F}$       E.  $\text{div } \vec{F} = \nabla \cdot \vec{F}$

41. A wire lies on the  $xy$ -plane along the curve  $y = x^2$ ,  $0 \leq x \leq 2$ . The mass density (per unit length) at any point  $(x, y)$  of the wire is equal to  $x$ . The mass of the wire is

- A.  $(17\sqrt{17} - 1)/12$       B.  $(17\sqrt{17} - 1)/8$       C.  $17\sqrt{17} - 1$   
 D.  $(\sqrt{17} - 1)/3$       E.  $(\sqrt{17} - 1)/12$

42. Evaluate  $\int_C \vec{F} \cdot d\vec{r}$  where  $\vec{F}(x, y) = y\vec{i} + x^2\vec{j}$  and  $C$  is composed of the line segments from  $(0, 0)$  to  $(1, 0)$  and from  $(1, 0)$  to  $(1, 2)$ .

- A. 0      B.  $\frac{2}{3}$       C.  $\frac{5}{6}$       D. 2      E. 3

43. Evaluate the line integral

$$\int_C x \, dx + y \, dy + xy \, dz$$

where  $C$  is parametrized by  $\vec{r}(t) = \cos t\vec{i} + \sin t\vec{j} + \cos t\vec{k}$  for  $-\frac{\pi}{2} \leq t \leq 0$ .

- A. 1      B. -1      C.  $\frac{1}{3}$       D.  $-\frac{1}{3}$       E. 0

44. Are the following statements true or false?

1. The line integral  $\int_C (x^3 + 2xy)dx + (x^2 - y^2)dy$  is independent of path in the  $xy$ -plane.
2.  $\int_C (x^3 + 2xy)dx + (x^2 - y^2)dy = 0$  for every closed oriented curve  $C$  in the  $xy$ -plane.
3. There is a function  $f(x, y)$  defined in the  $xy$ -plane, such that  
 $\text{grad } f(x, y) = (x^3 + 2xy)\vec{i} + (x^2 - y^2)\vec{j}$ .

- A. all three are false      B. 1 and 2 are false, 3 is true      C. 1 and 2 are true, 3 is false  
 D. 1 is true, 2 and 3 are false      E. all three are true

45. Evaluate  $\int_C y^2dx + 6xy \, dy$  where  $C$  is the boundary curve of the region bounded by  $y = \sqrt{x}$ ,  $y = 0$  and  $x = 4$ , in the counterclockwise direction.

- A. 0      B. 4      C. 8      D. 16      E. 32

46. If  $C$  goes along the  $x$ -axis from  $(0, 0)$  to  $(1, 0)$ , then along  $y = \sqrt{1 - x^2}$  to  $(0, 1)$ , and then back to  $(0, 0)$  along the  $y$ -axis, then  $\int_C xy \, dy =$

- A.  $-\int_0^1 \int_0^{\sqrt{1-x^2}} y \, dy \, dx$       B.  $\int_0^1 \int_0^{\sqrt{1-x^2}} y \, dy \, dx$       C.  $-\int_0^1 \int_0^{\sqrt{1-x^2}} x \, dy \, dx$   
 D.  $\int_0^1 \int_0^{\sqrt{1-x^2}} x \, dy \, dx$       E. 0

47. Evaluate  $\int_C \vec{F} \cdot d\vec{r}$ , if  $\vec{F}(x, y) = (xy^2 - 1)\vec{i} + (x^2y - x)\vec{j}$  and  $C$  is the circle of radius 1 centered at  $(1, 2)$  and oriented counterclockwise.

- A. 2      B.  $\pi$       C. 0      D.  $-\pi$       E.  $-2$

48. Green's theorem yields the following formula for the area of a simple region  $R$  in terms of a line integral over the boundary  $C$  of  $R$ , oriented counterclockwise. Area of  $R = \iint_R dA =$

- A.  $-\int_C y \, dx$       B.  $\int_C y \, dx$       C.  $\int_C x \, dx$       D.  $\frac{1}{2} \int_C y \, dx - x \, dy$       E.  $-\int x \, dy$

49. Evaluate the surface integral  $\iint_{\Sigma} x \, dS$  where  $\Sigma$  is the part of the plane  $2x + y + z = 4$  in the first octant.

- A.  $8\sqrt{6}$       B.  $\frac{8}{3}\sqrt{6}$       C.  $\frac{8}{3}\sqrt{14}$       D.  $\frac{\sqrt{14}}{3}$       E.  $\frac{\sqrt{10}}{3}$

50. If  $\Sigma$  is the part of the paraboloid  $z = x^2 + y^2$  with  $z \leq 4$ ,  $\vec{n}$  is the unit normal vector on  $\Sigma$  directed upward, and  $\vec{F}(x, y, z) = x\vec{i} + y\vec{j} + z\vec{k}$ , then  $\iint_{\Sigma} \vec{F} \cdot \vec{n} \, dS =$

- A. 0      B.  $8\pi$       C.  $4\pi$       D.  $-4\pi$       E.  $-8\pi$

51. If  $\vec{F}(x, y, z) = \cos z\vec{i} + \sin z\vec{j} + xy\vec{k}$ ,  $\Sigma$  is the complete boundary of the rectangular solid region bounded by the planes  $x = 0$ ,  $x = 1$ ,  $y = 0$ ,  $y = 1$ ,  $z = 0$  and  $z = \frac{\pi}{2}$ , and  $\vec{n}$  is the outward unit normal on  $\Sigma$ , then  $\iint_{\Sigma} \vec{F} \cdot \vec{n} \, dS =$

- A. 0      B.  $\frac{1}{2}$       C. 1      D.  $\frac{\pi}{2}$       E. 2

52. If  $\vec{F}(x, y, z) = x\vec{i} + y\vec{j} + z\vec{k}$ ,  $\Sigma$  is the unit sphere  $x^2 + y^2 + z^2 = 1$  and  $\vec{n}$  is the outward unit normal on  $\Sigma$ , then  $\iint_{\Sigma} \vec{F} \cdot \vec{n} \, dS =$

- A.  $-4\pi$       B.  $\frac{2\pi}{3}$       C. 0      D.  $\frac{4\pi}{3}$       E.  $4\pi$

## **ANSWERS**

1–C, 2–A, 3–B, 4–B, 5–A, 6–D, 7–B, 8–D, 9–E, 10–B, 11–A, 12–E, 13–B, 14–B,  
15–D, 16–C, 17–E 18–D, 19–A, 20–C, 21–E, 22–A, 23–D, 24–B, 25–E, 26–B, 27–C,  
28–B, 29–E, 30–A, 31–B, 32–D, 33–B, 34–B, 35–A, 36–E, 37–B, 38–C, 39–B, 40–C,  
41–A, 42–D, 43–D, 44–E, 45–D, 46–B, 47–D, 48–A, 49–B, 50–E, 51–A, 52–E