

## MA 224 FORMULAS

### THE SECOND DERIVATIVE TEST

Suppose  $f$  is a function of two variables  $x$  and  $y$ , and that all the second-order partial derivatives are continuous. Let

$$D = f_{xx}f_{yy} - (f_{xy})^2$$

and suppose  $(a, b)$  is a critical point of  $f$ .

1. If  $D(a, b) < 0$ , then  $f$  has a saddle point at  $(a, b)$ ,
2. If  $D(a, b) > 0$  and  $f_{xx}(a, b) < 0$ , then  $f$  has a relative maximum at  $(a, b)$ .
3. If  $D(a, b) > 0$  and  $f_{xx}(a, b) > 0$ , then  $f$  has a relative minimum at  $(a, b)$ .
4. If  $D(a, b) = 0$ , the test is inconclusive.

### LEAST-SQUARES LINE

The equation of the least-squares line for the  $n$  points  $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ , is  $y = mx + b$ , where  $m$  and  $b$  are solutions to the system of equations

$$\begin{aligned}(x_1^2 + x_2^2 + \dots + x_n^2)m + (x_1 + x_2 + \dots + x_n)b &= x_1y_1 + x_2y_2 + \dots + x_ny_n \\ (x_1 + x_2 + \dots + x_n)m + nb &= y_1 + y_2 + \dots + y_n\end{aligned}$$

### TRAPEZOIDAL RULE

$$\int_a^b f(x)dx \equiv \frac{\Delta x}{2} \left[ f(x_0) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(x_n) \right],$$

where  $a = x_0, x_1, x_2, \dots, x_n = b$  subdivides  $[a, b]$  into  $n$  equal subintervals of length  $\Delta x = \frac{b-a}{n}$ .

### ERROR ESTIMATE FOR THE TRAPEZOIDAL RULE

If  $M$  is the maximum value of  $|f''(x)|$  on the interval  $a \leq x \leq b$ , then

$$|E_n| \leq \frac{M(b-a)^3}{12n^2}$$

### GEOMETRIC SERIES

If  $0 < |r| < 1$ , then

$$\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}$$

### TAYLOR SERIES

The Taylor series of  $f(x)$  about  $x = a$  is the power series

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n = f(a) + f'(a)(x-a) + \frac{f^{(2)}(a)}{2!} (x-a)^2 + \dots$$

Examples:

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}, \text{ for } -\infty < x < \infty; \quad \ln x = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} (x-1)^n, \text{ for } 0 < x \leq 2$$