


**MA261-YOLCU      PRACTICE PROBLEMS FOR TEST 2      SPRING 2013**

- (1) Consider  $f(x, y) = x^4 + y^4 - 4xy + 1$ . Show that  $f$  has a local minimum at  $(1, 1)$  and  $(-1, -1)$  and that  $(0, 0)$  is a saddle point of  $f$ .
- (2) Find the extreme values of the function  $f(x, y) = x^2 + 2y^2$  on the circle  $x^2 + y^2 = 1$ . Answer:  $f(0, \pm 1) = 2$  is the maximum,  $f(\pm 1, 0) = 1$  is the minimum value.

- (3) Find the area of **one loop** of the rose  $r = \cos(2\theta)$  sketched below. Answer:  $\frac{\pi}{8}$



- (4) Find the value of the integral  $I = \int_0^{\sqrt{2}} \int_{y^2}^2 y e^{x^2} dx dy$  by interchanging the order of integration.

Answer:

$$I = \int_0^2 \int_0^{\sqrt{x}} y e^{x^2} dy dx = \frac{1}{4}(e^4 - 1).$$

See also similar problems on page: 996 (15.3(#49 – 54)).

- (5) Use the midpoint rule with  $m = n = 2$  to approximate

$$\iint_R (x^2 - 1)y \, dA$$

where  $R$  is the region  $\{(x, y) : 0 \leq x \leq 4, 2 \leq y \leq 4\}$ .

Answer: 96

- (6) Let  $R$  be the region in the first quadrant bounded by  $x = 0$ ,  $x - y = 0$ ,  $x^2 + y^2 = 9$  and  $x + y = 6$ . Evaluate

$$\iint_R \frac{x+y}{x^2+y^2} \, dA.$$

Answer:  $\frac{3}{2}\pi - 3$

- (7) Find the center of mass  $(\bar{x}, \bar{y})$  of the semicircular lamina described by  $\{(x, y) : x^2 + y^2 \leq a^2, y \geq 0\}$  if its density at the point  $(x, y)$  is  $\rho(x, y) = \sqrt{x^2 + y^2}$ .

Answer:  $\bar{x} = 0, \bar{y} = \frac{3a}{2\pi}$

- (8) Find the area of the region described by the intersection of two disks bounded by  $x^2 + y^2 = x$  and  $x^2 + y^2 = y$ .

Answer:  $\frac{\pi}{8} - \frac{1}{4}$

(9) Find  $a, b, c, d, e, f, g, h$  so that

$$\int_0^1 \int_0^y \int_{\sqrt{y}}^1 F(x, y, z) dx dz dy = \int_0^1 \int_a^b \int_c^d F(x, y, z) dy dx dz = \int_0^1 \int_e^f \int_g^h F(x, y, z) dz dy dx$$

Answer:  $a = \sqrt{z}$ ,  $b = 1$ ,  $c = z$ ,  $d = x^2$ ,  $e = 0$ ,  $f = x^2$ ,  $g = 0$ ,  $h = y$ .

(10) Find the volume the solid that lies above the cone  $z = \sqrt{x^2 + y^2}$  below the sphere  $x^2 + y^2 + z^2 = 2z$ .

Answer:  $\pi$

- (11) Let  $T$  be the solid region in the first octant that is bounded by the planes  $y = 2$ ,  $x = 0$ ,  $y = 2x$ ,  $z = 0$ , and  $z = 2y$ . What is the value of the tripple integral  $\iiint_T x \, dV$  ?

Answer: 1

- (12) Let  $c$  be a constant such that  $0 < c < \pi/2$  or  $\pi/2 < c < \pi$ . Show that the equation of the surface  $\phi = c$  converted to rectangular coordinates becomes  $z = \cot(c)\sqrt{x^2 + y^2}$ .

- (13) A lamina  $L$  occupies the triangular region in the  $xy$ -plane with vertices  $(0, 0)$ ,  $(0, 3)$  and  $(3, 3)$ . If the mass density at  $(x, y)$  is  $\rho(x, y) = x + 2y$ , then show that the  $y$ -coordinate of the center of mass of  $L$  is equal to  $\frac{9}{4}$ .

- (14) Let  $E$  be the solid region enclosed by the paraboloid  $z = x^2 + y^2$  and the plane  $2x + z = 4$ . Find the triple integral in cylindrical coordinates that gives the volume  $V(E)$  of solid  $E$ .

Answer:

$$V(E) = \int_0^{2\pi} \int_0^{-\cos\theta + \sqrt{4 + \cos^2\theta}} \int_{r^2}^{4 - 2r\cos\theta} r \, dz \, dr \, d\theta$$

(15) Convert

$$\int_0^3 \int_{-\sqrt{9-x^2}}^{\sqrt{9-x^2}} \int_0^{\sqrt{x^2+y^2}} x y z \, dz \, dy \, dx$$

to spherical coordinates.

Answer:

$$\int_{-\pi/2}^{\pi/2} \int_{\pi/4}^{\pi/2} \int_0^{3/\sin \phi} \rho^5 \cos \phi \sin^3 \phi \cos \theta \sin \theta \, d\rho \, d\phi \, d\theta$$

(16) What is the value of  $\int_{-2}^2 \int_0^{\sqrt{4-y^2}} \int_{-\sqrt{4-x^2-y^2}}^{\sqrt{4-x^2-y^2}} y^2 \sqrt{x^2 + y^2 + z^2} \, dz \, dx \, dy$  ?

Answer:  $\frac{64\pi}{9}$ .



- (17) Evaluate  $\iiint_E \sqrt{x^2 + z^2} \, dV$ , where  $E$  is the region bounded by the paraboloid  $y = x^2 + z^2$  and the plane  $y = 4$ .  
Answer:  $\frac{128\pi}{15}$ .

- (18) Find the surface area of the part of the paraboloid  $z = x^2 + y^2$  that lies under the plane  $z = 9$ .  
Answer:  $\frac{\pi}{6}(37\sqrt{37} - 1)$ .

(19) Find  $K$ ,  $L$ ,  $M$ , and  $N$  so that

$$\int_0^2 \int_{1-(x/2)}^{1-(x^2/4)} f(x, y) dy dx = \int_K^L \int_M^N f(x, y) dx dy.$$

Answer:  $K = 0$ ,  $L = 1$ ,  $M = 2 - 2y$ ,  $N = 2\sqrt{1 - y}$ .

(20) Write the double integral in polar coordinates representing the area of the planar region bounded on the right by  $x^2 + y^2 = 2$  and bounded on the left by  $x = 1$ .

Answer: 
$$\int_{-\pi/4}^{\pi/4} \int_{\sec \theta}^{\sqrt{2}} r dr d\theta = \frac{\pi}{2} - 1.$$