

MA 23200 - Practice Exam 3

1. Compute  $AB-C$  where  $A = \begin{bmatrix} 1 & 1 \\ 0 & 3 \\ 5 & 2 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 1 \\ -2 & 0 \end{bmatrix}$ , and  $C = \begin{bmatrix} 1 & 0 \\ 3 & 2 \\ 6 & 4 \end{bmatrix}$

A.  $\begin{bmatrix} 0 & 1 \\ -9 & -2 \\ 5 & 1 \end{bmatrix}$

B.  $\begin{bmatrix} 0 & 1 \\ 9 & -2 \\ -5 & -1 \end{bmatrix}$

C.  $\begin{bmatrix} 0 & -1 \\ -9 & -2 \\ -5 & 1 \end{bmatrix}$

D.  $\begin{bmatrix} 0 & 1 \\ -9 & 2 \\ 5 & -1 \end{bmatrix}$

E.  $\begin{bmatrix} 0 & 1 \\ -9 & -2 \\ -5 & -1 \end{bmatrix}$

2. Solve the initial value problem

$$\begin{cases} f''(x) = e^{-\frac{1}{3}x} \\ f(0) = 1, f'(0) = 2 \end{cases}$$

A.  $-\frac{1}{9}e^{-\frac{1}{3}x} + 5x - 8$

B.  $9e^{-\frac{1}{3}x} + 5x - 8$


C.  $\frac{1}{9}e^{-\frac{1}{3}x} + 5x - 8$

D.  $\frac{1}{9}e^{-\frac{1}{3}x} + 5x + 8$

E.  $-9e^{-\frac{1}{3}x} + 5x - 8$

MA 23200 - Practice Exam 3

3. Which differential equation has the direction field shown below?



- A.  $\frac{dy}{dx} = 1 + x$
- B.  $\frac{dy}{dx} = x^2$
- C.  $\frac{dy}{dx} = \frac{x}{y}$
- D.  $\frac{dy}{dx} = \ln y$
- E.  $\frac{dy}{dx} = x + y$

4. Use Euler's method with  $\Delta x = .1$  to approximate  $y(0.2)$  if  $y'(x) = x + 2y$  and  $y(0) = 2$ .  
Euler's method:  $y_{n+1} = y_n + f(x_n, y_n)\Delta x$ , where  $y_n \approx y(x_n)$ .

- A. 2.40
- B. 3.09
- C. 2.89
- D. 2.68
- E. 3.49

MA 23200 - Practice Exam 3

5. Which best describes the stability of the equilibrium values of the autonomous differential equation

$$y' = -y^3 - y^2 + 6y.$$

- A. 2 stable and 1 unstable
- B. 1 stable, 1 semistable, and 1 unstable
- C. 2 unstable and 1 semistable
- D. 2 semistable and 1 unstable
- E. 1 stable and 2 unstable

6. Find the interval on which a unique solution exists for the initial value problem

$$(x + 3)y' - \frac{1}{x + 4}y = \frac{e^{4x}}{x - 2}; \quad y(1) = 3.$$

- A.  $(-4, 2)$
- B.  $(-\infty, 2)$
- C.  $(-3, \infty)$
- D.  $(-3, 2)$
- E.  $(-4, \infty)$

MA 23200 - Practice Exam 3

7. Find the integrating factor  $G(x)$  of the differential equation

$$(\cos x + 2)y' - (\sin x)y = 2.$$

- A.  $\cos x$
- B.  $\sin x + 2$
- C.  $\cos x - 2$
- D.  $\sin x$
- E.  $\cos x + 2$

8. Solve the initial value problem

$$\frac{dy}{dx} = \frac{(y^2 + 1) \sin x}{y^3 + y}; \quad y(0) = 2.$$

- A.  $y(x) = -\sqrt{-2 \cos x + 6}$
- B.  $y(x) = \sqrt{-2 \cos x - 6}$
- C.  $y(x) = \sqrt{-2 \cos x + 3}$
- D.  $y(x) = -\sqrt{-2 \cos x + 3}$
- E.  $y(x) = \sqrt{-2 \cos x + 6}$

MA 23200 - Practice Exam 3

9. The population of a city is increasing at a rate inversely proportional to the population at time  $t$ , where  $t$  is the number of years after 1993. If the population in 1993 ( $t = 0$ ) was 2 thousand, and in 1994 was 4 thousand, what was the population in 1998?

- A. 6 thousand
- B. 7 thousand
- C. 8 thousand
- D. 9 thousand
- E. 10 thousand

10. Given the initial value problem

$$y' - 2axy = -bx; \quad y(0) = \frac{b}{a},$$

where  $a$  and  $b$  are nonzero constants. The integrating factor  $G(x)$  of the differential equation  $y' - 2axy = -bx$  is  $G(x) = e^{-ax^2}$ . Solve for  $y(x)$ .

- A.  $y(x) = \frac{b}{2a} + \frac{b}{2a}e^{ax^2}$
- B.  $y(x) = -\frac{b}{2a} + \frac{b}{2a}e^{-ax^2}$
- C.  $y(x) = -\frac{b}{a} + \frac{b}{a}e^{ax^2}$
- D.  $y(x) = \frac{b}{a} + \frac{b}{a}e^{ax^2}$
- E.  $y(x) = \frac{b}{2a} + \frac{b}{a}e^{-ax^2}$


MA 23200 - Practice Exam 3

11. How many of the following differential equations are separable, linear or autonomous?

$$y' = ye^{-3x} + e^{-3x} \quad 2(y^2 + 1)y' = \ln y \quad y' + 3y = 1.$$

- A. 1 separable, 1 linear, and 1 autonomous
- B. 1 separable, 2 linear, and 2 autonomous
- C. 3 separable, 2 linear, and 1 autonomous
- D. 3 separable, 2 linear, and 2 autonomous
- E. 2 separable, 2 linear, and 2 autonomous

12. The Leslie diagram for a bird population is shown below



In Year 1, the hatchling population is 33 and the adult population is 18. Let  $a$  and  $b$  be the hatchling and adult population respectively in Year 3. Choose the correct statement.

- A.  $\begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 1.22 & 0.75 \\ 0.6 & 0.29 \end{bmatrix} \left( \begin{bmatrix} 1.22 & 0.75 \\ 0.6 & 0.29 \end{bmatrix} \begin{bmatrix} 33 \\ 18 \end{bmatrix} \right)$
- B.  $\begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0.75 & 0.29 \\ 0.6 & 1.22 \end{bmatrix} \left( \begin{bmatrix} 0.75 & 0.29 \\ 0.6 & 1.22 \end{bmatrix} \begin{bmatrix} 33 \\ 18 \end{bmatrix} \right)$
- C.  $\begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0.6 & 1.22 \\ 0.75 & 0.29 \end{bmatrix} \left( \begin{bmatrix} 0.6 & 1.22 \\ 0.75 & 0.29 \end{bmatrix} \begin{bmatrix} 33 \\ 18 \end{bmatrix} \right)$
- D.  $\begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 0.29 & 0.6 \\ 0.75 & 1.22 \end{bmatrix} \left( \begin{bmatrix} 0.29 & 0.6 \\ 0.75 & 1.22 \end{bmatrix} \begin{bmatrix} 33 \\ 18 \end{bmatrix} \right)$
- E.  $\begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 1.22 & 0.6 \\ 0.75 & 0.29 \end{bmatrix} \left( \begin{bmatrix} 1.22 & 0.6 \\ 0.75 & 0.29 \end{bmatrix} \begin{bmatrix} 33 \\ 18 \end{bmatrix} \right)$