

Study Guide # 2

1. Constrained extreme values via **Lagrange Multipliers**: Max/min -ize  $f(\mathbf{v})$  subject to constraint  $g(\mathbf{v}) = C$ , solve the system  $\nabla f = \lambda \nabla g$  and  $g(\mathbf{v}) = C$ .

2. Double integrals; Double Riemann sums:  $\iint_R f(x, y) dA \approx \sum_{i=1}^m \sum_{j=1}^n f(x_i^*, y_j^*) \Delta A$ ;

3. Type I region  $D : \begin{cases} g_1(x) \leq y \leq g_2(x) \\ a \leq x \leq b \end{cases}$  ; Type II region  $D : \begin{cases} h_1(y) \leq x \leq h_2(y) \\ c \leq y \leq d \end{cases}$  ;

iterated integrals over Type I and II regions:  $\iint_D f(x, y) dA = \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx$  and

$\iint_D f(x, y) dA = \int_c^d \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy$ , respectively; Reversing Order of Integration (regions that are both Type I and Type II); properties of double integrals.

4. Integral inequalities:  $mA \leq \iint_D f(x, y) dA \leq MA$ , where  $A = \text{area of } D$  and  $m \leq f(x, y) \leq M$  on  $D$ .

5. Polar:  $r^2 = x^2 + y^2$ ,  $x = r \cos \theta$ ,  $y = r \sin \theta$ ,  $\tan \theta = \frac{y}{x}$  (make sure  $\theta$  in correct quadrant).

Change of Variables Formula in Polar Coordinates: if  $D : \begin{cases} h_1(\theta) \leq r \leq h_2(\theta) \\ \alpha \leq \theta \leq \beta \end{cases}$ , then

$$\iint_D f(x, y) dA = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} f(r \cos \theta, r \sin \theta) \underset{\uparrow}{r} dr d\theta.$$

6. Applications of double integrals:

(a) Area of region  $D$  is  $A(D) = \iint_D dA$

(b) Volume of solid under graph of  $z = f(x, y)$ , where  $f(x, y) \geq 0$ , is  $V = \iint_D f(x, y) dA$

(c) Mass of  $D$  is  $m = \iint_D \rho(x, y) dA$ , where  $\rho(x, y) = \text{density (per unit area)}$ ; sometimes write  $m = \iint_D dm$ , where  $dm = \rho(x, y) dA$ .

(d) Moment about the  $x$ -axis  $M_x = \iint_D y \rho(x, y) dA$ ; moment about the  $y$ -axis  $M_y = \iint_D x \rho(x, y) dA$ .

(e) Center of mass  $(\bar{x}, \bar{y})$ , where  $\bar{x} = \frac{M_y}{m} = \frac{\iint_D x \rho(x, y) dA}{\iint_D \rho(x, y) dA}$ ,  $\bar{y} = \frac{M_x}{m} = \frac{\iint_D y \rho(x, y) dA}{\iint_D \rho(x, y) dA}$

(f) Surface Area  $A(S) = \iint_D \sqrt{1 + f_x^2 + f_y^2} dA$

**7.** Elementary solids  $E \subset \mathbb{R}^3$  of Type 1, Type 2, Type 3; triple integrals over solids  $E$ :

$$\iiint_E f(x, y, z) dV = \iint_D \int_{u(x,y)}^{v(x,y)} f(x, y, z) dz dA \text{ for } E = \{(x, y) \in D, u(x, y) \leq z \leq v(x, y)\};$$

volume of solid  $E$  is  $V(E) = \iiint_E dV$ ; applications of triple integrals, mass of a solid, moments about the coordinate planes  $M_{xy}, M_{xz}, M_{yz}$ , center of mass of a solid  $(\bar{x}, \bar{y}, \bar{z})$ .

**8. Cylindrical Coordinates**  $(r, \theta, z)$ :

$$\text{From CC to RC : } \begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases}$$

Going from RC to CC use  $x^2 + y^2 = r^2$  and  $\tan \theta = \frac{y}{x}$  (make sure  $\theta$  is in correct quadrant).

**9. Spherical Coordinates**  $(\rho, \theta, \phi)$ , where  $0 \leq \phi \leq \pi$ :

$$\text{From SC to RC : } \begin{cases} x = (\rho \sin \phi) \cos \theta \\ y = (\rho \sin \phi) \sin \theta \\ z = \rho \cos \phi \end{cases}$$

Going from RC to SC use  $x^2 + y^2 + z^2 = \rho^2$ ,  $\tan \theta = \frac{y}{x}$  and  $\cos \phi = \frac{z}{\rho}$ .

**10.** Triple integrals in Cylindrical Coordinates:  $\begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases}, \quad dV = r dz dr d\theta$

$$\iiint_E f(x, y, z) dV = \iiint_E f(r \cos \theta, r \sin \theta, z) r dz dr d\theta$$

↑

**11.** Triple integrals in Spherical Coordinates:  $\begin{cases} x = (\rho \sin \phi) \cos \theta \\ y = (\rho \sin \phi) \sin \theta \\ z = \rho \cos \phi \end{cases}, \quad dV = \rho^2 \sin \phi d\rho d\phi d\theta$

$$\iiint_E f(x, y, z) dV = \iiint_E f(\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi) \rho^2 \sin \phi d\rho d\phi d\theta$$

↑

**12.** Vector fields on  $\mathbb{R}^2$  and  $\mathbb{R}^3$  :  $\mathbf{F}(x, y) = \langle P(x, y), Q(x, y) \rangle$  and

$$\mathbf{F}(x, y, z) = \langle P(x, y, z), Q(x, y, z), R(x, y, z) \rangle;$$

$\mathbf{F}$  is a conservative vector field if  $\mathbf{F} = \nabla f$ , for some real-valued function  $f$  (potential).

**13.** Line integral of a function  $f(x, y)$  along  $C$ , parameterized by  $x = x(t)$ ,  $y = y(t)$  and  $a \leq t \leq b$ , is

$$\int_C f(x, y) ds = \int_a^b f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt.$$

(independent of orientation of  $C$ , other properties and applications of line integrals of  $f$ )

**Remarks:**

(a)  $\int_C f(x, y) ds$  is sometimes called the “line integral of  $f$  with respect to arc length”

(b) 
$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t)) x'(t) dt$$

(c) 
$$\int_C f(x, y) dy = \int_a^b f(x(t), y(t)) y'(t) dt$$

**14.** Line integral of vector field  $\mathbf{F}(x, y)$  along  $C$ , parameterized by  $\mathbf{r}(t)$  and  $a \leq t \leq b$ , is given by

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt.$$

(depends on orientation of  $C$ , other properties and applications of line integrals of  $\mathbf{F}$ )


**15.** Connection between line integral of vector fields and line integral of functions:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C (\mathbf{F} \cdot \mathbf{T}) ds$$

where  $\mathbf{T}$  is the unit tangent vector to the curve  $C$ .

**16.** If  $\mathbf{F}(x, y) = P(x, y) \mathbf{i} + Q(x, y) \mathbf{j}$ , then  $\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C P(x, y) dx + Q(x, y) dy$ ; Work =  $\int_C \mathbf{F} \cdot d\mathbf{r}$ .

**17.** FUNDAMENTAL THEOREM OF CALCULUS FOR LINE INTEGRALS:  $\int_C \nabla f \cdot d\mathbf{r} = f(\mathbf{r}(b)) - f(\mathbf{r}(a))$ :



**18.** A vector field  $\mathbf{F}(x, y) = P(x, y) \mathbf{i} + Q(x, y) \mathbf{j}$  is *conservative* (i.e.  $\mathbf{F} = \nabla f$ ) if  $\frac{\partial Q}{\partial x} = \frac{\partial P}{\partial y}$ ;  
how to determine a potential function  $f$  if  $\mathbf{F} = \nabla f$ .