


**Study Guide # 3****You also need Study Guides # 1 and # 2 for the Final Exam**

- 1.** GREEN'S THEOREM:  $\int_C P(x, y) dx + Q(x, y) dy = \iint_D \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$  ( $C$  = boundary of  $D$ ):




- 2.** DEL OPERATOR:  $\nabla = \frac{\partial}{\partial x} \vec{i} + \frac{\partial}{\partial y} \vec{j} + \frac{\partial}{\partial z} \vec{k}$ ; if  $\vec{F}(x, y, z) = P(x, y, z) \vec{i} + Q(x, y, z) \vec{j} + R(x, y, z) \vec{k}$ , then

$$\text{curl } \vec{F} = \nabla \times \vec{F} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix} \quad \text{and} \quad \text{div } \vec{F} = \nabla \cdot \vec{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

Properties of curl and divergence:

- (i) If  $\text{curl } \vec{F} = \vec{0}$ , then  $\vec{F}$  is a conservative vector field ( $\vec{F} = \nabla f$ ) in a simply-connected domain.
- (ii) If  $\text{curl } \vec{F} = \vec{0}$ , then  $\vec{F}$  is *irrotational*; if  $\text{div } \vec{F} = 0$ , then  $\vec{F}$  is *incompressible*.
- (iii) *Laplace's Equation:*  $\nabla^2 f = \text{div}(\nabla f) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = 0$ .
- (iv) For functions with continuous partials,  $\text{curl}(\nabla f) = \vec{0}$ , and  $\text{div}(\text{curl } \vec{F}) = 0$ .

- 3.** Parametric surface  $S$ :  $\vec{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle$ , where  $(u, v) \in D$ :



Normal vector to surface  $S$ :  $\vec{n} = \vec{r}_u \times \vec{r}_v$ ; tangent planes and normal lines to parametric surfaces.

**4.** Surface area of a surface  $S$ :

$$(i) \ A(S) = \iint_D |\vec{r}_u \times \vec{r}_v| dA$$

$$(ii) \text{ If } S \text{ is the graph of } z = h(x, y) \text{ above } D, \text{ then } A(S) = \iint_D \sqrt{1 + (\partial h / \partial x)^2 + (\partial h / \partial y)^2} dA;$$

Remark:  $dS = |\vec{r}_u \times \vec{r}_v| dA$  = differential of surface area; while  $d\vec{S} = (\vec{r}_u \times \vec{r}_v) dA$

**5.** The surface integral of  $f$  over the surface  $S$ :

$$(i) \ \iint_S f(x, y, z) dS = \iint_D f(\vec{r}(u, v)) |\vec{r}_u \times \vec{r}_v| dA.$$

$$(ii) \text{ If } S \text{ is the graph of } z = h(x, y) \text{ above } D, \text{ then}$$


$$\iint_S f(x, y, z) dS = \iint_D f(x, y, h(x, y)) \sqrt{1 + (\partial h / \partial x)^2 + (\partial h / \partial y)^2} dA.$$

**6.** The surface integral of  $\vec{F}$  over the surface  $S$  (recall,  $d\vec{S} = (\vec{r}_u \times \vec{r}_v) dA$ ):


$$\iint_S \vec{F} \cdot d\vec{S} = \iint_D \vec{F} \cdot (\vec{r}_u \times \vec{r}_v) dA = \iint_S (\vec{F} \cdot \vec{n}) dS = \text{flux of } \vec{F} \text{ across the surface } S.$$

If  $S$  is the graph of  $z = h(x, y)$  above  $D$ , with  $\vec{n}$  oriented upward, and  $\vec{F} = \langle P, Q, R \rangle$ , then

$$\iint_S \vec{F} \cdot d\vec{S} = \iint_D \left( -P \frac{\partial h}{\partial x} - Q \frac{\partial h}{\partial y} + R \right) dA$$




**7.** STOKES' THEOREM:  $\int_C \vec{F} \cdot d\vec{r} = \iint_S \text{curl } \vec{F} \cdot d\vec{S}$  (recall,  $\text{curl } \vec{F} = \nabla \times \vec{F}$ ).



$$\int_C \vec{F} \cdot d\vec{r} = \text{circulation of } \vec{F} \text{ around } C.$$

**8.** THE DIVERGENCE THEOREM/GAUSS' THEOREM:  $\iint_S \vec{F} \cdot d\vec{S} = \iiint_E \text{div } \vec{F} dV$

(recall,  $\text{div } \vec{F} = \nabla \cdot \vec{F}$ ).



## 9. Summary of Line Integrals and Surface Integrals:

LINE INTEGRALS	SURFACE INTEGRALS
$C : \vec{r}(t)$ , where $a \leq t \leq b$	$S : \vec{r}(u, v)$ , where $(u, v) \in D$
$ds =  \vec{r}'(t)  dt =$ differential of arc length	$dS =  \vec{r}_u \times \vec{r}_v  dA =$ differential of surface area
$\int_C ds =$ length of $C$	$\iint_S dS =$ surface area of $S$
$\int_C f(x, y, z) ds = \int_a^b f(\vec{r}(t))  \vec{r}'(t)  dt$ (independent of orientation of $C$ )	$\iint_S f(x, y, z) dS = \iint_D f(\vec{r}(u, v))  \vec{r}_u \times \vec{r}_v  dA$ (independent of normal vector $\vec{n}$ )
$d\vec{r} = \vec{r}'(t) dt$	$d\vec{S} = (\vec{r}_u \times \vec{r}_v) dA$
$\int_C \vec{F} \cdot d\vec{r} = \int_a^b \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt$ (depends on orientation of $C$ )	$\iint_S \vec{F} \cdot d\vec{S} = \iint_D \vec{F}(\vec{r}(u, v)) \cdot (\vec{r}_u \times \vec{r}_v) dA$ (depends on normal vector $\vec{n}$ )
$\int_C \vec{F} \cdot d\vec{r} = \int_C (\vec{F} \cdot \vec{T}) ds$ The <i>circulation</i> of $\vec{F}$ around $C$	$\iint_S \vec{F} \cdot d\vec{S} = \iint_S (\vec{F} \cdot \vec{n}) dS$ The <i>flux</i> of $\vec{F}$ across $S$ in direction $\vec{n}$

## 10. Integration Theorems:


FUNDAMENTAL THEOREM OF CALCULUS:  $\int_a^b F'(x) dx = F(b) - F(a)$




FUNDAMENTAL THEOREM OF CALCULUS FOR LINE INTEGRALS:  $\int_a^b \nabla f \cdot d\vec{r} = f(\vec{r}(b)) - f(\vec{r}(a))$



GREEN'S THEOREM:  $\iint_D \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA = \int_C P(x, y) dx + Q(x, y) dy$



STOKES' THEOREM:  $\iint_S \text{curl } \vec{F} \cdot d\vec{S} = \int_C \vec{F} \cdot d\vec{r}$



DIVERGENCE THEOREM:  $\iiint_E \text{div } \vec{F} dV = \iint_S \vec{F} \cdot d\vec{S}$

