Study Guide for Exam 2

- 1. You are supposed to know how to carry out the integration by parts
 - Indefinite form $\int u dv = uv \int v du$
 - Definite form $\int_a^b u dv = [uv]_a^b \int_a^b v du$

Example Problems

- 1.1. Compute the following integrals:
 - (i) $\int xe^x dx$ (ii) $\int e^x \sin x dx$ (iii) $\int \ln x \ dx$ (iv) $\int \sin^{-1} x \ dx$
- 1.2. Evaluate the following integrals
 - (i) $\int_{0}^{\frac{\pi}{6}} x \cos(2x) \, dx$ (ii) $\int_{0}^{1/2} \sin^{-1} x \, dx$ (iii) $\int_{0}^{1} \tan^{-1} x \, dx$
- 1.3. Find the volume of the solid obtained by rotating the following region about the y-axis: the region is bounded by $y = f(x) = \sin x$ and the x-axis on the interval $[0, \pi]$.
- 2. You are supposed to know how to comopute the integration of the form
 - $(1) \int \sin^m x \cos^n x \ dx$
 - Case: $m \text{ odd} \rightarrow \text{Use } u = \cos x \text{ substitution}$
 - Case: $n \text{ odd} \rightarrow \text{Use } u = \sin x \text{ substitution}$
 - \bullet Case: $m\ \&\ n$ even \rightarrow Reduce the degree by double angle formula
 - (2) $\int \tan^m x \sec^n x \ dx$
 - Case: n > 0 even \rightarrow Use $u = \tan x$ substitution
 - Case: n > 0 odd & m odd \rightarrow Use $u = \sec x$ substitution
 - Case: n > 0 & m even \rightarrow Integration by parts
 - Case: $n = 0 \to \text{Use } \tan^2 x = \sec^2 x 1$ to reduce to

the case n > 0 and to the lower degree case

Example Problems

2.1. Compute the following integrals:

(i)
$$\int \sin^3 x \cos^2 x \, dx$$

(ii)
$$\int \sin^2 x \cos^3 x \, dx$$

(iii)
$$\int \sin^3 x \cos^3 x \, dx$$

(iv)
$$\int \sin^4 x \cos^2 x \, dx$$

2.2. Compute the following integrals:

(i)
$$\int \tan^2 x \sec^4 x \, dx$$
(ii)
$$\int \sec^3 \tan x \, dx$$
(iii)
$$\int \sec^3 x \, dx.$$
(iii)
$$\int \tan^2 x \sec x \, dx.$$
(iii)
$$\int \tan^5 x \, dx.$$

2.3.

(i) Compute
$$\int \sec x \, dx = \int \frac{1}{\cos^2 x} \cos x \, dx = \int \frac{1}{1 - \sin^2 x} \cos x \, dx$$
 using the substitution $u = \sin x$ and then using the partial fraction.

(ii) Check that the result obtained in (i) coincides with the well-known formula $\int \sec x \, dx = \ln|\sec x + \tan x| + C$.

2.4.

We would like to compute

$$\int \tan x \ dx = \int \frac{\sin x}{\cos x} \ dx$$

in the following two ways.

(i) Use substitution $u = \sin x$ to get

$$\int \tan x \, dx = \int \frac{\sin x}{\cos x} \, dx = \int \frac{\sin x}{\cos^2 x} \cos x dx$$
$$= \int \frac{\sin x}{1 - \sin^2 x} \cos x dx = \int \frac{u}{1 - u^2} \, du$$

and then use the partial fractions.

(ii) Use the same substitution as above and then use another substitution $v = 1 - u^2$ to get

$$\int \tan x \ dx = \int \frac{u}{1 - u^2} \ du = \int -\frac{1}{2} \frac{dv}{v}$$

and compute.

- (iii) Check that the results obtained in (i) and (ii) coincide with the well known formula $\int \tan x \ dx = \ln|\sec x| + C$.
- 3. You are supposed to know how to use the 3 types of trigonometric substitution, and carry out the integration accordingly.
 - (1) $\sqrt{a^2 x^2}$, $x = a \sin \theta$, $dx = a \cos \theta$, $\sqrt{a^2 x^2} = a \cos \theta$,
 - (2) $\sqrt{a^2 + x^2}$, $x = a \tan \theta$, $dx = a \sec^2 \theta$, $\sqrt{a^2 + x^2} = a \sec \theta$,
 - (3) $\sqrt{x^2 a^2}$, $x = a \sec \theta$, $dx = a \tan \theta \sec \theta$, $\sqrt{x^2 a^2} = a \tan \theta$.

Example Problems

3.1. Compute the following integrals:

(i)
$$\int \frac{dx}{\sqrt{4 - x^2}}$$
(ii)
$$\int \sqrt{5 - 4x^2} \, dx$$
(iii)
$$\int_{3}^{6} \frac{\sqrt{x^2 - 9}}{x} \, dx$$
(iv)
$$\int \sqrt{x^2 + 1} \, dx$$
(v)
$$\int_{1}^{\sqrt{3}} \frac{dx}{x^2 \sqrt{x^2 + 1}}$$
(vi)
$$\int \frac{x}{\sqrt{3 - 2x - x^2}} dx$$
(vii)
$$\int \frac{x^2 - 2x + 2}{\sqrt{x^2 - 2x + 10}} dx$$
(viii)
$$\int \frac{x}{\sqrt{3 + 2x - x^2}} dx$$

- 3.2. Verify that the area of a circle of radius r is πr^2 .
- 3. You are supposed to know
 - the proper form of the partial fractions,
- how to determine the appropariate constants appearing in the partial fraction,
 - o how to compute the integral accordingly.

Example Problems

3.1. Determine the proper form of th partial fractions for the following. (You do not have to calculate the constants.)

(i)
$$\frac{1}{(x+2)(x^2-4)((x^2+x+1)^2)}$$
(ii)
$$\frac{x^3}{(x-1)(x^3-1)(x^2+4x+5)}$$

3.2. Compute the following integrals:

(i)
$$\int \frac{x^2}{(x-1)^2} dx$$
(ii)
$$\int \frac{x+2}{x^2+2x+2} dx$$
(iii)
$$\int \frac{x}{(x+1)(x-1)(x-2)} dx$$
(iv)
$$\int \frac{x^2}{(x-1)^2(x^2+1)} dx$$
(v)
$$\int \frac{x^2+x+2}{x^2+4x+5} dx$$

4. You are supposed to know why a given improper integral is improper, and accordingly to be able to determine if the given improper integral is convergent/divergent. In case it is convergent, you should be able to compute its value.

Example Problems

4.1. Evaluate the following improper integrals

(i)
$$\int_0^\infty \frac{e^x}{e^{2x} + 1} dx$$
(ii)
$$\int_0^\infty \frac{e^{2x}}{e^{2x} + 1} dx$$
(iii)
$$\int_0^9 \frac{1}{x - 1} dx$$
(iv)
$$\int_0^9 \frac{1}{\sqrt[3]{x - 1}} dx$$
(v)
$$\int_0^\infty x dx$$
(vi)
$$\int_0^\infty xe^{-x} dx$$
(vi)
$$\int_0^\infty xe^{-x^2} dx$$

5. You are supposed to be able to determine if a given sequence is convergent/divergent. In case it is convergent, you should be able to compute its limit.

Example Problems

(i)
$$a_n = \frac{(-1)^n n}{n^2 + 1}$$

(ii) $a_n = \tan^{-1} \left(\frac{n^3 + 5}{n^2}\right)$
(iii) $a_n = (-1)^n \sin\left(\frac{\pi}{2} - \frac{1}{n}\right)$
(iv) $a_n = \cos(n\pi)$
(v) $a_n = \sin(n\pi)$
(vi) $a_n = \frac{n!n^2}{(n+2)!} \cos(1/2n)$
(vii) $a_n = n \tan(1/n)$