

Name _____ 10-digit PUID _____

Recitation Instructor _____ Recitation Section Number _____

Lecture's Name _____

Instructions:

1. Academic Integrity

- a. Students may not open the exam until instructed to do so.
- b. Students must obey the orders and requests by all proctors, TAs, and lecturers.
- c. No student may leave in the first 20 minutes or in the last 10 minutes of the exam.
- d. Books, notes, calculators, or any electronic devices are not allowed on the exam, and they should not even be in sight in the exam room. Students may not look at anybody else's test, and may not communicate with anybody else except, if they have a question, with their TA or lecturer.
- e. After time is called, the students have to put down all writing instruments and remain in their seats, while the TAs will collect the scantrons and the exams.
- f. Any violation of these rules and any act of academic dishonesty may result in severe penalties. Additionally, all violators will be reported to the Office of the Dean of Students.

I have read and understand the exam rules stated above:

STUDENT NAME (print): _____

STUDENT SIGNATURE: _____

2. Fill in all the information requested above. On the scantron sheet fill in your name, student ID number, and the section number of your recitation with an extra 0 at the left. Blacken the correct circles.
3. This booklet contains 12 problems. Problems 1 through 4 are worth 9 points each. Problems 5 through 12 are worth 8 points each. The maximum score is 100 points.
4. For each problem mark your answer on the scantron sheet and also circle it in this booklet.
5. Work only on the pages of this booklet.

Mark **TEST 01** on your scantron!

1. Find the area between the curves $y = 3 - x^2$ and $y = 2x$.

A. $\frac{14}{3}$

B. $\frac{32}{3}$

C. 9

D. $\frac{34}{3}$

E. $\frac{35}{3}$

2. The triangular region with vertices $(0, 0)$, $(2, 0)$ and $(2, 4)$ is rotated about the line $x = 0$. Using the method of disks (or washers), find the volume of the resulting solid.

A. $\int_0^4 \pi \left(\frac{1}{2}y\right)^2 dy$

B. $\int_0^4 \pi \left(\left(\frac{1}{2}y\right)^2 - 2^2\right) dy$

C. $\int_0^4 \pi \left(4^2 - \left(\frac{1}{2}y\right)^2\right) dy$

D. $\int_0^4 \pi \left(2^2 - \left(\frac{1}{2}y\right)^2\right) dy$

E. $\int_0^2 \pi \left(2^2 - \left(\frac{1}{2}y\right)^2\right) dy$

3. The triangular region with vertices $(0,0)$, $(2,2)$ and $(2,4)$ is rotated about the line $x = 2$. Using the method of cylindrical shells, find the volume of the resulting solid.

A. $\int_0^4 2\pi x(2x) dx$

B. $\int_0^2 2\pi x(4 - 2x) dx$

C. $\int_0^2 2\pi(2 - x)(x) dx$

D. $\int_0^2 2\pi(2 - x)(4x) dx$

E. $\int_0^2 2\pi x(2x) dx$

4. If 3 ft-lbs is required to stretch a spring $\frac{1}{2}$ foot beyond its natural length, how much work is required to stretch the spring $\frac{1}{3}$ foot beyond its natural length?

A. $\frac{1}{6}$ ft-lbs

B. 2 ft-lbs

C. $\frac{4}{3}$ ft-lbs

D. $\frac{3}{4}$ ft-lbs

E. $\frac{5}{3}$ ft-lbs

5. $\int_1^3 x \ln x \, dx =$

A. $\frac{9}{2} \ln 3 - 2$

B. $\frac{9}{2} \ln 3 - \frac{5}{2}$

C. $\frac{9}{4} \ln 3 - 2$

D. $\frac{9}{2} \ln 3$

E. $\frac{9}{4} \ln 3 - \frac{5}{4}$

6. Find the equation of a sphere if one of its diameters has endpoints $(2, 3, 0)$ and $(0, 1, -2)$.

A. $x^2 + y^2 + z^2 + x + 2y - z = 3$

B. $x^2 + y^2 + z^2 - 2x - 4y + 2z = -3$

C. $x^2 + y^2 + z^2 + 2x + 4y - 2z = 3$

D. $x^2 + y^2 + z^2 + x + 2y - z = 6$

E. $x^2 + y^2 + z^2 - x + 2y - z = -3$

7. Find the angle between $\langle 4, 1, 1 \rangle$ and $\langle 2, 2, -1 \rangle$.

- A. $\frac{\pi}{6}$
- B. $\frac{\pi}{3}$
- C. $\frac{2\pi}{3}$
- D. $\frac{5\pi}{6}$
- E. $\frac{\pi}{4}$

8. Find a vector that is perpendicular to the plane which contains the points $P(1, 2, 2)$, $Q(3, 1, -1)$ and $R(3, 3, 1)$.

- A. $\langle 4, -4, 1 \rangle$
- B. $\langle 1, 1, 1 \rangle$
- C. $\langle 4, -4, -1 \rangle$
- D. $\langle -2, -4, 4 \rangle$
- E. $\langle 1, -1, 1 \rangle$

9. If $\vec{v} = 2\vec{i} - \vec{j}$ and $\vec{w} = \vec{i} + \vec{j}$, find $\text{proj}_{\vec{v}}\vec{w}$.

A. $\left\langle \frac{2}{\sqrt{5}}, \frac{-1}{\sqrt{5}} \right\rangle$

B. $\left\langle \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right\rangle$

C. $\left\langle \frac{2}{5}, \frac{-1}{5} \right\rangle$

D. $\left\langle \frac{1}{2}, \frac{1}{2} \right\rangle$

E. $\left\langle \frac{-2}{5}, \frac{1}{5} \right\rangle$

10 Find $\vec{a} \cdot \vec{b}$, where $|\vec{a}| = 8$, $|\vec{b}| = 3$ and the angle between \vec{a} and \vec{b} is $\frac{2\pi}{3}$.

A. $12\sqrt{3}$

B. $-12\sqrt{3}$

C. 12

D. -12

E. $6\sqrt{2}$

11. Find the volume of the solid obtained by rotating the region bounded by $x = 2\sqrt{3y}$, $x = 0$ and $y = 5$ about the y -axis.

A. 150π

B. 75π

C. $\frac{25\sqrt{3}\pi}{2}$

D. $10\sqrt{3}\pi$

E. 300π

12. Find all values of c such that $f(c)$ is the average value of $f(x) = (x - 3)^2$ on the interval $[2, 5]$.

A. $c = 1$

B. $c = 3$

C. $c = 1, 3$

D. $c = 2, 4$

E. $c = 4$