

PROBLEM OF THE WEEK  
Solution of Problem No. 14 (Fall 2001 Series)

**Problem:** Consider the motion of a mass in the  $(x, y)$  plane. Given that the angular velocity  $\omega = x\dot{y} - y\dot{x}$  ( $\cdot$  is  $\frac{d}{dt}$ ) and the Lenz vector  $(\ell_x, \ell_y)$ , where  $\ell_x = -\frac{\omega}{k}\dot{y} + \frac{x}{r}$ ,  $\ell_y = \frac{\omega}{k}\dot{x} + \frac{y}{r}$  ( $r = (x^2 + y^2)^{1/2}$ ) are constant (independent of  $t$ ), show that the acceleration vector points toward the point where  $r = 0$  and its magnitude is inversely proportional to  $r^2$ .

**Solution** (by Mike Hamburg, Sr. St. Joseph H.S., South Bend, edited by the Panel)

First we note that  $\dot{r} = \frac{x\dot{x} + y\dot{y}}{r}$ . Now, we have

$$\begin{aligned}(\dot{\ell}_x, \dot{\ell}_y) &= \left(-\frac{\omega}{k}\ddot{y} + \frac{\dot{x}r - \frac{x(x\ddot{x} + y\dot{y})}{r}}{r^2}, \quad \frac{\omega}{k}\ddot{x} + \frac{\dot{y}r - \frac{y(x\dot{x} + y\dot{y})}{r}}{r^2}\right) \\&= \left(-\frac{\omega}{k}\ddot{y} + \frac{\dot{x}y^2 - xy\dot{y}}{r^3}, \quad \frac{\omega}{k}\ddot{x} + \frac{x^2\dot{y} - xy\dot{x}}{r^3}\right) \\&= \left(-\frac{\omega}{k}\ddot{y} - \frac{y\omega}{r^3}, \quad \frac{\omega}{k}\ddot{x} + \frac{x\omega}{r^3}\right) \\&= \omega\left(-\frac{\ddot{y}}{k} - \frac{y}{r^3}, \quad \frac{\ddot{x}}{k} + \frac{x}{r^3}\right).\end{aligned}$$

If  $\omega \neq 0$ , since  $(\dot{\ell}_x, \dot{\ell}_y) = 0$ , it follows that the acceleration

$$(\ddot{x}, \ddot{y}) = -\frac{k}{r^2}\left(\frac{x}{r}, \frac{y}{r}\right),$$

which is parallel to  $(x, y)$ , points toward the origin as long as  $k > 0$ , and has magnitude  $\frac{|k|}{r^2}$  as required.

Mike Hamburg observes that the conclusion does not hold if  $\omega = 0$ .

Also solved by:

Undergraduates: Eric Tkaczyk (Jr. EE/MA)

Faculty: Steven Landy (Phys. at IUPUI)