

Gap References

Mark Ashbaugh
Department of Mathematics
University of Missouri
Columbia, MO 65211
mark@math.missouri.edu

Rodrigo Bañuelos
Department of Mathematics
Purdue University
West Lafayette, IN 47906
banuelos@math.purdue.edu

References

- [1] S. Abramovich, *On frequencies of strings and deformations of beams*, Quart. Appl. Math. **63** (2005), 291–299.
- [2] S. Abramovich, *The gap between the first two eigenvalues of a one-dimensional Schrödinger operator with symmetrical potential*, Proc. Amer. Math. Soc. **111** (1991), 451–453.
- [3] M. S. Ashbaugh, “Open problems” section of “Some eigenvalue comparison results for domains in $\mathbb{R}^n$ and for annular domains in $\mathbb{R}^n$ ”, pp. 2–11 of **ESI - Workshop on Geometrical Aspects of Spectral Theory, Matrei in East Tyrolia, Austria July 5–12, 1999**, L. Friedlander and T. Hoffmann–Ostenhof, editors, published electronically as preprint #768 at <http://www.esi.ac.at/ESI-Preprints.html> (October 12, 1999).
- [4] M. Ashbaugh and R. Benguria, *Best constant for the ratio of the first two eigenvalues of one-dimensional Schrödinger operators with potentials*, Proc. Amer. Math. Soc., **99** (1987), 598–599.
- [5] M. Ashbaugh and R. Benguria, *Optimal lower bounds for eigenvalue gaps for Schrödinger operators with symmetric single well potentials and related results*, Maximum principles and eigenvalue problems in partial differential equations. Longman, White Plains, NY (1988).
- [6] M. Ashbaugh and R. Benguria, *Optimal lower bounds for the gap between the first two eigenvalues of one-dimensional Schrödinger operators with symmetric single-well potentials*, Proc. Amer. Math. Soc., **105** (1989), 419–424.
- [7] M. S. Ashbaugh and R. Benguria, *A sharp bound for the ratio of the first two eigenvalues of Dirichlet Laplacians and extensions*, Ann. Math. **135**(2) (1992), 601–628.
- [8] C. Bandle, *Isoperimetric Inequalities and Applications*, Monographs and Studies in Mathematics, Pitman 1980.
- [9] R. Bañuelos, *Intrinsic ultracontractivity and eigenfunction estimates for Schrödinger operators*, J. Funct. Anal. **100** (1991), 181–206.
- [10] R. Bañuelos and T. Carroll, *Brownian motion and the fundamental frequency of a drum*, Duke Math. J. **75** (1994), no. 3, 575–602.

- [11] R. Bañuelos and K. Burdzy, *On the “hot spots” conjecture of J. Rauch*, *J. Funct. Anal.* 164 (1999), 1–33.
- [12] R. Bañuelos, R. Latała, P. J. Méndez-Hernández, *A Brascamp-Lieb-Luttinger-type inequality and applications to symmetric stable processes*, *Proc. Amer. Math. Soc.* 129(10) (2001), 2997–3008.
- [13] R. Bañuelos and P.J. Méndez-Hernández, *Sharp inequalities for heat kernels of Schrödinger operators and applications to spectral gaps*, *J. Funct. Anal.* 176(2) (2000), 368–399.
- [14] R. Bañuelos and P.J. Méndez-Hernández, *Hot-Spots for Conditioned Brownian Motion*, *Illinois J. Math*, (**to appear**)
- [15] R. Bañuelos and P. Kröger, *Gradient estimates for the ground state Schrödinger eigenfunction and applications*, *Comm. Math. Phys.*, 224(2) 2001, 545–550.
- [16] R. Bañuelos and T.Kulczycki, *Eigenvalue gaps for the Cauchy process and a Poincaré inequality* (**preprint**).
- [17] R. Bañuelos and T.Kulczycki, *Spectral gap for the Cauchy process on convex symmetric domains*, (**preprint**).
- [18] M. van den Berg, *On condensation in the free-boson gas and the spectrum of the Laplacian*, *J. Statist. Phys.* , 31 (1983), 623–637.
- [19] D. Betsakos, *Symmetrization, symmetric stable processes, and Riesz capacities*, *Trans. Amer. Math. Soc.* 356 (2004), no. 2, 735–755
- [20] D. Betsakos, *Polarization, continuous Markov processes, and second order elliptic equations*, *Indiana Univ. Math. J.* 53 (2004), no. 2, 331–345.
- [21] S. G. Bobkov, *Isoperimetric and Analytic Inequalities for Log-Concave Probability Measures*, *Ann. Probab.* 27, no. 4 (1999), 1903–1921.
- [22] S. G. Bobkov and M. Ledoux, *Poincaré’s inequalities and Talagrand’s concentration phenomenon for the exponential distribution*, *Probab. Theory Related Fields* 107 (1997), no. 3, 383–400
- [23] C. Borell, *Diffusion equations and geometric inequalities*, *Potential Anal.* 12 (2000), 49–71.

- [24] H.J. Brascamp, E. H. Lieb and J.M. Luttinger, A General Rearrangement Inequality for Multiple Integrals, *Jour. Funct. Anal.* **17** (1975), 227-237
- [25] H. J. Brascamp and E. H. Lieb, *Some inequalities for Gaussian measures*, **Functional Integration and Its Applications**, A. M. Arthurs, editor, Oxford University Press, London, 1975, pp. 1-14.
- [26] H.J. Brascamp and E.H. Lieb, *On extensions of the Brunn-Minkowski and Prékopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation*, *J. Functional Analysis*, **22** (1976), 366-389.
- [27] R. M. Brown, P. D. Hislop, and A. Martinez, *Lower bounds on the interaction between cavities connected by a thin tube*, *Duke Math. J.* **73** (1994), 163-176.
- [28] A. Burchard and M. Schmuckenschläger, *Comparison theorems for exit times*, *Geom. Funct. Anal.*, **11** (2001), 651-692.
- [29] K. Burdzy and W. Werner, *A counterexample to the "hot spots" conjecture*. *Ann. Math.* **149** (1999), 309-317.
- [30] R. Carmona, W. C. Masters and B. Simon, *Relativistic Schrödinger operators: asymptotic behavior of eigenfunctions*, *J. Funct. Anal.*, **91** (1990), 117-142.
- [31] J. Cheeger, *A lower bound for the smallest eigenvalue of the Laplacian*, in Gunning (ed.), *Problems in Analysis*, Princeton University Press. (1970), 195-199.
- [32] S.Y. Cheng, *Eigenvalue comparison theorems and its geometric applications*, *Math. Z.* **143** (1975), 289-297.
- [33] R. Chen, *Gap estimates of Schrödinger operator*, *Pac. J. Math.* **178** (1997), 225-240.
- [34] R. Chen, *Neumann eigenvalue estimate on a compact Riemannian manifold*, *Proc. Amer. Math. Soc.* **108** (1990), 961-970.
- [35] M. F. Chen and F.-Y. Wang, *General formula for lower bound of the first eigenvalue on Riemannian manifolds*, *Sci. Sin. (A)* **40** (1997), 384-394.

- [36] S.Y. Cheng and K. Oden, *Isoperimetric inequalities and the gap between the first and second eigenvalues of an Euclidean domain*, . Geom. Anal. 7 (1997), no. 2, 217–239.
- [37] F. R.K. Chung, *Spectral Graph Theory*, AMS Publications, 1997.
- [38] F. R.K. Chung and K. Oden, *Weighted graph Laplacians and isoperimetric inequalities*, Pacific J. Math. 192 (2000), 257–273.
- bibitemDa84b E. B. Davies, *Double well Hamiltonians*, Proc. London Math. Soc. **49** (1984), 193–206.
- [39] E.B. Davies, *Structural isomers, double wells, resonances an Dirichlet decoupling*, Ann. Phys., 157 (1984), 166–182.
- [40] E.B. Davies, *Heat Kernels and Spectral Theory*, Cambridge University Press, Cambridge, (1989).
- [41] E.B. Davies and M. Pang, *The eigenvalue gap for second-order elliptic operators with Dirichlet boundary conditions*, J. Differential Equations 88 (1990), no. 1, 46–70.
- [42] B. Davis, *On the spectral gap for fixed membranes*, Ark. Mat. 39(1) (2001), 65–74.
- [43] B. Davis and M. Hosseini, *On ratio inequalities for heat content*, J. London Math. Soc. (2) 69 (2004), no. 1, 97–106.
- [44] P. Diaconis and D.W. Stroock, *Geometric bounds for eigenvalues of Markov Chains*, Snnals of Applied Prob., 1 (1991), 36–61.
- [45] P. Diaconis and L. Saloff-Coste, *Comparison theorems for reversible Markov chains*, Annals of Prob. 3 (1993), 696–730.
- [46] J.L. Doob, *Conditioned Brownian motion and the boundary limits of harmonic functions*, Bull. Soc. Math. France, 85 (1957), 431–458.
- [47] C. Draghici, *Rearrangement inequalities with application to ratios of heat kernels*, Potential Anal., 22 (2005), 351–374.
- [48] C. Draghici, *A general rearrangement inequality*, Proc. Amer. Math. Soc. 133 (2005), no. 3, 735–743.
- [49] C. Draghici, *Inequalities for integral means over symmetric sets*, (2005 preprint).

- [50] J.A. Fill, *Eigenvalue bounds on convergence to stationary for non-reversible Markov chains, with applications to the exclusion process*, *Annals of Applied Probability*. **1** (1991), 62–87.
- [51] E. Harrell, *Double wells*, *Comm. Math. Phys.*, **75** (1980), 239–261.
- [52] E. Harrell, *On the rate of asymptotic eigenvalue degeneracy*, *Comm. Math. Phys.*, **60** (1978), 73–95.
- [53] A. Henrot, **Extremum Problems for Eigenvalues of Elliptic Operators**, 2006 (to appear)
- [54] M. Hildebrand, *Rates of Convergence for non-reversible Markov chain sampler* (**preprint**)
- [55] M. Horvath, *On the first two eigenvalues of Sturm-Liouville operators*, *Proc. Amer. Math. Soc.* **131** (2003), 1215–1224.
- [56] H.-K. Huang, *Optimal estimates of the eigenvalue gap and eigenvalue ratio with variational analysis*, Masters thesis (unpublished), Department of Applied Mathematics, National Sun Yat-sen University, Kaohsiung, Taiwan, Republic of China, May, 2004 (51 pp.).
- [57] M.-J. Huang, *On the eigenvalue ratio for vibrating strings*, *Proc. Amer. Math. Soc.* **127** (1999), 1805–1813.
- [58] W. Kirsch, *Estimates on the difference between succeeding eigenvalues and Lifshitz tails for random Schrödinger operators*, **Stochastic Processes—Mathematics and Physics, II (Bielefeld, 1985)**, *Lect. Notes in Math.* **1250**, Springer, Berlin, 1987, pp. 138–151.
- [59] W. Kirsch, *On the difference between eigenvalues of Sturm-Liouville operators and the semiclassical limit*, **Differential Equations and Mathematical Physics (Birmingham, Alabama, 1986)**, *Lect. Notes in Math.* **1285**, Springer, Berlin, 1987, pp. 230–235.
- [60] W. Kirsch and B. Simon, *Universal lower bounds on eigenvalue splittings for one-dimensional Schrödinger operators*, *Comm. Math. Phys.*, **97** (1985), 435–460.
- [61] W. Kirsch and B. Simon, *Comparison theorems for the gap of Schrödinger operators*, *J. Funct. Anal.*, **75** (1986), 337–340.
- [62] M. Kiss, *Eigenvalue ratios of vibrating strings*, *Acta Math. Hungar.* **110** (2006), 253–259.

- [63] R. Lavine, *The eigenvalue gap for one dimensional convex potentials*, *Proc. Amer. Math. Soc.*, 121 (1994), 815–821.
- [64] Y. I. Lee and A. N. Wang, *Estimate of $\lambda_2 - \lambda_1$ on spheres*, *Chinese J. Math.* **15** (1987), 95–97.
- [65] J. Ling, *A lower bound for the gap between the first two eigenvalues of Schrödinger operators on convex domains in S^n and $\mathbb{R}^n$* , *Michigan Math. J.*, 40 (1993), 259–270.
- [66] J. Ling, *Estimates on the lower bound of the first gap*, 2005 preprint (available at the Los Alamos archive).
- [67] S. Nakamura, *A remark on eigenvalue splittings for one-dimensional double-well Hamiltonians*, *Lett. Math. Phys.* **11** (1986), 337–340.
- [68] A.D. Melas, *On the nodal line of the second eigenfunction of the Laplacian in $\mathbb{R}^2$* , *J. Differential Geom.*, 35 (1992), 255–266.
- [69] P.J. Méndez-Hernández, *Brascamp-Lieb-Luttinger Inequalities for Convex Domains of Finite Inradius*, *Duke Math. J.*, 13 (2002), 93–131
- [70] K. Oden, C.J. Sung and J. Wang, *Spectral gap estimates on Compact manifolds*, *Trans. Amer. Math. Soc.* 351 (1999), 3533–3548.
- [71] L. E. Payne, *Isoperimetric inequalities and their applications*, *SIAM Rev.*, 9 (1967), 453–488.
- [72] L. Payne, *On two conjectures in the fixed membrane eigenvalue problem*, *J. Appl. Math. Phys. (ZAMP)*, 24 (1973), 721–728.
- [73] L. E. Payne and H. F. Weinberger, *An optimal Poincaré inequality for convex domains*, *Arch. Rational Mech. Anal.*, 5 (1960), 286–292.
- [74] R. G. Pinsky, *Comparison theorems for the spectral gap of diffusions on an interval and their connection to corresponding results for one-dimensional Schrödinger operators*, this paper can be found on the web at <http://www.math.technion.ac.il/~pinsky/mypapers/speccom.ps> (15 pp.).
- [75] R. G. Pinsky, *Spectral gap and rate of convergence to equilibrium for a class of conditioned Brownian motions*, *Stochastic Process. Appl.* **115** (2005), 875–889. [This paper can also be found on the web at

<http://www.math.technion.ac.il/~pinsky/mypapers/specgap.ps> (18 pp.).

- [76] R. G. Pinsky, *Comparison theorems for the spectral gap of diffusion processes and Schrödinger operators on an interval*, J. London Math. Soc. (2) **72** (2005), 621–631.
- [77] M. H. Protter, *Lower bounds for the spectrum of second order operators and systems of operators*, **Maximum Principles and Eigenvalue Problems in Partial Differential Equations**, P. W. Schaefer, editor, Pitman Research Notes in Mathematics Series, vol. **175**, Longman Scientific and Technical, Harlow, Essex, United Kingdom, 1988, pp. 82–93.
- [78] I. M. Singer, B. Wong, S.-T. Yau, S. S.-T. Yau, *An estimate of the gap of the first two eigenvalues in the Schrödinger operator*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **12**(2) (1985), 319–333.
- [79] L. Saloff-Coste, *Precise estimates on the rate at which certain diffusions tend to equilibrium*, Math. Z. **217** (1994), no. 4, 641–677.
- [80] R. Smits, *Spectral gaps and rates to equilibrium for diffusions in convex domains*, Michigan Math. J., **43** (1996), 141–157.
- [81] F.-Y. Wang, *On estimation of the Dirichlet spectral gap*, Arch. Math. **75** (2000), 450–455.
- [82] S.-T. Yau, **Nonlinear Analysis in Geometry**, Enseignement Math., Geneva, 1986 (this also appeared as an article *Nonlinear analysis in geometry*, Enseign. Math. **33** (1987), 109–158).
- [83] S.-T. Yau, *Open problems in geometry*, in “Chern—a great geometer of the twentieth century,” Edited by S.-T. Yau, International Press, Hong Kong, 1992.
- [84] D. You, *Sharp inequalities for ratios and partition functions of Schrödinger operators*, Potential Anal. **18** (2003), no. 3, 219–250.
- [85] Q.-H. Yu and J.-Q. Zhong, *Lower bounds of the gap between the first and second eigenvalues of the Schrödinger operator*, Trans. Amer. Math. Soc., **294** (1986), 341–349.
- [86] W. K. Yuen, *Applications of geometric bounds to the convergence rate of Markov chains on R^n* , Stochastic Process. Appl. **87** (2000), no. 1, 1–23.