

MA 528 Exam 1 Spring 2016 Name & Section _____

Show your work to get credit. Box your answers.

1. (20 pts) For the surface given by $z = x^3 + y^3$, find a downward pointing normal vector at $(1, 2, 10)$ and an equation for the tangent plane at that point.

$$\varphi(x, y, z) = z - x^3 - y^3 - 1$$

$$\vec{N} = \nabla \varphi \Big|_{(1, 2, 10)} = -3x^2 \hat{i} - 3y^2 \hat{j} + \hat{k} \Big|_{(1, 2, 10)} = -3\hat{i} - 12\hat{j} + \hat{k}$$

Downward normal : $\vec{N} = 3\hat{i} + 12\hat{j} - \hat{k}$

Equation of tangent plane

$$3(x-1) + 12(y-2) - (z-10) = 0$$

2. (20 pts) Find a potential function for the curl free vector field

$$\vec{F} = \left(\frac{2xy}{1+x^2} \right) \hat{i} + (2yz + \ln(1+x^2)) \hat{j} + (1+y^2) \hat{k}$$

$$(A): \frac{\partial \varphi}{\partial x} = \frac{2xy}{(1+x^2)} \quad \varphi = \int \underbrace{\frac{2xy}{(1+x^2)}}_u dx = y \ln(1+x^2) + g(y, z)$$

$$(B): \frac{\partial \varphi}{\partial y} = \frac{2}{2y} \left[y \ln(1+x^2) + g(y, z) \right] \stackrel{\text{want}}{=} 2yz + \ln(1+x^2)$$

$$\ln(1+x^2) + \frac{\partial g}{\partial y} =$$

$$\text{So } \frac{\partial g}{\partial y} = 2yz.$$

$$g(y, z) = \int 2yz dy = y^2 z + h(z)$$

$$\text{So } \varphi = y \ln(1+x^2) + y^2 z + h(z)$$

$$(C): \frac{\partial \varphi}{\partial z} = \frac{2}{2z} \left[y \ln(1+x^2) + y^2 z + h(z) \right]$$

$$= 0 + y^2 + h'(z) \stackrel{\text{want}}{=} 1 + y^2$$

$$h'(z) = 1 \quad . \quad \text{So } h(z) = z + C$$

$$\varphi = y \ln(1+x^2) + y^2 z + z + C, \quad \text{arbitrary const.}$$

3. (10 pts) If γ is the positively oriented boundary curve of an ellipse E of area 7 centered at $(2, 3)$, compute

$$\int_{\gamma} -xy \, dx + x^2 \, dy = \iint_E \frac{\partial}{\partial y} \left(\frac{2x^2}{2x} - \frac{\partial(-xy)}{\partial y} \right) dA$$

↑
Green's
 E

$$= 3 \iint_E x \, dA = 3 \bar{x} \cdot \text{Area}(E) = 3 \cdot 2 \cdot 7 = 42$$

\uparrow
x-coord of
center of mass
centroid

4. (10 pts) Let $\vec{F} = x^2yz\hat{i} + xy^2z\hat{j} + xyz^2\hat{k}$. Compute $\text{Curl } \vec{F}$ and $\text{Div } \vec{F}$.


$$\begin{aligned} \text{Curl } \vec{F} &= \det \begin{bmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^2yz & xy^2z & xyz^2 \end{bmatrix} \\ &= \hat{i}(xz^2 - xy^2) - \hat{j}(yz^2 - yx^2) + \hat{k}(y^2z - x^2z) \\ &= x(z^2 - y^2)\hat{i} + y(x^2 - z^2)\hat{j} + z(y^2 - x^2)\hat{k} \end{aligned}$$

$$\text{Div } \vec{F} = 2xyz + 2xyz + 2xyz = 6xyz$$

5. (20 pts) Let $\vec{F} = P_1(y)\hat{i} + P_2(z)\hat{j} + z\hat{k}$ where $P_1(y)$ is a polynomial in y and $P_2(z)$ is a polynomial in z . Let S denote the surface given by $z = 4 - x^2 - y^2$ above the xy -plane. Compute

$$\int_S \vec{F} \cdot \hat{n} dA$$

where \hat{n} denotes the upward pointing unit normal vector to S .



$$\operatorname{Div} \vec{F} = 0 + 0 + 1 = 1$$

$$\iint_S \vec{F} \cdot \hat{n} dA = \iint_{S_{\text{BOTTOM}}} [\underbrace{*, *, 0}_{=0}] \cdot (\hat{k}) dA = 0$$

$$\iint_S \vec{F} \cdot \hat{n} dA = \left(\iint_{S'} + \iint_{S'_1} \right) \vec{F} \cdot \hat{n} dA$$

$$\stackrel{\text{Div}}{\uparrow} \iint_{\Omega} \underbrace{\operatorname{Div} \vec{F}}_1 dV = \text{Volume}(\Omega)$$

$$= \int_0^{2\pi} \int_0^2 4 - r^2 r dr d\theta = 2\pi \int_0^2 4r - r^3 dr$$

$$= 2\pi \left[2r^2 - \frac{1}{4}r^4 \right]_0^2 = 2\pi (8 - 4) = 8\pi$$


6. (20 pts) Let $\vec{F} = xy\hat{i} + xz\hat{j} + z\hat{k}$. Let S denote the surface given by $z = 4 - x^2 - y^2$ above the xy -plane. Write out (but DO NOT COMPUTE) a double integral in the form

$$\int_{\square}^{\square} \int_{\square}^{\square} \underline{\hspace{10em}} dy dx$$

that yields

$$\int_S \vec{F} \cdot \hat{n} dA,$$

where \hat{n} denotes the upward pointing unit normal vector to S . Be sure to put values in all the spots where boxes and blanks appear in the double integral above. Do NOT compute the integral.



$$\vec{r}(x, y) = x\hat{i} + y\hat{j} + (4 - x^2 - y^2)\hat{k}$$

$$\vec{r}_x \times \vec{r}_y = \det \begin{bmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 0 & -2x \\ 0 & 1 & -2y \end{bmatrix} = 2x\hat{i} + 2y\hat{j} + \hat{k}$$

↑
upward

$$\iint_S \vec{F} \cdot \hat{n} dA = \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} [xy\hat{i} + x(4-x^2-y^2)\hat{j} + (4-x^2-y^2)\hat{k}] \cdot [2x, 2y, 1] dy dx$$

$$= \int_{-2}^2 \int_{-\sqrt{4-x^2}}^{\sqrt{4-x^2}} 2x^2y + 2xy(4-x^2-y^2) + (4-x^2-y^2) dy dx$$