



# Lesson 12

## §3.5 - Derivatives of Trig Functions

Note:




$$A = \pi r^2$$



$\theta$  in radians

$$A_{\text{sector}} = \left(\frac{\theta}{2\pi}\right) \pi r^2 = \frac{1}{2} r^2 \theta$$



area  $\triangle AOD < A_{\text{sector}} < \text{area } BOC$

$$\frac{1}{2} (\cos x)(\sin x) < \frac{1}{2} (1)^2 x < \frac{1}{2} (1) \frac{\sin x}{\cos x}$$

$$\cos x < \frac{x}{\sin x} < \frac{1}{\cos x} \quad (\text{if } x > 0)$$

$$\frac{1}{\cos x} > \frac{\sin x}{x} > \cos x \quad \text{As } x \rightarrow 0^+$$

$$\downarrow \quad \therefore \quad \downarrow \quad \downarrow$$

$$1 \quad \quad \quad 1 \quad \quad \quad 1$$

Hence,

Thm: If  $x$  is in radians, then

2

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

$$\lim_{x \rightarrow 0} \frac{x}{\sin x} = 1$$

$$\lim_{x \rightarrow 0} \frac{\cos x - 1}{x} = 0$$

Ex 1 Compute these limits:

$$(a) \lim_{x \rightarrow 0} \frac{\sin 5x}{3x} = \lim_{x \rightarrow 0} \left( \frac{\sin 5x}{5x} \right) \left( \frac{5x}{3x} \right)$$

let  $t = 5x$

$$= \lim_{t \rightarrow 0} \left( \frac{\sin t}{t} \right) \left( \frac{5}{3} \right) = \frac{5}{3} \quad \checkmark$$

$$(b) \lim_{x \rightarrow 0} \frac{\tan 3x}{\sin \pi x} = \lim_{x \rightarrow 0} \frac{\frac{\sin 3x}{\cos 3x}}{\sin \pi x} \quad \boxed{3}$$

$$= \lim_{x \rightarrow 0} \frac{\sin 3x}{(\sin \pi x)(\cos 3x)}$$

$$= \lim_{x \rightarrow 0} \left( \frac{\sin 3x}{3x} \right) \left( \frac{\pi x}{\sin \pi x} \right) \left( \frac{1}{\cos 3x} \right) \left( \frac{3x}{\pi x} \right)$$

let  $t = 3x$                       let  $t = \pi x$

$$= (1) (1) (1) \left( \frac{3}{\pi} \right) = \frac{3}{\pi} \checkmark$$

$$(c) \lim_{x \rightarrow 3} \frac{\sin 2(x-3)}{8(3-x)} = \lim_{t \rightarrow 0} \frac{\sin 2t}{-8t}$$

let  $\boxed{t = x-3}$

$$= \lim_{t \rightarrow 0} \left( \frac{\sin 2t}{2t} \right) \left( \frac{2t}{-8t} \right)$$

$$= (1) \left( -\frac{2}{8} \right) = -\frac{1}{4} \checkmark$$

Thm: If  $x$  is in radians then

4

$$\boxed{\frac{d(\sin x)}{dx} = \cos x}$$

$$\boxed{\frac{d(\cos x)}{dx} = -\sin x}$$

Reason:

$$\frac{d(\sin x)}{dx} = \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{(\sin x \overset{\vee}{\cos h} + \cos x \overset{\vee}{\sin h}) - \sin x}{h}$$

$$= \lim_{h \rightarrow 0} (\sin x) \left( \frac{\overset{\vee}{\cos h} - 1}{h} \right) + (\cos x) \left( \frac{\overset{\vee}{\sin h}}{h} \right)$$

$$= \cos x \checkmark$$

Ex2 Find derivative

$$\textcircled{a} \underline{y = x \sin x + \cos x}$$

$$\Rightarrow y' = x(\cos x) + (\sin x)(1) + (-\sin x)$$

$$y' = x \cos x \checkmark$$

(b)  $y = \tan x = \frac{\sin x}{\cos x}$

$\Rightarrow \frac{dy}{dx} = \frac{(\cos x)(\cos x) - (\sin x)(-\sin x)}{\cos^2 x}$

$= \frac{1}{(\cos x)^2} = \sec^2 x$  Hence

If  $x$  is in radians, then

$(\sin x)' = \cos x$	$(\cos x)' = -\sin x$
$(\tan x)' = \sec^2 x$	$(\cot x)' = -\csc^2 x$
$(\sec x)' = \sec x \tan x$	$(\csc x)' = -\csc x \cot x$