

Def'n. Let $A \subseteq \mathbb{R}$ and $f: A \rightarrow \mathbb{R}$.

We say that f is uniformly

continuous on A if for every

$\epsilon > 0$, there is a $\delta(\epsilon) > 0$

such that if $x_1, x_2 \in A$

are any numbers satisfying

$|x_1 - x_2| < \delta(\epsilon)$, then

$|f(x_1) - f(x_2)| < \epsilon$.

The point is that if we want to guarantee that $|f(x_1) - f(x_2)|$, it suffices to choose δ sufficiently small, say $|x_1 - x_2| < \delta(\epsilon)$.

Thm. If $I = [a, b]$ is a closed bounded interval, and f is continuous on I , then f is uniformly continuous on I .

Pf. If f is not uniformly
 continuous on I , then there
 is a number $\epsilon_0 > 0$, such that
 for any number $\delta > 0$, there
 are numbers $u = u(\delta)$ and
 $v = v(\delta)$ such that
 $|u(\delta) - v(\delta)| < \delta$, but
 that $|f(u(\delta)) - f(v(\delta))| \geq \epsilon_0$

In fact, for every $n \in \mathbb{N}$,

there are numbers u_n and v_n

in I such that $|u_n - v_n| < \frac{1}{n}$

but that $|f(u_n) - f(v_n)| \geq \epsilon_0$.

Since I is bounded, the

Bolzano-Weierstrass Thm

implies that the sequence

(u_n) has a subsequence

$\{u_{n_k}\}$ that converges
to a number x in $[a, b]$.

Since $a \leq u_{n_k} \leq b$ for all
 $k=1, 2, \dots$, it follows that

$x = \lim_{k \rightarrow \infty} u_{n_k}$ also is in $[a, b]$.

Note that

$$|v_{n_k} - x| \leq |v_{n_k} - u_{n_k}| + |u_{n_k} - x|$$

We know $|v_n - u_n| < \frac{1}{n} \rightarrow 0$

In particular, $\lim_{k \rightarrow \infty} |v_{n_k} - u_{n_k}|$

approaches 0. In addition,

we know that $u_{n_k} - x$ also

approaches 0. We conclude

that $\lim_{k \rightarrow \infty} v_{n_k} = x$. ~~Since~~

It is clear that both

u_{n_k} and v_{n_k} approach x .

Since f is continuous at x ,

both $f(u_{n_k})$ and $f(v_{n_k})$

converge to $f(x)$. But

this is impossible since

$$|f(u_n) - f(v_n)| \geq \epsilon_0.$$

Thus, our assumption that

f is not uniformly continuous

implies that f is not

continuous at some point x in I .

Consequently, if f is continuous at every point of I , then f is uniformly continuous on I .

Lipschitz Functions.

Def'n. Let $A \subseteq \mathbb{R}$ and let $f: A \rightarrow \mathbb{R}$.

If there exists a constant $K > 0$

such that $|f(x) - f(u)| \leq K|x - u|$,

(1)

for all $x, u \in A$, then


f is said to be a Lipschitz

function on A .

Geometrically, the Lipschitz

condition can be written as

$$\left| \frac{f(x) - f(u)}{x - u} \right| \leq k$$



Thus, the slopes of all the segments joining two points on the graph of $y = f(x)$ are bounded by a constant K .

Thm. If $f: A \rightarrow \mathbb{R}$ is a Lipschitz function, then f is uniformly continuous

Pf If (1) is true, then

given $\epsilon > 0$, we can take

$$\delta = \frac{\epsilon}{K}. \quad \text{If } x, u \in A$$

satisfy $|x - u| < \delta$, then

$$\begin{aligned} |f(x) - f(u)| &\leq K|x - u| \\ &\leq K \cdot \frac{\epsilon}{K} = \epsilon. \end{aligned}$$

Ex. The function $g(x) = \sqrt{x}$
is continuous on $[0, 1]$,

but it is not Lipschitz,

because if

$$g(x) - g(0) \leq K(x - 0) = Kx,$$


then $\sqrt{x} \leq Kx$ for all $x \in [0, 1]$.

Thus $1 \leq K\sqrt{x}$. But this

cannot happen if x is small in $[0, 1]$

Def'n. Let $I \subseteq \mathbb{R}$ be an interval and let $s: I \rightarrow \mathbb{R}$.

Then s is called a step function if it has only a finite number of values. Moreover, on each interval, the step function takes on only one value in the interior of each interval.



Very closed

Thm. Let $I = [a, b]$ be a closed bounded interval, and let

$f: I \rightarrow \mathbb{R}$ be continuous on I .

If $\varepsilon > 0$, then there exists

a step function $S : I \rightarrow \mathbb{R}$

such that $|f(x) - S(x)| < \varepsilon$

for all $x \in I$.

Pf. The function f is

uniformly continuous, so

given $\varepsilon > 0$, there is a

number $\delta(\varepsilon)$ such that

if $x, y \in I$ and $|x - y| < \delta$,

then $|f(x) - f(y)| < \epsilon$.

Let $I = [a, b]$ and let m

be sufficiently large so

that $h = (b-a)/m < \delta(\epsilon)$

Now we divide $[a, b]$ into

m disjoint intervals of

length h .

$$a = x_0 < x_1 < \dots < x_{m-1} < x_m = b.$$

$$\text{where } x_i - x_{i-1} = h = \frac{b-a}{m}.$$

Now define

$$S_{\xi}(x) = f(a + kh), \text{ for}$$

$$x \in I_k, \quad k=1, \dots, m,$$

so S_{ξ} is constant on each

interval (The value of S_{ξ}

on I_k is the value of f

at the right endpoint of I_k

Hence, if $x \in I_k$, then

$$|f(x) - S_\varepsilon(x)| = |f(x) - f(a+kh)|$$

$$< \varepsilon.$$

Hence $|f(x) - S_\varepsilon(x)| < \varepsilon$

for all $x \in I$.