

3.5 Cauchy's Criterion.

Def'n. A sequence $X = (x_n)$ is

a Cauchy sequence if

for all $\epsilon > 0$, there exists a

number H in \mathbb{N} so that

if $n, m \geq H$, then

$$|x_n - x_m| < \epsilon$$

Even though the definition
does not mention a limit x ,
Still, the numbers x_n and x_m
get closer as $n, m \rightarrow \infty$

Lemma. If a sequence approaches
a limit x , then the sequence
 (x_n) is Cauchy

Proof of Lemma. If $x = \lim (x_n)$,
 then given $\epsilon > 0$, there is a
 natural number K , such that
 if $n \geq K$, then $|x_n - x| < \frac{\epsilon}{2}$.

Thus, if

$n, m \geq K$, then we have

$$|x_n - x_m| = |(x_n - x) + (x - x_m)|$$

$$\leq |x_n - x| + |x - x_m| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

Since $\varepsilon > 0$ is arbitrary,
it follows that (x_n) is a
Cauchy sequence.

Lemma. A Cauchy sequence
is bounded.

Pf. Let $X = (x_n)$ be Cauchy,
and set $\epsilon = 1$. There is H in \mathbb{N} so
that, if $n \geq H$, then

$$|x_n - x_H| < 1. \quad \text{By the}$$

Triangle Inequality, we have

$$|x_n| \leq |x_H + (x_n - x_H)|$$

$$\leq |x_H| + 1$$

If we set

$$M = \max \left\{ |x_1|, |x_2|, \dots, |x_{H-1}|, |x_H| + 1 \right\},$$

then it follows that

$$|x_n| \leq M, \quad \text{for all } n.$$

Cauchy Convergence Thm.

A sequence $X = (x_n)$ is convergent if it is a Cauchy sequence. We have to find x !!

We already showed that if X is convergent, then it is Cauchy. To prove the other direction, suppose X is Cauchy. We showed above that X is therefore bounded. By the Bolzano-Weierstrass theorem, there exists a subsequence

$X' = (x_{n_k})$ of X that
converges to a number x^* .

We will show that $\lim x_n = x^*$.

Since $X = (x_n)$ is a Cauchy
sequence, given $\varepsilon > 0$, there
is a natural number H

such that if $n, m \geq H$

then $|x_n - x_m| < \frac{\varepsilon}{2}$. (1)

Since the subsequence

$X' = \{x_{n_k}\}$ converges to x^* ,

there is a natural number

$K \geq H$ that belongs to the set $\{n_1, n_2, \dots\}$ such that

$$|x_K - x^*| < \frac{\epsilon}{2}$$

Since $K \geq H$, it follows

from (1) with $m = K$ that

$$|x_n - x_k| < \frac{\epsilon}{2} \quad \text{for } n \geq H.$$

Therefore, if $n \geq H$,

we have

$$\begin{aligned} |x_n - x^*| &= |(x_n - x_k) + (x_k - x^*)| \\ &\leq |x_n - x_k| + |x_k - x^*| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2}. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, we obtain that $\lim(x_n) = x^*$.

Ex. The polynomial equation

$$x^3 - 5x + 1 = 0 \text{ has a root}$$

π with $0 < \pi < 1$.

We define an iteration procedure to construct a sequence (x_n) that approaches the root π .

We define x_1 to be any number with $0 < x_1 < 1$.

and we define

$$x_n^3 - 5x_{n+1} + 1 = 0,$$

$$\text{or } x_{n+1} = \frac{1}{5}(x_n^3 + 1)$$

It is easy to verify that if

$$0 \leq x_n \leq 1, \text{ then } 0 \leq x_{n+1} \leq 1.$$

Hence, $0 \leq x_n \leq 1$ for all $n \geq N$.

We can estimate $|x_{n+2} - x_{n+1}|$ by

$$\begin{aligned} |x_{n+2} - x_{n+1}| &= \\ &= \left| \frac{1}{5}(x_{n+1}^3 + 1) - \frac{1}{5}(x_n^3 + 1) \right| \\ &= \frac{1}{5} |x_{n+1}^3 - x_n^3| \end{aligned}$$

$$= \frac{1}{5} \left| x_{n+1}^2 + x_{n+1}x_n + x_n^2 \right| |x_{n+1} - x_n|$$

$$\leq \frac{3}{5} |x_{n+1} - x_n|.$$

Definition. We say that a sequence (x_n) of real numbers is contractive if there is a constant C , $0 < C < 1$, such that

$$|x_{n+2} - x_{n+1}| \leq C |x_{n+1} - x_n|$$

for all $n \in \mathbb{N}$. C is the constant of the sequence.

We now prove

Thm. Every contractive
sequence is a Cauchy
Sequence and therefore
is convergent.

Observe that in the above
example, (x_n) is contractive
with $C = \frac{3}{5}$.

Pf. Using the contractive inequality, we get:

$$\begin{aligned} |x_{n+2} - x_{n+1}| &\leq C |x_{n+1} - x_n| \\ &\leq C^2 |x_n - x_{n-1}| \\ &\vdots \\ &\leq C^n |x_2 - x_1| \end{aligned}$$

To show that (x_n) is Cauchy, we have

Hence

14

$$|x_m - x_n| \quad (2)$$

$$\leq |x_m - x_{m-1}| + |x_{m-1} - x_{m-2}| \\ + \dots + |x_{n+1} - x_n|$$

$$\leq (c^{m-2} + \dots + c^{n-1}) |x_2 - x_1|$$

$$= c^{n-1} \left(\frac{1 - c^{m-n}}{1 - c} \right) |x_2 - x_1|$$

$$\leq c^{n-1} \left(\frac{1}{1 - c} \right) |x_2 - x_1|$$

We conclude that (x_n)

is a Cauchy sequence

and therefore convergent.

Observe that we can estimate the accuracy of (x_n) :

$$|x_m - x_n| \leq \frac{c^{n-1}}{1-c} |x_2 - x_1|$$

Since $\lim (x_m) = \tilde{x}$, we have

$$|\tilde{x} - x_n| \leq \frac{c^{n-1}}{1-c} |x_2 - x_1|$$

which shows that the error approaches 0 exponentially.

$$\text{Since } x_{n+1} = \frac{1}{5} (x_n^3 + 1),$$

we can take the limit as

$n \rightarrow \infty$, so

$$x = \frac{1}{5} (x^3 + 1)$$

$$\rightarrow 5x = x^3 + 1, \quad \text{or}$$

$$x^3 - 5x + 1 = 0$$