

Infinite Limits 4.3

Def'n. Let $A \subseteq \mathbb{R}$, let $f: A \rightarrow \mathbb{R}$

and let c be a cluster point of A .


We say f tends to ∞ as $x \rightarrow c$

and write $\lim_{x \rightarrow c} f = \infty$

if for all $\alpha \in \mathbb{R}$, there is $\delta > 0$

so that if $x \in A$ and $0 < |x - c| < \delta$

then $f(x) > \alpha$



Ex. Show $\lim_{x \rightarrow 0} \frac{1}{\sqrt{x}} = \infty$. $x > 0$.

We need $\frac{1}{\sqrt{x}} > \alpha$

$$\rightarrow \frac{1}{x} > \alpha^2 \rightarrow x < \frac{1}{\alpha^2}$$

\therefore Set $\delta = \frac{1}{\alpha^2}$.

$$\text{If } 0 < x < \frac{1}{\alpha^2} \rightarrow \sqrt{x} < \frac{1}{\alpha}$$

$$\rightarrow \frac{1}{\sqrt{x}} > \alpha \quad \checkmark$$

Limits at ∞ .

Def'n. Let $A \subseteq \mathbb{R}$, and let $f: A \rightarrow \mathbb{R}$.

Suppose that $(a, \infty) \subseteq A$ for

some $a \in \mathbb{R}$. We say L is a

limit of f as $x \rightarrow \infty$, and we write

$\lim_{x \rightarrow \infty} f(x) = L$ if given any $\epsilon > 0$

there is $K > a$ so that

if $x > K$, then $|f(x) - L| < \epsilon$

Ex. Show that $\lim_{x \rightarrow \infty} \frac{x^2 - 3x - 1}{2x^2 + 1} = \frac{1}{2}$

It's easy to show that $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$.

Note $\frac{x^2 - 3x - 1}{2x^2 + 1} = \frac{x^2 \left(1 - \frac{3}{x} - \frac{1}{x^2}\right)}{x^2 \left(2 + \frac{1}{x^2}\right)}$

$= \frac{1 - \frac{3}{x} - \frac{1}{x^2}}{2 + \frac{1}{x^2}}$

Using the
analogs of
the limit rules

$\lim_{x \rightarrow \infty} \frac{1}{x} = 0, \quad \frac{1}{x^2} = 0, \text{ etc}$

we obtain $\lim_{x \rightarrow \infty} \frac{1 - \frac{3}{x} - \frac{1}{x^2}}{2 + \frac{1}{x^2}} = \frac{1 - 0 - 0}{2 + 0}$

$$= \frac{1}{2}$$

5.2 Continuous Functions

Def'n. Let $A \subseteq \mathbb{R}$, let $f: A \rightarrow \mathbb{R}$

and let $c \in A$. We say f is

continuous at c if for any

$\epsilon > 0$, there exists $\delta > 0$ such

that if x satisfies $x \in A$ with

$|x - c| < \delta$, then $|f(x) - f(c)| < \epsilon$.

If f is continuous at c , then

three conditions must hold:

(i) f must be defined at c ,

(ii) The limit of f at c must exist,

(iii) These two values must be equal.

Of course we have the following result.

Sequential Criterion for Continuity.

A function $f: A \rightarrow \mathbb{R}$ is continuous at c if and only if for every sequence (x_n) in A that converges

to c , $f(x_n) \rightarrow f(c)$.

converges to c , the sequence $(f(x_n))$ converges to $f(c)$.

And we have a

Discontinuity Thm. Let $A \subseteq \mathbb{R}$,

let $f: A \rightarrow \mathbb{R}$, and let $c \in A$.

Then f is discontinuous at c

if and only if there exists

a sequence (x_n) in A such that

(x_n) converges to c , but the

sequence $(f(x_n))$ does NOT converge

to $f(c)$.

Pf. If f is discontinuous at c . 8


there is a number $\epsilon_0 > 0$

such that for every $n \in \mathbb{N}$,

there is a number $x_n \in A$

with $|x_n - c| < \frac{1}{n}$ and

$$|f(x_n) - f(c)| \geq \epsilon_0$$



Choose $\epsilon_0 = \frac{d}{2}$

If $\frac{1}{n}$ is sufficiently small,

$$|f(x_n) - f(c)| \geq \epsilon_0$$

Def'n. Let $A \subseteq \mathbb{R}$, let $f: A \rightarrow \mathbb{R}$. 8.1

If B is a subset of A , we say

that f is continuous on the set B

if f is continuous at every

point of B .

Ex. Let $A = \mathbb{R}$, and define

the Dirichlet function f by

$$f(x) = \begin{cases} 1 & \text{if } x \text{ is rational} \\ 0 & \text{if } x \text{ is irrational} \end{cases}$$

We show that f is discontinuous at every point of \mathbb{R} . First,

we suppose that c is rational,

so that $f(c) = 1$. Let (x_n)

be a sequence

of irrational

numbers that

converge to c

.....

~~|||||~~
c

$$\text{Set } x_n = c + \frac{\sqrt{2}}{n}$$

Then $f(x_n) = 0$ for all $n \in \mathbb{N}$

Since $f(c)=1$, it follows that

$f(x_n)$ does not converge to $f(c)$.

By the Discontinuity Criterion

f is not continuous at c .

Similarly, suppose c is an

irrational number. Since the

rational numbers are dense in \mathbb{R} , for

every n we can find a rational

number $x_n \in (c, c + \frac{1}{n})$,

so that $\lim_{n \rightarrow \infty} f(x_n) = 1$.

Since (x_n) converges to c ,

and $f(c) = 0$, it follows that

$\lim_{n \rightarrow \infty} (f(x_n))$ does not converge

to $f(c)$. By the Discontinuity

Criterion, it follows that


f is discontinuous at c .

$\therefore f$ is discontinuous at each
point of \mathbb{R} .

Ex. Thomae Fcn. We define

$h: \mathbb{R} \rightarrow \mathbb{R}$ by

$$h(x) = \begin{cases} \frac{1}{q} & \text{if } x = \frac{p}{q} \text{ and } p, q \text{ have} \\ & \text{no common factor } > 1 \\ 0 & \text{if } x \text{ is irrational} \end{cases}$$



We show that the function h is continuous at each irrational number x' and discontinuous at each rational number x'' .

It's easy to show that h is discontinuous at each rational.

In fact, as above, if c is rational, then $h(c) = 1/q$ for some

positive q . But let

As above, let x'

(x_n) be a sequence of
irrational numbers that
converges to c . Then

$f(x_n)$ does NOT converge to $f(c)$.

Hence the Discontinuity Thm

implies that h is discontinuous
at c .

Now we show h is continuous
at each irrational number b .

Let ε be any positive ^{number} $\sqrt{\quad}$. Then

there is a number n_0 with

$\frac{1}{n_0} < \varepsilon$. There are only

a finite number of rationals

with denominator less than n_0

in the interval $(b-1, b+1)$.

Hence we can choose $\delta > 0$

so small that the neighborhood

$(b-\delta, b+\delta)$ contains no

rational numbers with denominator

less than n_0 . It follows that

for $|x - b| < \delta$, $x \in A$ we have

$$|h(x) - h(b)| = |h(x)| \leq \frac{1}{n_0} < \varepsilon.$$

Thus h is continuous at

the irrational number b