

## 5.6 Inverse Functions 1


If  $f$  is continuous on a

closed bounded interval

$I = [a, b]$ , we showed that

there are 2 points  $x_*$  and  $x^*$

such that  $f(x_*) \leq f(x) \leq f(x^*)$ .



We will say that a function  $f$  is strictly increasing on an interval  $I$  if whenever  $x' < x''$  then  $f(x') < f(x'')$ .

Let's assume that  $f$  is strictly increasing and continuous on  $[x_*, x^*]$ .

Suppose that  $k$  satisfies


$$f(x') < k < f(x'')$$

Then the Intermediate Value

Thm (IVT) says that

there is a number  $x_0 \in (x_*, x^*)$

such that  $f(x_0) = k$ .



In fact, this  $x_0$  must be

unique, for if  $x'_0$  is another

number with  $f(x_0) = k = f(x'_0)$ ,

then  $f$  would not be strictly

increasing. Hence  $x_0$  is unique.

We can define the inverse

by setting  $g(y) = x$ ,

whenever  $f(x) = y$ .

Thus the function  $g(y)$  is

well-defined for all  $y$

that satisfy

$$f(x_*) \leq y \leq f(x^*)$$

Note that if  $x \in [x_*, x^*]$ ,


then  $f(x) \in [f(x_*), f(x^*)]$

$= J$ . If we set  $y = f(x)$ ,

then  $y \in \text{Range of } f$ .

Thus  $g(y) = x$ , so

$$J = f(I)$$



We want to show that  
the inverse function  $g$   
is continuous on  $J$ .


Let  $c \in I$ . For any small  
number  $\epsilon$ , set  $y_+ = f(c + \epsilon)$   
and set  $y_- = f(c - \epsilon)$ .

This implies

$$g(y_+) = c + \epsilon \text{ and } g(y_-) = c - \epsilon$$

If  $c - \epsilon < x < c + \epsilon$ , then

$$y_- < f(x) < y_+$$



Now set

$$\delta = \min \{ |y_+ - f(c)|, |y_- - f(c)| \}$$

It follows that if

$y \in V_\delta(f(c))$ , then

$g(y) \in V_\varepsilon(c)$ .

It follows that  $g$  is continuous at  $f(c)$ . Since  $c$  is arbitrary,

it follows that

$g: [f(a), f(b)] \rightarrow [a, b]$  is

continuous at  $c$ .



Thus we've proved that if  $f$  is continuous on an interval  $I$ , and if  $f$  is strictly increasing on  $I$ , then there is a continuous function  $g$  on  $J = [f(a), f(b)]$  such that

$$g(f(x)) = x, \quad \text{all } x \in [a, b].$$

## 6.1 The derivative.

Def'n. Let  $I \subseteq \mathbb{R}$  be an

interval, let  $f: I \rightarrow \mathbb{R}$ ,

and let  $c \in I$ . We say that

$L$  is the derivative of  $f$  at  $c$

if given any  $\epsilon > 0$ , there is

$\delta(\epsilon) > 0$  so that if  $x \in I$  and

satisfies

$0 < |x - c| < \delta(\epsilon)$ , then

$$\left| \frac{f(x) - f(c)}{x - c} - L \right| < \epsilon.$$

We write  $f'(c) = L$

Thus the derivative of  $f$   
at  $c$  is given by

$$f'(c) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$$

A useful theorem:

Thm. If  $f: I \rightarrow \mathbb{R}$  has a derivative at  $c \in I$ , then  $f$  is continuous at  $c$ .

Pf. For all  $x \in I$ ,  $x \neq c$ , we have

$$f(x) - f(c) = \left( \frac{f(x) - f(c)}{x - c} \right) (x - c).$$

Since  $\lim \left( \frac{f(x) - f(c)}{x - c} \right)$  and

$\lim (x - c)$  exist, the

product rule implies that

$$\lim_{x \rightarrow c} (f(x) - f(c))$$

$$= \lim_{x \rightarrow c} \left( \frac{f(x) - f(c)}{x - c} \right) \cdot \lim_{x \rightarrow c} (x - c)$$

$$= f'(c) \cdot 0 = 0$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c).$$

This shows that  $f$  is  
continuous at  $c$

These limit laws are very important:

Thm. Suppose that both  $f$  and  $g$  are differentiable at  $c \in I$ . Then:

$$(a) \quad (bf)'(c) = bf'(c)$$

$$(b) \quad (f+g)'(c) = f'(c) + g'(c).$$

(c) Product Rule.

$$(fg)'(c) = f'(c)g(c) + f(c)g'(c)$$

(d) Quotient Rule If  $g(c) \neq 0$ ,

$$\text{then } \left(\frac{f}{g}\right)'_c = \frac{f'(c)g(c) - f(c)g'(c)}{(g(c))^2}.$$

We'll prove the Product

and Quotient Rules:

(c) (Prod. Rule) Let  $p = fg$ .

$$\frac{p(x) - p(c)}{x - c} = \frac{f(x)g(x) - f(c)g(c)}{x - c}$$

$$= \frac{f(x)g(x) - f(c)g(x) + f(c)g(x) - f(c)g(c)}{x-c}$$

$$= \frac{f(x) - f(c)}{x-c} g(x) + f(c) \cdot \frac{g(x) - g(c)}{x-c}$$

Since  $g(x)$  is differentiable at  $c$ , it's also continuous at  $c$ .

$$\lim_{x \rightarrow c} \frac{p(x) - p(c)}{x - c} = f'(c)g(c) + f(c)g'(c)$$

The Quotient Rule is:

$$\left( \text{Set } q(x) = \frac{f(x)}{g(x)} \right)$$



Since  $g$  is differentiable, it's also continuous at  $c$ . Hence

$g(x) \neq 0$  in a neighborhood of  $c$

(since  $g(c) \neq 0$ )

$$\frac{g(x) - g(c)}{x - c} = \frac{f(x)/g(x) - f(c)/g(c)}{x - c}$$

$$= \frac{f(x)g(c) - f(c)g(x)}{g(x)g(c)(x - c)}$$

$$= \frac{f(x)g(c) - f(c)g(c) + f(c)g(x) - f(c)g(x)}{g(x)g(c)(x-c)}$$

$$= \frac{1}{g(x)g(c)} \left[ \frac{f(x)-f(c)}{x-c} g(c) - f(c) \frac{g(x)-g(c)}{x-c} \right]$$

Using the continuity of  $g$ , we get


$$g'(c) = \lim_{x \rightarrow c} \frac{g(x) - g(c)}{x - c}$$

$$= \frac{f'(c)g(c) - f(c)g'(c)}{(g(c))^2}$$

If we use the notation

$$\frac{d}{dx} (y(x)) = \lim_{x \rightarrow c} \frac{y(x) - y(c)}{x - c}$$

we get  $= \lim_{x \rightarrow c} \frac{\Delta y}{\Delta x}$



As  $x \rightarrow c$ ,  $\Delta x = x - c \rightarrow 0$

$\therefore \lim \frac{\Delta y}{\Delta x} =$  slope of the  
tangent line  
at  $(c, y(c))$