

## 3.1 Sequences

A sequence  $X$  is a function from  $\mathbb{N}$  to  $\mathbb{R}$ . Sometimes  $X$  is defined by a formula for the  $n$ -th term  $x_n$  such as

$$x_n = \frac{2^n}{n+1} \cdot \text{Sometimes we just}$$

define the first few terms,

$$X = \left\{ \frac{1}{3}, \frac{1}{5}, \frac{1}{7}, \dots \right\} \text{ or}$$

$$x_n = \frac{1}{2n+1}$$

We can also give a recursive formula for  $x_n$ :

$$x_n = \frac{x_{n-1}}{x_{n-1}^2 + 1}, \quad x_1 = 3.$$

It is very important to compute the limit of a sequence.

**Definition.** We say a sequence

$X$  converges to  $x$  if for all  $\varepsilon > 0$ ,

there is a number  $K$  in  $\mathbb{N}$ , so that

if  $n \geq K$ , then  $|x_n - x| < \varepsilon$ .

The number  $x$  is the limit of  $X$ , and we say  $X$  is convergent.

If  $X$  is not convergent, we say  $X$  is divergent.

A sequence can only have at most one limit. Suppose  $\lim X = x'$

and  $\lim X = x''$ . Set  $\epsilon = \frac{|x' - x''|}{2}$ .

Choose  $K_1$  so  $|x_n - x'| < \epsilon$

if  $n \geq K_1$ .

and choose  $K_2$  so that

$$|x_n - x''| < \epsilon \quad \text{if } n \geq K_2.$$

Now set  $K = \text{maximum of } \{K_1, K_2\}$ .

Then if  $n \geq K$ ,

$$|x' - x''| = |(x' - x_n) - (x'' - x_n)|$$

$$\leq |x' - x_n| + |x'' - x_n|$$

$$< \epsilon + \epsilon = 2\epsilon$$

$$= |x' - x''|.$$

Dividing by  $|x' - x''|$  we get  $1 < 1$ .

The contraction implies that

$$x' = x''.$$

Some examples:

Compute  $\lim \frac{1}{n}$ .

We proved that for any  $\varepsilon > 0$ ,

there is a  $K$  so that if  $n \geq K$ ,

$\frac{1}{n} < \varepsilon$ . We obtain that

$|\frac{1}{n} - 0| = \frac{1}{n} < \varepsilon$ . It follows

that  $\lim(\frac{1}{n}) = 0$ .

Ex. Prove that  $\lim \left( \frac{3}{n+5} \right) = 0$ .

Note that  $\frac{3}{n+5} < \frac{3}{n}$ .

For a given  $\epsilon > 0$ , choose  $K > 0$  so that if  $n \geq K$ , then  $\frac{1}{n} < \frac{\epsilon}{3}$ .

If  $n \geq K$ , then

$$\left| \frac{3}{n+5} - 0 \right| = \frac{3}{n+5} < \frac{3}{n} < 3 \cdot \frac{\epsilon}{3} = \epsilon.$$

Hence  $\lim \left( \frac{3}{n+5} \right) = 0$ .



Ex. Show that  $\lim (-1)^n$  does not exist.

Assuming  $\lim (-1)^n = x$ ,

set  $\epsilon = 1$ . Then there

is a  $K \in \mathbb{N}$  so that if  $n \geq K$ ,

then  $|(-1)^n - x| < 1$ .

If  $n$  is even and  $\geq K$ , then

$$|x - 1| < 1 \rightarrow x - 1 > -1 \rightarrow x > 0$$

8.

If  $n$  is odd and  $\geq K$ , then

$$|x+1| = |x - (-1)^n| < 1.$$

Hence,  $x+1 < 1$ , which implies that  $x < 0$ .

This contradiction implies that  $\lim \{-1\}^n$  does not exist.



We now prove

Let  $(x_n)$  be a sequence of numbers and let  $x \in \mathbb{R}$ .

If  $(a_n)$  is a sequence of positive numbers with  $\lim(a_n) = 0$  and if for some constant  $C > 0$  and some  $m \in \mathbb{N}$ , we have

$$|x_n - x| \leq C a_n \quad \text{for all } n \geq m,$$

then it follows that  $\lim x_n = x$ .

Proof. If  $\epsilon > 0$  is given, then

since  $\lim (a_n) = 0$ , we know

there exists  $K$  such that

$n \geq K$  implies  $a_n = |a_n - 0| < \epsilon/c$ .

It follows that if both  $n \geq K$

and  $n \geq m$ , then

$$|x_n - x| \leq C a_n < C(\epsilon/c) = \epsilon.$$

Since  $\epsilon$  is arbitrary, we conclude

that  $x = \lim (x_n)$ .

We will use this to show that

if  $0 < b < 1$ , then  $\lim (b^n) = 0$ .

But first we prove:

Ex. If  $a > 0$ , show  $\lim \left( \frac{1}{1+na} \right) = 0$

Since  $a > 0$ , then

$$0 < na < 1+na, \quad \text{and}$$

$$\text{therefore } 0 < \frac{1}{1+na} < \frac{1}{na}.$$

Thus we have

$$\left| \frac{1}{1+na} - 0 \right| \leq \frac{1}{a} \cdot \frac{1}{n} \quad \text{for all } n \in \mathbb{N}.$$

Since  $\lim \left( \frac{1}{n} \right) = 0$ ,

the above theorem with  $C = \frac{1}{a}$

and  $m=1$  implies that

$$\lim \left( \frac{1}{1+na} \right) = 0.$$

Recall that Bernoulli's  
Inequality states that

if  $x > -1$ , then

$$(1+x)^n \geq 1+nx, \quad \text{all } n \in \mathbb{N}.$$

---

We now show that if

$$0 < b < 1, \quad \text{then } \lim(b^n) = 0.$$

Since  $0 < b < 1$ , we can write

$$b = \frac{1}{1+a}$$



where  $a = \left(\frac{1}{b}\right)^{-1}$ , so that

$a > 0$ . By Bernoulli's Inequality,

we have

$$(1+a)^n \geq 1+na, \text{ where } a > -1.$$

Hence,

$$0 < b^n = \frac{1}{(1+a)^n} \leq \frac{1}{1+na}$$

From the above theorem, we

conclude that  $\lim (b^n) = 0$ .



### 3.2. Limit Theorems.

Using the results of this section, we can analyse the convergence of many sequences.

**Definition.** A sequence  $X = (x_n)$

is bounded if there exists

a number  $M > 0$  such that

$$|x_n| \leq M, \quad \text{for all } n \in \mathbb{N}.$$

Thm. A convergent sequence of real numbers is bounded.

Pf. Suppose that  $\lim x_n = x$

and let  $\epsilon = 1$ . Then there is

a  $K \in \mathbb{N}$  such that  $|x_n - x| < 1$

for all  $n \geq K$ . The Triangle

Inequality with  $n \geq K$  implies

that

$$|x_n| = |x_n - x + x| \leq |x_n - x| + |x|$$

$$< 1 + |x|.$$

If we set

$$M = \max \left\{ |x_1|, |x_2|, \dots, |x_{K-1}|, 1 + |x| \right\}$$

then it follows that

$$|x_n| \leq M, \quad \text{for all } n \in \mathbb{N}.$$

We want to learn how

taking limits interacts

with the operations of

addition, subtraction,

multiplication and division.

Given two sequences  $X = (x_n)$   
and  $Y = (y_n)$ , we define

$$X + Y = (x_n + y_n)$$

$$X - Y = (x_n - y_n)$$

$$XY = (x_n y_n)$$

$$cX = (cx_n)$$

and

$$X/Y = \left( \frac{x_n}{y_n} \right) \quad \left( \begin{array}{l} \text{providing} \\ y_n \neq 0 \end{array} \right)$$

Suppose  $X = (x_n)$  and  $Y = (y_n)$   
converge to  $x$  and  $y$   
respectively. Let  $\epsilon > 0$ .

Addition.

Choose  $K_1$  and  $K_2$  so that

$$|x_n - x| < \frac{\epsilon}{2} \quad \text{if } n \geq K_1 \quad \text{and}$$

$$|y_n - y| < \frac{\epsilon}{2} \quad \text{if } n \geq K_2.$$

Now set  $K = \text{Max} \{ K_1, K_2 \}$

If  $n \geq K$ , then  $n \geq K_1$  and  $n \geq K_2$ . Hence,

$$\begin{aligned} & |(x_n + y_n) - (x + y)| \\ &= |(x_n - x) + (y_n - y)| \\ &\leq |x_n - x| + |y_n - y| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

Hence  $\lim (x_n + y_n) = x + y$ .



For subtraction, we use the same argument. Just replace

$x_n + y_n$  by  $x_n - y_n$  and

$x + y$  by  $x - y$ .

Multiplication. This is a bit

more complicated. Note that

$$|x_n y_n - xy| = \left| \begin{array}{l} (x_n y_n - x_n y) \\ + (x_n y - xy) \end{array} \right|$$

$$\leq |x_n(y_n - y)| + |(x_n - x)y|$$

$$\leq |x_n||y_n - y| + |x_n - x||y|$$

By the boundedness theorem,

there is  $M_1 > 0$  such that

$$|x_n| \leq M_1, \quad \text{all } n.$$

Now set  $M = \text{Max}\{M_1, |y|\}$ .

We conclude that

$$|x_n y_n - x y| \leq M |y_n - y| + M |x_n - x|$$

Now let  $\epsilon > 0$  be given.

Then there exists  $K_1$

such that

$$|x_n - x| < \frac{\epsilon}{2M} \quad \text{if } n \geq K_1.$$

Similarly, there exists  $K_2$

such that

$$|y_n - y| < \frac{\epsilon}{2M} \quad \text{if } n \geq K_2.$$

Now set  $K = \text{Max} \{ K_1, K_2 \}$

If  $n \geq K$ , then

$$|x_n y_n - xy|$$

$$\leq M|y_n - y| + M|x_n - x|$$

$$< M \cdot \frac{\epsilon}{2M} + M \cdot \frac{\epsilon}{2M} = \epsilon.$$

This proves

$$\lim (x_n y_n) = xy.$$