

## 3.6 Properly Divergent Series

Let  $(x_n)$  be a sequence.

- (i) We say  $(x_n)$  tends to  $+\infty$  and write  $\lim(x_n) = +\infty$  if for every  $\alpha \in \mathbb{R}$ , there exists a nat. number  $K(\alpha)$  such that if  $n \geq K(\alpha)$ , then  $x_n > \alpha$ .

(iii) We say  $\{x_n\}$  tends to  $-\infty$

and write  $\lim (x_n) = -\infty$

if for every  $B \in \mathbb{R}$ ,

there exists

a nat. number  $K(B)$  such

that if  $n \geq K(B)$ , then

$x_n < B$ .

In either case, we say  $\{x_n\}$

is properly divergent.

Ex.  $\lim\{n\} = +\infty$ ,

because if  $\alpha$  is given,

let  $K(\alpha)$  be any natural

number & such that  $K(\alpha) > \alpha$ .

If  $n \geq K(\alpha)$ , then  $n > \alpha$ .



Ex.  $\lim(n^2) = +\infty$  Because

if  $K(\alpha) > \alpha$ , and if  $n \geq K(\alpha)$

then  $n^2 \geq n > \alpha$ .

Ex. If  $c > 1$ , then  $\lim c^n = +\infty$

In fact, let  $c = 1+b$ . If

$\alpha$  is given, let  $K(\alpha)$  be a natural number such that

$K(\alpha) > \frac{\alpha}{b}$ . If  $n \geq K(\alpha)$ ,

it follows from Bernoulli's

Inequality that

$$c^n = (1+b)^n \geq 1+nb > 1+\alpha > \alpha.$$



Note that the inequalities

$n \geq K(\alpha) > \frac{\alpha}{b}$  imply that

$$n > \frac{\alpha}{b} \iff nb > \alpha.$$

---

Recall that the Monotone

Convergence Thm states

that a monotone sequence

is convergent if and only if

it's bounded.

Similarly, we have:

Thm. A monotone sequence

is properly divergent if and

only if it is unbounded.

(a) If  $(x_n)$  is an unbounded

sequence, then  $\lim(x_n) = +\infty$   
 increasing

(b) If  $(x_n)$  is an unbounded

decreasing sequence, then

$$\lim(x_n) = -\infty.$$

## Comparison Test :

Thm. Let  $(x_n)$  and  $(y_n)$

be two sequences and

suppose that  $x_n \leq y_n$ , all  $n \in \mathbb{N}$

(a) If  $\lim(x_n) = +\infty$ ,

then  $\lim(y_n) = +\infty$

(b) If  $\lim(y_n) = -\infty$ , then

$\lim(x_n) = -\infty$

Ex.  $\lim (\sqrt{n}) = +\infty$ .

Let  $K(\alpha)$  be any natural number.

with  $K(\alpha) > \alpha^2$ . If

$n \geq K(\alpha)$ , then  $n > \alpha^2$ .

which implies  $\sqrt{n} > \alpha$ .

Compute  $\lim (\sqrt{n+2})$

Note that if we use the same  $K(\alpha)$  as above,

Then, if  $n > \alpha^2$ , then

$$\sqrt{n+2} > \sqrt{n} > \alpha.$$

which implies  $\lim(\sqrt{n+2}) = +\infty$ .

OR, we could have used the  
above convergence test,

With  $x_n = \sqrt{n}$  and  $y_n = \sqrt{n+2}$ .

Since  $\lim(\sqrt{n}) = +\infty$ , we get

$$\lim(\sqrt{n+2}) = +\infty.$$

Compute  $\lim \left\{ \frac{\sqrt{n^2+1}}{\sqrt{n}} \right\}$ .

$$\frac{\sqrt{n^2+1}}{\sqrt{n}} \rightarrow \frac{\sqrt{n^2}}{\sqrt{n}} = \sqrt{n}.$$

$\downarrow$

Set  $y_n$

Set  $x_n$

$\therefore$  Comp Test  $\Rightarrow \lim \left\{ \frac{\sqrt{n^2+1}}{\sqrt{n}} \right\} = +\infty$ .



Ex. What about  $\frac{\sqrt{n}}{(n^2+1)}$

Note that  $\frac{\sqrt{n}}{(n^2+1)} < \frac{\sqrt{n}}{n^2} < \frac{n}{n^2} = \frac{1}{n}$ .

Since  $\lim \frac{1}{n} = 0$ ,

so does  $\lim \frac{\sqrt{n}}{(n^2+1)} = 0$

Limit Comparison Test.

Suppose  $\{x_n\}$  and  $\{y_n\}$

are positive, and that

$$\lim \left\{ \frac{x_n}{y_n} \right\} = L \neq 0.$$

Then  $\lim x_n = +\infty$

if and only if  $\lim y_n = +\infty$

$$\text{Set } x_n = \frac{\sqrt{2n^2+1}}{\sqrt{3n-1}}$$

$$\text{and } y_n = \frac{n}{\sqrt{n}} = \sqrt{n}.$$

Check  $\lim \frac{x_n}{y_n}$ .

$$\frac{x_n}{y_n} = \frac{\sqrt{2n^2+1}}{\sqrt{3n-1}} \cdot \frac{1}{\sqrt{n}}$$

$$= \frac{\sqrt{2n^2+1}}{\sqrt{n} \cdot \sqrt{3n-1}}$$

$$= \frac{n \sqrt{2 + \frac{1}{n^2}}}{\sqrt{n} \cdot \sqrt{n} \cdot \sqrt{3 - \frac{1}{n}}}$$

$$= \sqrt{\frac{2 + \frac{1}{n^2}}{3 - \frac{1}{n}}} \rightarrow \sqrt{\frac{2}{3}}$$

# Proof of Limit Comparison Test.

We have  $\lim \frac{x_n}{y_n} = L > 0$ .

Set  $\epsilon = \frac{L}{2}$ .

$$\rightarrow L - \frac{L}{2} < \frac{x_n}{y_n} < L + \frac{L}{2}$$

if  $n$  is  $> K_1$ .

$$\rightarrow \frac{L}{2} < \frac{x_n}{y_n} < \frac{3L}{2}$$

$$\text{or } \frac{L}{2} y_n < x_n < \frac{3L}{2} y_n$$

Hence the usual Comparison

Test implies :

If  $\lim y_n = +\infty$ , then

$$\lim x_n = +\infty.$$

and if

$\lim x_n = +\infty$ , then

$$\lim y_n = +\infty.$$

$$\text{Compute } \lim (3^n)^{\frac{1}{2n}}$$

{ We use  $\lim n^{\frac{1}{n}} = 1$  at end  
at 3.1. }

$$= \lim 3^{\frac{1}{2n}} \cdot n^{\frac{1}{2n}}$$

$$= \lim (3^{\frac{1}{n}})^{\frac{1}{2}} \cdot (n^{\frac{1}{n}})^{\frac{1}{2}}$$

Note that  $1 \leq 3 \leq n$

$$\therefore 1^{\frac{1}{n}} \leq 3^{\frac{1}{n}} \leq n^{\frac{1}{2n}}$$

↓

conv. to 1

$$\therefore \{3^{\frac{1}{n}}\}^{\frac{1}{2}} \rightarrow \sqrt{1}$$

Also,  $n^{\frac{1}{n}} \rightarrow 1$ ,

$$\text{so } n^{\frac{1}{n}} \{n^{\frac{1}{n}}\}^{\frac{1}{2}} \rightarrow \sqrt{1}.$$

Compute  $\lim \left(1 + \frac{1}{2n}\right)^{3n}$

$$= \lim \left\{ \left(1 + \frac{1}{2n}\right)^{2n} \right\}^{3/2}$$

conv. to e

Because this a subsequence

of  $\left(1 + \frac{1}{k}\right)^k$ .