

15.3 We learned how to  
integrate a function defined  
on a rectangle  $R$

$$\text{If } R = \left\{ (x, y) \mid \begin{array}{l} a \leq x \leq b \\ c \leq y \leq d \end{array} \right\}.$$

then

$$\iint_R f(x, y) dA = \int_a^b \int_c^d f(x, y) dy dx$$


or

$$z = \int_c^d \int_a^b f(x,y) dx dy$$

What if  $R$  is replaced by  
 a more complicated region?

We can write


$$F(x,y) = \begin{cases} f(x,y) & \text{if } (x,y) \text{ is in } D \\ 0 & \text{if } (x,y) \text{ is in } R \\ & \text{but not in } D \end{cases}$$



Geometrically

$$\iiint f(x,y) dA = \text{volume of region}$$

above  $D$ , under  $z = f(x,y)$




A plane region  $D$  is of type I

if it lies between the graphs

of two functions:

$$D = \{ (x, y) \mid a \leq x \leq b, g_1(x) \leq y \leq g_2(x) \}$$



$$= \int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx$$

y-integral

Ex Let  $D = \left\{ (x, y) \mid \begin{array}{l} 0 \leq x \leq 1 \\ x^2 \leq y \leq x \end{array} \right\}$

Compute  $\int_0^1 \int_{x^2}^x xy^2 dy dx$

$$= \int_0^1 \left. \frac{xy^3}{3} \right|_{y=x^2}^{y=x} dx$$

$$= \int_0^1 \left( \frac{x \cdot x^3}{3} - \frac{x \cdot x^6}{3} \right) dx$$


$$= \int_0^1 \frac{x^4}{3} - \frac{x^7}{3} dx$$

$$= \frac{x^5}{15} - \frac{x^8}{24} \Big|_0^1$$

$$= \frac{1}{15} - \frac{1}{24} = 120 \cdot \frac{8-5}{120} = \underline{\underline{\frac{1}{40}}}$$

A region  $D$  is of type II

$$\text{if } D = \left\{ (x, y) \mid \begin{array}{l} c \leq y \leq d \\ h_1(y) \leq x \leq h_2(y) \end{array} \right.$$



Ex. Let  $D$  be the region between

the graphs of  $x = y^2$  and  $x = 2y + 3$

Compute  $\iint_D y + x \, dA$

When do  $x=y^2$  and  $x=2y+3$

coincide?

$$y^2 = x = 2y + 3$$

$$\rightarrow y^2 - 2y - 3 = 0$$


$$(y-3)(y+1) = 0 \quad y = -1, y = 3$$

Which curve is above?

Plug into  $y=0$  into both

equations  $0 = x$   $2 \cdot 0 + 3 = y$

$\therefore 2y+3$  is  $>$   $y^2$





$$\rightarrow \int_{-1}^3 \int_{y^2}^{2y+3} y+x \, dx \, dy$$

$$= \int_{-1}^3 \left. xy + \frac{x^2}{2} \right|_{x=y^2}^{x=2y+3} dy$$

$$= \int_{-1}^3 (2y+3)y + \frac{(2y+3)^2}{2} - y^2 \cdot y - \frac{y^4}{2} \uparrow dy$$

$$= \int_{-1}^3 \frac{2y^2 + 3y}{3} + 2y^2 + 6y + \frac{9}{2} - y^3 - \frac{y^4}{2} dy$$

$$= \int_{-1}^3 \frac{-y^4}{2} - y^3 + \frac{8y^2}{3} + 9y + \frac{9}{2} dy$$

Usually it's better to avoid

square roots

$$y^2 = x = 2y$$

$$\rightarrow y^2 - 2y = 0 \rightarrow \begin{cases} y=0 \\ y=2 \end{cases}$$

$$\text{Plug } y=1 \quad y^2=1 \quad 2 \cdot 1 = 2$$

$\therefore x=2y$  is bigger (in  $x$ -direction)

$$\int_0^2 \int_{y^2}^{2y} xy \, dx \, dy$$

Ex. Sometimes only Type I or

Type II is possible

Evaluate  $\int_0^1 \int_x^1 \sin(y^2) dy dx$


$$= \iint_D \sin(y^2) dx dy$$

$$= \int_0^1 \int_x^1 \sin y^2 dx dy$$

$y=1$

$y=x$

$= ?$



$$\iint_D \sin(y^2) dA$$

$$= \int_0^1 \int_0^y \sin(y^2) dx dy$$

$$= \int_0^1 \left[ x \sin(y^2) \right]_{x=0}^{x=y} dy$$

$$= \int_0^1 y \sin(y^2) dy$$

$$= \frac{1}{2} \int_0^1 \sin(y^2) 2y dy$$

$$= -\frac{1}{2} \cos(y^2) \Big|_0^1$$

$$= -\frac{1}{2} \cos 1 + \frac{1}{2} \cos 0$$

$$= \frac{1}{2} (1 - \cos 1)$$



## Properties of Double Integrals

$$\iint_D [f(x,y) + g(x,y)] dA$$

$$= \iint_D f(x,y) dA + \iint_D g(x,y) dA$$

And:

$$\iint_D c f(x,y) dA = c \iint_D f(x,y) dA$$

If  $f(x,y) \geq g(x,y)$ , for  $(x,y)$  in  $D$ ,

then

$$\iint_D f(x,y) dA \geq \iint_D g(x,y) dA$$


Also if  $D = D_1 \cup D_2$ , where

$D_1$  and  $D_2$  don't intersect, then

(except on

boundaries)

$$\iint_D f(x,y) dA = \iint_{D_1} f(x,y) dA + \iint_{D_2} f(x,y) dA$$



$$D = D_1 \cup D_2$$

# 31. Find volume of region

bounded by the cylinder  $x^2 + y^2 = 1$

and the planes  $y = z$ ,  $x = 0$ , and

$z = 0$  in the first octant.