


15.7 Triple Integrals

$$\text{Given a box } B = \left\{ (x, y, z) \mid \begin{array}{l} a \leq x \leq b \\ c \leq y \leq d \\ r \leq z \leq s \end{array} \right\},$$

we divide B into sub-boxes

of width Δx , Δy , and Δz ,

$$\text{where } \Delta x = \frac{b-a}{k}, \quad \Delta y = \frac{d-c}{m}, \quad \Delta z = \frac{s-r}{n}$$



We define the triple Riemann sum by

$$\sum_{i=1}^l \sum_{j=1}^m \sum_{k=1}^n f(x_i, y_j, z_k) \Delta x \Delta y \Delta z,$$

where (x_i, y_j, z_k) is in the box

$$B_{ijk} = [x_{i-1}, x_i] \times [y_{j-1}, y_j] \times [z_{k-1}, z_k]$$

If we let $m \rightarrow \infty$, then

$$\iiint_B f(x, y, z) dV = \begin{array}{l} \text{limit of Riemann} \\ \text{sum as } l, m, n \\ \rightarrow \infty \end{array}$$

This is called the triple integral of f over the box B . To calculate

it, we use:

Fubini's Thm.

$$\iiint_B f(x, y, z) = \int_a^b \int_c^d \int_r^s f(x, y, z) dx dy dz$$

Integrate first with respect to

x , then y , then z .

Or one could integrate first with respect

to y , then z , then x

$$= \int_a^b \int_n^s \int_c^d f(x, y, z) dy dz dx, \text{ etc.}$$

Ex. If $B = \left\{ (x, y, z) \mid 0 \leq x \leq 2, 1 \leq y \leq 2, 0 \leq z \leq 1 \right\}$

calculate $\int \int \int_B xyz^2 dV$

$$= \int_0^2 \int_1^2 \int_0^1 xy^2z \, dz \, dy \, dx$$

$$= \int_0^2 \int_1^2 \left. \frac{xy^2z^2}{2} \right|_0^1 dy \, dx$$

$$= \int_0^2 \int_1^2 \frac{xy^2}{2} dy \, dx$$

$$= \int_0^2 \left. \frac{xy^3}{6} \right|_{y=1}^{y=2} dx$$

$$= \int_0^2 \left(\frac{x \cdot 4}{3} - \frac{x}{6} \right) dx = \int_0^2 \frac{x}{2} dx$$

$$= \frac{x^2}{4} \Big|_0^2 = \underline{\underline{1}}$$

A solid region E is said to be of

type 1, if it lies between the

graphs of 2 functions of x and y

over a region D :

$$\therefore E = \left\{ (x, y, z) \mid (x, y) \in D, u_1(x, y) \leq z \leq u_2(x, y) \right\}$$

$$\iiint_E f(x, y, z) dV = \iint_D \left[\int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) dz \right] dA$$

If D is a type II region, then

$$\iiint_E f(x, y, z) dV = \int_c^d \int_{h_1(y)}^{h_2(y)} \int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) dz \, dx \, dy$$

or if D is of type I:

$$\iiint_E f(x, y, z) dV = \int_a^b \int_{g_1(x)}^{g_2(x)} \int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) dz \, dy \, dx$$


Ex. Let $D =$ solid region bounded

by $z = x^2 + 2y + 1$ and $z = y + 2$


in the first octant. Find Vol.

$$x^2 + 2y + 1 = y + 2$$

$$\rightarrow y = 1 - x^2$$



$$\text{Vol} = \iiint 1 \, dV$$



$$= \int_0^1 \int_0^{1-x^2} \int_{x^2+2y+1}^{y+2} 1 \, dz \, dy \, dx$$

$$= \int_0^1 \int_0^{1-x^2} z \Big|_{x^2+2y+1}^{y+2} dy dx$$

$$= \int_0^1 \int_0^{1-x^2} (y+2) - (x^2+2y+1) dy dx$$

$$= \int_0^1 \int_0^{1-x^2} (1-x^2-y) dy dx$$

~~$$= \int_0^1 \left[y - x^2 y - \frac{y^2}{2} \right]_0^{1-x^2} dx$$~~

$$= \int_0^1 \left[y(1-x^2) - \frac{y^2}{2} \right]_0^{1-x^2} dx$$

$$= \int_0^1 (1-x^2)^2 - \frac{(1-x^2)^2}{2} dx$$

$$= \int_0^1 \frac{1}{2} (1-x^2)^2 dx$$

$$= \frac{1}{2} \int_0^1 (1 - 2x^2 + x^4) dx$$

$$= \frac{1}{2} \left(x - \frac{2x^3}{3} + \frac{x^5}{5} \right) \Big|_0^1 = \frac{4}{15}$$

Ex. Let $R =$ triangular region

in the xy -plane between

$y=x$ and $y=1$ for $0 \leq x \leq 1$,

and let $E =$ solid region

between the surfaces


$z = -2$ and $z = 1 - x^2$ for $(x,y) \in R$

Evaluate $\int \int \int_E (1-x^2 - (-2)) dz dA$

$\int_{-2}^{1-x^2}$

Evaluate $\int \int \int_E (x+1) dV.$

$$= \int_0^1 \int_x^1 \int_{-2}^{1-x^2} (x+1) dz dy dx$$



$$= \int_0^1 \int_x^1 (x+1)z \Big|_{-2}^{1-x^2} dy dx$$

$$= \int_0^1 \int_x^1 (x+1) (1-x^2 - (-2)) dy dx$$

$$= \int_0^1 \int_x^1 (-x^3 - x^2 + 3x + 3) dy dx$$

$$= \int_0^1 \left(-x^3 - x^2 + 3x + 3 \right) y \Big|_x^1$$

$$= \int_0^1 \left(x^4 - 4x^2 + 3 \right) dx = \frac{28}{15}$$

Ex. Suppose that a tetrahedron is



bounded by $x+2y+z=2$, $x=0$, $y=0$

and $z=0$.

and that the density

is $\rho = 2z$. Calculate mass m

$$\text{Set } z=0 \rightarrow x+2y=2 \rightarrow y=1-\frac{x}{2}$$



$$= \int_0^2 \int_0^{1-\frac{x}{2}} \left(1 - \frac{x}{2} - y\right)^2 dy dx$$

$$= \int_0^2 \int_0^{1-\frac{x}{2}} \left(y + \frac{x}{2} - 1\right)^2 dy dx$$

$$= \int_0^2 \left. \frac{\left(y + \frac{x}{2} - 1\right)^3}{3} \right|_{y=0}^{y=1-\frac{x}{2}} dx$$

$$= \int_0^2 \frac{\left(1 - \frac{x}{2} + \frac{x}{2} - 1\right)^3}{3} - \frac{\left(\frac{x}{2} - 1\right)^3}{3} dx$$

$$= \int_0^2 \left(1 - \frac{x}{2}\right)^3 dx$$

$$= \int_0^2 \left(1 - \frac{3x}{2} + 3\frac{x^2}{4} - \frac{x^3}{8}\right) dx$$

$$= \left. x - \frac{3x^2}{4} + \frac{x^3}{4} - \frac{x^4}{32} \right|_0^2$$

$$= 2 - 3 + 2 - \frac{1}{2} = \underline{\underline{\frac{1}{2}}}$$

Various Applications

$$\text{Vol}(E) = \iiint_E dV = \iiint_E 1 dV$$

Center
of Mass = $(\bar{x}, \bar{y}, \bar{z})$.

where $\frac{M_{yz}}{m} = \bar{x}$, $\frac{M_{xz}}{m} = \bar{y}$.

and $\frac{M_{xy}}{m} = \bar{z}$

where $M_{yz} = \iiint x \rho \, dV$

$M_{xz} = \iiint y \rho \, dV$, and

$M_{xy} = \iiint z \rho \, dV$

Ex. Suppose that $E =$ solid tetrahedron


bounded by $2x + y + 2z = 2$, $x = 0$, $y = 0$,

and $z = 0$. Suppose that the density

is $= x$. Calculate mass.

$$m = \int_0^2 \int_0^{2-2x} \int_0^{\frac{2-2x-y}{2}} x \, dz \, dy \, dx$$

Set $z=0 \rightarrow 2x+y=2$



solve for z

$$z = \frac{2 - 2x - y}{2}$$

$$m = \int_0^2 \int_0^{2-2x} xz \Big|_0^{1-x-y/2}$$

$$= \int_0^2 \int_0^{2-2x} x \left(1-x-\frac{y}{2}\right) dy dx$$

$$= \int_0^2 \int_0^{2-2x} (x-x^2) - \frac{xy}{2} dy dx$$

$$= \int_0^2 (2-2x)(1-x^2) dx - \int_0^2 \int_0^{2-2x} \frac{xy^2}{4} dx$$

$$= \int_0^2 2x^3 - 2x^2 - 2x + 2 - \int_0^2 \frac{xy^3}{12} \Big|_0^{2-2x}$$

$$= \left. \frac{x^4}{2} - \frac{2x^3}{3} - x^2 + 2x \right|_0^2 \rightarrow = \frac{1}{2} \cdot 2^4 - \frac{2}{3} \cdot 2^3 - 2^2 + 2 \cdot 2$$

$$= \int_0^2 \frac{x}{12} (2-2x)^3 dx$$