

16.4 Green's Theorem

Let C be the boundary of a

region D . We adopt the

convention the curve C has


a positive orientation if

C has the counterclockwise orientation

Thus if C is parameterized

by $\vec{r}(t)$, $a \leq t \leq b$, then the

domain D should be on the left side



Green's Thm. Let C be positively oriented, and suppose also that the boundary of D is parameterized by a single closed curve C .

If P and Q have continuous partial derivatives, then

$$\int_C P dx + Q dy = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$

Sometimes you'll see the notation

$$\oint_C P dx + Q dy \text{ to emphasize that}$$

C is parameterized counter-clockwise.

In one variable, the Fund. Thm
of Calculus states that

$$\int_a^b F'(x) dx = F(b) - F(a)$$

On the interior
of $[a, b]$

On the boundary
of $[a, b]$

Why is Green's Thm. true?


We assume that D is
of type I. (Maybe later)

Ex. Use Green's Thm. to

evaluate $\int_C \begin{matrix} 2xy \, dx + x^2 \, dy \\ -x \end{matrix}$,

where C is the triangle with

vertices at $(0,0)$, $(1,1)$, $(1,3)$



$$P(x,y) = 2xy +$$

$$Q(x,y) = x^2 - x$$

$$\frac{\partial Q}{\partial x} = 2x - 1$$

$$\frac{\partial P}{\partial y} = 2x$$

6.

$$\therefore \int_C P dx + Q dy$$

$$= \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$$


$$= \iint_D \left((2x-1) - (2x) \right) dA$$

$$= - \iint_D 1 dA = - \text{Area} = - \left(1 \cdot 2 \cdot \frac{1}{2} \right) = -1$$

Ex. Compute $\int_C \cos y \, dx + x^2 \sin y \, dy$,

where C = rectangle with

vertices at $(0,0)$, $(5,0)$, $(5,2)$
and $(0,2)$.



$$P(x,y) = \cos y$$

$$Q(x,y) = x^2 \sin y$$

$$\int_C = \iint_R (2x \sin y) + \sin y \, dA$$

$$= \int_0^5 \int_0^2 (2x+1) \sin y \, dy \, dx$$

$$= \int_0^5 (2x+1) (-\cos y) \Big|_0^2$$

$$= \int_0^5 (2x+1) (-\cos 2 + 1) \, dx$$

$$= (1 - \cos 2) (x^2 + x) \Big|_0^5$$

$$= \underline{\underline{(1 - \cos 2) \cdot (30)}}$$

Ex. Use Green's Thm. to evaluate

$$\int_C \vec{F} \cdot d\vec{n}, \quad \text{where } \vec{F} = \langle y - \cos y, x \sin y \rangle$$

and $C = \text{circle}$

$$(x-3)^2 + (y+4)^2 = 4.$$

$$\text{cent.} = (3, -4)$$

$$\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA =$$

$$= \iint_D \sin y - (1 + \sin y) dA$$

$$= - \int_D 1 \, dA = -\pi \cdot 2^2 = \underline{\underline{-4\pi}}$$

Ex. Use Green's Thm. to find
the work done by the force

$$\vec{F} = x(x+y)\vec{i} + xy^2\vec{j} \quad \text{in moving}$$


a particle from the origin

along the x -axis, then along

the line segment to $(0,1)$ and

then back along the y-axis

to the origin



$$\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} = (y^2 - x)$$

$$\therefore \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) = \iint_D (y^2 - x) \, dA$$

$$= \int_0^1 \int_0^{1-x} (y^2 - x) \, dA$$

$$= \int_0^1 \left(\frac{y^3}{3} - xy \right) \Big|_0^{1-x}$$


$$\int_0^1 \frac{(1-x)^3}{3} - x(1-x) \, dx$$

$$= \int_0^1 \left\{ \frac{1}{3} - x + x^2 - \frac{x^3}{3} - x + x^2 \right\} dx$$


$$= \int_0^1 \left\{ \frac{1}{3} - 2x + 2x^2 - \frac{x^3}{3} \right\} dx$$

$$= x = \frac{1}{3} - 1 + \frac{2}{3} - \frac{1}{12} = \underline{\underline{-\frac{1}{12}}}$$

What about more complicated
regions



Note that the inside is
parameterized clockwise.



We can use Green's Thm.

to compute the area of a

domain D

There are 3 ways :

1. Set $P = 0$, $Q = x$

$$\therefore Q_x - P_y = 1 - 0 = 1$$

2. Set $P = -y$, $Q = 0$

$$\therefore Q_x - P_y = 0 - (-1) = 1$$

or

$$3. \quad Q = \frac{1}{2}x, \quad P = -\frac{1}{2}y$$

$$\therefore Q_x - P_y = \frac{1}{2} - \left(-\frac{1}{2}\right) = 1$$

Ex. Find area of ellipse. (Use 3)

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$A = \frac{1}{2} \int_C X dy - Y dx = \frac{1}{2} \iint_D 1+1 dA = A$$

$$\text{Set } x = a \cos t \quad y = b \sin t$$

$$A = \frac{1}{2} \int_C x dy - y dx$$

$$= \frac{1}{2} \int_0^{2\pi} a \cos t \cdot b \cos t - b \sin t \cdot (-a \sin t) dt$$

$$= \frac{1}{2} \int_0^{2\pi} ab (\cos^2 t + \sin^2 t) dt$$

$$= \underline{\underline{\pi ab}}$$

Ex. 10 Let D be the region

between the circles $x^2 + y^2 = 4$

and $x^2 + y^2 = 9$

$C =$ boundary of D



$$\int_C (1 - y^3) dx + (x^2 + e^{y^2}) dy$$

$$= \iint_D (3x^2 + 0) - (0 - 3y^2) dA$$

$$= 3 \iint_D x^2 + y^2 dA$$

$$\frac{1}{2} \int_C x dy - y dx$$

$$= \frac{1}{2} \int_0^{2\pi} a \cos t (-\sin t) - b \sin t dt.$$