

2.1 Algebraic and Order

Properties of \mathbb{R} .

On \mathbb{R} , there are two

operations, addition +

multiplication. They satisfy:

$$(A_1) \quad a+b = b+a, \quad \begin{cases} \text{commutative} \\ \text{addition} \end{cases}$$

$$(A_2) \quad (a+b)+c = a+(b+c) \quad \begin{cases} \text{associative} \\ \text{addition} \end{cases}$$

(A₃) There is an element 0

in \mathbb{R} so $a+0 = a$
(0-element exists)

(A4) For each a in \mathbb{R} , there is

an element $-a$ in \mathbb{R} so

that

$$a + (-a) = 0 \text{ and } (-a) + a = 0$$

{negative element}

(M1) $a \cdot b = b \cdot a$ {commutative}

{multiplication}

(M2) $(a \cdot b) \cdot c = a \cdot (b \cdot c)$

{associative}

{multiplication}

(M3) There is an element 1 in \mathbb{R}

so that $a \cdot 1 = 1 \cdot a = a$

{unit element}
exists}

(M4). For each $a \neq 0$ in \mathbb{R} ,

there exists an element

$\frac{1}{a}$ such that

$$a \cdot \left\{ \frac{1}{a} \right\} = 1 \text{ and}$$

$$\left\{ \frac{1}{a} \right\} \cdot a = 1$$

{existence
of reciprocal}

$$(D) \quad a \cdot (b+c) = (a \cdot b) + (a \cdot c)$$

and

$$(b+c) \cdot a = (b \cdot a) + (c \cdot a)$$

(distributive property)

In a word, \mathbb{R} is a field

By applying some of the

above properties, one

can show that the

(1) zero element 0, the

(2) unit element 1, and

(3) the reciprocal $\frac{1}{a}$ are

all unique.

For example, suppose $a \neq 0$

and $a \cdot b = 1$. Then

$$b = 1 \cdot b = \left(\left(\frac{1}{a} \right) \cdot a \right) \cdot b$$

(M3) (M4)

$$= \left(\frac{1}{a} \right) \cdot \{ a \cdot b \} = \left(\frac{1}{a} \right) \cdot 1 = \frac{1}{a}$$

(M2) (D) (M3)

This proves (3)

Also, if $a \in \mathbb{R}$, then $a \cdot 0 = 0$

In fact,

$$a + a \cdot 0 = a \cdot 1 + a \cdot 0 = a \cdot (1+0)$$

$$\text{by } (M_3) \qquad \qquad \text{by } (D)$$

$$= a \cdot 1 = a$$

$$\text{by } (A_3) \qquad \text{by } (M_3)$$

Adding $(-a)$ to both sides, we get

$$a \cdot 0 = 0.$$

$$\text{Also, } 0 = (-1)(-1+1) = (-1)(-1) + (-1).$$

Adding 1 to both sides, we get

$$(-1)(-1) = 1$$

We define subtraction by

$$a - b = a + (-b)$$

and also we write

$$ab = a \cdot b,$$

and $a^2 = aa$ and

$$a^3 = a^2 a \text{ and}$$


$$a^{n+1} = a^n a, \text{ etc.}$$

\mathbb{Q}, \mathbb{R} are both fields.

Thm. There does not exist

a rational number r such

that $r^2 = 2$



Suppose by contradiction

that $r = \frac{p}{q}$. Then

$$r^2 = \left(\frac{p}{q}\right)^2 = 2 \rightarrow p^2 = 2q^2.$$

We can assume that

p and q have no common

factor. Then at most one
of p and q is even.

Since $p^2 = 2q^2$, we see

that p^2 is even. This implies

that p is also even (because

if $p = 2n+1$ is odd, then

$$p^2 = 4n^2 + 4n + 1 \text{ is also odd.}$$
)

Hence we can write $p = 2m$,

so that

$$p^2 = 4m^2 = 2q^2.$$

Dividing by 2,

$$2m^2 = q^2.$$

Hence q^2 must be even,

which implies q is even.

This shows that both

p and q are even, which

is a contradiction.

It follows that

\mathbb{R} must include numbers

that are ; irrational

(i.e., not rational).

For this purpose we need to
study Order Properties.

i.e., $<$ and $>$.

Order Properties of \mathbb{R}

There is a nonempty subset

\mathbb{P} of \mathbb{R} , called the set of positive real numbers such that

(i) If $a, b \in \mathbb{P}$, then $a+b \in \mathbb{P}$

(ii) If $a, b \in \mathbb{P}$, then $ab \in \mathbb{P}$

(iii) If $a \in \mathbb{P}$, then exactly one of the following holds:

$a \in \mathbb{P}$, $a=0$, $-a \in \mathbb{P}$

Trichotomy Property

If $\underline{-a \in P}$, we say a is negative,

and we write $\underline{a < 0}$ or $\underline{0 > a}$.

If $\underline{a \in P}$, we write $\underline{a > 0}$

or $\underline{0 < a}$

If $\underline{a \in P \cup \{0\}}$, we write $\underline{a \geq 0}$.

If $\underline{-a \in P \cup \{0\}}$, then we
write $\underline{a \leq 0}$.

If (i)-(iii) hold, then we say

\mathbb{R} is an ordered field.

Applying the Trichotomy Property
to $a-b$, we get

If $a-b \in \mathbb{P}$, i.e. $a > b$.

If $-(a-b) \in \mathbb{P}$, then $(b-a) \in \mathbb{P}$

$\Rightarrow b > a$

If $a-b=0$, then $a=b$

Here are the Rules for

Inequalities :

Thm. Let $a, b, c \in \mathbb{R}$.

2.1.7

(a) If $a > b$ and $b > c$, then

$$\underline{a > c}$$

(b) If $a > b$, then $a+c > b+c$

(c) If $a > b$ and $c > 0$, then

$$\underline{ca > cb}$$

If $a > b$ and $c < 0$, then

$$\underline{ac < bc}$$

Proof of (a): $a-b > 0$, $b-c > 0$

$$\text{then } (a-b)+(b-c) > 0$$

$$\text{or } a-c > 0 \rightarrow a > c$$

(b) If $a-b > 0$, then

$$(a+c) - (b+c) = a-b > 0$$

$$\rightarrow a+c > b+c$$

(c) If $a > b$ and $c > 0$, then

$$ca - cb = c(a-b) > 0.$$

$$\rightarrow ca > cb$$

If $c < 0$, then $-c > 0$. Hence

$$c(b-a) = -c(a-b) > 0$$

$$\rightarrow cb - ca > 0 \rightarrow cb > ca.$$

The Order Properties

in 2.1.5. and 2.1.6 lead to

2.1.10 and 2.1.11, which are

useful for solving inequalities:

1. Suppose that $ab > 0$. If

$a > 0$, then $b > 0$.

2. If $ab > 0$ and $a < 0$, then $b < 0$

3. If $ab < 0$ and $a > 0$, then $b < 0$

4. If $ab < 0$ and $a < 0$, then $b > 0$

Finally, we need to prove several facts:

Thm 2.1.8

(a) if $a \in \mathbb{R}$ and $a \neq 0$, then

$$a^2 > 0$$

(b) if $n \in \mathbb{N}$, then $n > 0$

Since $1 = 1^2$, (a) $\Rightarrow 1 > 0$

(c) If $n \in \mathbb{N}$, then $n > 0$.

Apply (b) and (i) from Order

Properties. Use Math. Ind.

(d) If $a > 0$, then $a^{-1} > 0$.

(e) If $0 < a < b$, then

$$a^{-1} > b^{-1}.$$

Pf. of (d). Suppose that

$a^{-1} \leq 0$. Then

$$1 = a a^{-1} \leq a \cdot 0 = 0.$$

This contradiction shows

$$a^{-1} > 0.$$

Pf. of (e). If $0 < a < b$,

then $a^{-1} - b^{-1} = (ab)^{-1}(b-a) > 0$

Since $(ab)^{-1} > 0$ and $b-a > 0$,

we get $a^{-1} > b^{-1}$.

22.

Ex. Find all real numbers x
 such that $3x + 4 \leq 12$.

Justify each step.

$$3x + 4 \leq 12 \Leftrightarrow 3x \leq 8 \Leftrightarrow x \leq \frac{8}{3}$$

↑ ↑

By (b) of 2.1.7

By (c) of 2.1.7

Ex. Solve $x^2 - 4x - 5 < 0$.

$$x^2 - 4x - 5 = (x-5)(x+1) < 0$$

\Leftrightarrow

By Property
 (3) above

If $x-5 > 0$, then $x+1 < 0$



No solution.

Or, by Property 4, if

$x-5 < 0$, then $x+1 > 0$.

\therefore Solution is $-1 < x < 5$.



Finally, we have

~~Thm. 2.1.8 :~~

(a) if $a \in \mathbb{R}$ and $a \neq 0$,

~~then~~ $a^2 > 0$.

~~(b) $l \neq 0$. Since $l = l^2$~~

~~this follows from (a)~~

A

We will define \mathbb{R} as
the set of infinite
decimal expansions:

$$x = \pm B. b_1 b_2 \dots ,$$

where B is a non-negative
integer and b_j is the
coefficient of 10^{-j} and

$$0 \leq b_j \leq 9$$

B

For example,

$$\pi = 3.14159265\dots$$

$$e = 2.71828182845\dots$$

$$\sqrt{2} = 1.4142135623\dots$$

It turns out that

rational numbers are

those decimal expansions

that are periodic.

C

Express $x = 45.2343434\dots$

Multiply by 10.

$$10x = 452.3434\dots$$

Multiply $10x$ by 100

$$1000x = 45234.3434\dots$$

Subtract:

$$990x = (45234 - 452)$$

$$x = \frac{44782}{990}.$$