

3.6 Properly Divergent Series

Let (x_n) be a sequence.

(i) We say (x_n) tends to

$+\infty$ and write $\lim (x_n) = +\infty$

if for every $\alpha \in \mathbb{R}$, there

exists a nat. number $K(\alpha)$

such that if $n \geq K(\alpha)$,

then $x_n > \alpha$.

(iii) We say $\{x_n\}$ tends to $-\infty$

and write $\lim (x_n) = -\infty$

if for every $B \in \mathbb{R}$,

there exists

a nat. number $K(B)$ such

that if $n \geq K(B)$, then

$x_n < B.$

In either case, we say $\{x_n\}$

is properly divergent.

Ex. $\lim(n) = +\infty$,

because if α is given,

let $K(\alpha)$ be any natural

number & such that $K(\alpha) > \alpha$.

If $n \geq K(\alpha)$, then $n > \alpha$.

Ex. $\lim(n^2) = +\infty$ Because

if $K(\alpha) > \alpha$, and if $n \geq K(\alpha)$

then $n^2 \geq n > \alpha$.

Ex. If $c > 1$, then $\lim c^n = +\infty$

In fact, let $c = 1+b$. If

α is given, let $K(\alpha)$ be a

natural number such that

$K(\alpha) > \frac{\alpha}{b}$. If $n \geq K(\alpha)$,

it follows from Bernoulli's

Inequality that

$$c^n = (1+b)^n \geq 1+nb > 1+\alpha > \alpha.$$

↑

Note that the inequalities

$n \geq K(\alpha) > \frac{\alpha}{b}$ imply that

$$n > \frac{\alpha}{b} \iff nb > \alpha.$$

Recall that the Monotone

Convergence Thm states

that a monotone sequence

is convergent if and only if

it's bounded.

Similarly, we have:

Thm. A monotone sequence

is properly divergent if and only if it is unbounded.

(a) If (x_n) is an unbounded

sequence, then $\lim(x_n) = +\infty$
increasing

(b) If (x_n) is an unbounded

decreasing sequence, then

$$\lim(x_n) = -\infty.$$

Comparison Test:

Thm. Let (x_n) and (y_n)

be two sequences and

suppose that $x_n \leq y_n$, all $n \in \mathbb{N}$

(a) If $\lim(x_n) = +\infty$,

then $\lim(y_n) = +\infty$

(b) If $\lim(y_n) = -\infty$, then

$\lim(x_n) = -\infty$

Ex. $\lim (\sqrt{n}) = +\infty$.

Let $K(\alpha)$ be any natural number.

with $K(\alpha) > \alpha^2$. If

$n \geq K(\alpha)$, then $n > \alpha^2$.

which implies $\sqrt{n} > \alpha$.

Compute $\lim (\sqrt{n+2})$

Note that if we use the same $K(\alpha)$ as above,

Then, if $n > \alpha^2$, then

$$\sqrt{n+2} > \sqrt{n} > \alpha.$$

which implies $\lim(\sqrt{n+2}) = +\infty$.

OR, we could have used the
above convergence test,

$$\text{With } x_n = \sqrt{n} \text{ and } y_n = \sqrt{n+2}.$$

Since $\lim(\sqrt{n}) = +\infty$, we get

$$\lim(\sqrt{n+2}) = +\infty.$$

Compute $\lim \left(\frac{\sqrt{n^2+1}}{\sqrt{n}} \right)$.

$$\frac{\sqrt{n^2+1}}{\sqrt{n}} \rightarrow \frac{\sqrt{n^2}}{\sqrt{n}} = \sqrt{n}.$$

↓ ↓
Set y_n Set x_n

\therefore Comp Test $\Rightarrow \lim \left(\frac{\sqrt{n^2+1}}{\sqrt{n}} \right) = +\infty$.

Ex. What about $\frac{\sqrt{n}}{(n^2+1)}$

Note that $\frac{\sqrt{n}}{(n^2+1)} < \frac{\sqrt{n}}{n^2} < \frac{n}{n^2} = \frac{1}{n}$.

Since $\lim \frac{1}{n} = 0$,

so does $\lim \frac{\sqrt{n}}{(n^2+1)} = 0$

Limit Comparison Test.

Suppose $\{x_n\}$ and $\{y_n\}$

are positive, and that

$$\lim \left\{ \frac{x_n}{y_n} \right\} = L \neq 0.$$

Then $\lim x_n = +\infty$

if and only if $\lim y_n = +\infty$

$$\text{Set } x_n = \frac{\sqrt{2n^2+1}}{\sqrt{3n-1}}$$

$$\text{and } y_n = \frac{n}{\sqrt{n}} = \sqrt{n}.$$

Check $\lim \frac{x_n}{y_n}$.

$$\frac{x_n}{y_n} = \frac{\frac{\sqrt{2n^2+1}}{\sqrt{3n-1}}}{\sqrt{n}}$$

$$= \frac{\sqrt{2n^2+1}}{\sqrt{n} \cdot \sqrt{3n-1}}$$

$$= \frac{n \sqrt{2 + \frac{1}{n^2}}}{\sqrt{n} \cdot \sqrt{n} \cdot \sqrt{3 - \frac{1}{n}}}$$

$$= \sqrt{\frac{2 + \frac{1}{n^2}}{3 - \frac{1}{n}}} \rightarrow \sqrt{\frac{2}{3}}$$

Proof of Limit Comparison Test.

We have $\lim \frac{x_n}{y_n} = L > 0$.

Set $\epsilon = \frac{L}{2}$.

$$\rightarrow L - \frac{L}{2} < \frac{x_n}{y_n} < L + \frac{L}{2}$$

if n is $> K_1$.

$$\rightarrow \frac{L}{2} < \frac{x_n}{y_n} < \frac{3L}{2}$$

$$\text{or } \frac{L}{2} y_n < x_n < \frac{3L}{2} y_n$$

Hence the usual Comparison

Test implies :

If $\lim y_n = +\infty$, then

$$\lim x_n = +\infty.$$

and if

$\lim x_n = +\infty$, then

$$\lim y_n = +\infty.$$