

3.1 Sequences

A sequence X is a function from \mathbb{N} to \mathbb{R} . Sometimes X is defined by a formula for the n -th term x_n such as

$$x_n = \frac{2^n}{n+1} \cdot \text{ Sometimes we just}$$

define the first few terms,

$$X = \left\{ \frac{1}{3}, \frac{1}{5}, \frac{1}{7}, \dots \right\} \text{ or}$$

$$x_n = \frac{1}{2n+1}$$

We can also give a recursive formula for x_n :

$$x_n = \frac{x_{n-1}}{x_{n-1}^2 + 1} \quad x_1 = 3.$$

It is very important to compute the limit of a sequence.

Definition. We say a sequence X converges to x if for all $\varepsilon > 0$, there is a number K in N , so that if $n \geq K$, then $|x_n - x| < \varepsilon$.

The number x is the limit of X , and we say X is convergent.

If X is not convergent, we say

X is divergent.

A sequence can only have at most

one limit. Suppose $\lim X = x'$

and $\lim X = x''$. Set $\varepsilon = \frac{|x'' - x'|}{2}$.

Choose K_1 so $|x_n - x'| < \varepsilon$
if $n \geq K_1$

and choose K_2 so that

$$|x_n - x''| < \epsilon \text{ if } n > K_2.$$

Now set $K = \max\{K_1, K_2\}$.

Then if $n \geq K$,

$$\{x' - x''\} = \{(x' - x_n) + (x'' - x_n)\}$$

$$\leq \{x' - x_n\} + \{x'' - x_n\}$$

$$< \epsilon + \epsilon = 2\epsilon$$

$$= |x' - x''|.$$

Dividing by $|x' - x''|$ we get $1 < 1$.

The contradiction implies:

$$x' = x''.$$

Some examples:

Compute $\lim \frac{1}{n}$.

We proved that for any $\epsilon > 0$,

there is a K so that if $n \geq K$,

$\frac{1}{n} < \epsilon$. We obtain that

$$\left| \frac{1}{n} - 0 \right| = \frac{1}{n} < \epsilon. \text{ It follows}$$

$$\text{that } \left| \lim \left(\frac{1}{n} \right) \right| = 0.$$

Ex. Prove that $\lim\left(\frac{3}{n+5}\right) = 0$.

Note that $\frac{3}{n+5} < \frac{3}{n}$.

For a given $\epsilon > 0$, choose $K > 0$

so that if $n \geq K$, then $\frac{1}{n} < \frac{\epsilon}{3}$.

If $n \geq K$, then

$$\begin{aligned} \left| \frac{3}{n+5} - 0 \right| &= \frac{3}{n+5} < \frac{3}{n} < 3 \cdot \frac{\epsilon}{3} \\ &= \epsilon. \end{aligned}$$

Hence $\lim\left(\frac{3}{n+5}\right) = 0$.

Ex. Show that $\lim (-1)^n$ does not exist.

Assuming $\lim (-1)^n = x$,

set $\epsilon = 1$. Then there

is a $K \in \mathbb{N}$ so that if $n \geq K$,

then $\{|(-1)^n - x|\} < 1$.

If n is even and $\geq K$, then

$$\{|x - 1| < 1 \rightarrow x - 1 > -1 \rightarrow x > 0\}$$

If n is odd and $\geq K$, then

$$|x+1| = |x - \{-1\}^n| < 1.$$

Hence, $x+1 < 1$, which

implies that $x < 0$.

This contradiction implies

that $\lim \{-1\}^n$ does not exist.

3.2. Limit Theorems.

Using the results of this section, we can analyse the convergence of many sequences.

Definition. A sequence $X = (x_n)$

is bounded if there exists

a number $M > 0$ such that

$$|x_n| \leq M, \quad \text{for all } n \in N.$$

Thm. A convergent sequence
of real numbers is bounded.

Pf. Suppose that $\lim x_n = x$

and let $\epsilon = 1$. Then there is

a $K \in \mathbb{N}$ such that $|x_n - x| < 1$

for all $n \geq K$. The Triangle

Inequality with $n \geq K$ implies

that

$$\begin{aligned} |x_n| &= |x_n - x + x| \leq |x_n - x| + |x| \\ &< 1 + |x|. \end{aligned}$$

If we set

$$M := \max \left\{ |x_1|, |x_2|, \dots, |x_{K-1}|, 1 + |x| \right\}$$

then it follows that

$$|x_n| \leq M, \quad \text{for all } n \in \mathbb{N}.$$

We want to learn how

taking limits interacts

with the operations of

addition, subtraction,

multiplication and division.

Given two sequences $X = (x_n)$

and $Y = (y_n)$, we define

$$X + Y = (x_n + y_n)$$

$$X - Y = (x_n - y_n)$$

$$XY = (x_n y_n)$$

$$cX = (cx_n)$$

and

$$X/Y = \left\{ \frac{x_n}{y_n} \right\} \quad \begin{array}{l} \text{(provided)} \\ y_n \neq 0 \end{array}$$

Suppose $X = (x_n)$ and $Y = (y_n)$

converge to x and y

respectively. Let $\epsilon > 0$.

Addition.

Choose K_1 and K_2 so that

$$|x_n - x| < \frac{\epsilon}{2} \text{ if } n \geq K_1 \quad \text{and}$$

$$|y_n - y| < \frac{\epsilon}{2} \text{ if } n \geq K_2.$$

Now set $K = \max\{K_1, K_2\}$

If $n \geq K$, then $n \geq K_1$ and
 $n \geq K_2$. Hence,

$$\begin{aligned} & |(x_n + y_n) - (x + y)| \\ &= |(x_n - x) + (y_n - y)| \\ &\leq |x_n - x| + |y_n - y| \\ &\leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

Hence $\lim (x_n + y_n) = x + y$.

For subtraction, we use the same argument. Just replace

$x_n + y_n$ by $x_n - y_n$ and

$x + y$ by $x - y$.

Multiplication. This is a bit

more complicated. Note that

$$|x_n y_n - xy| = |(x_n y_n - x_n y) + (x_n y - xy)|$$

$$\leq |x_n(y_n - y)| + |(x_n - x)y|$$

$$\leq |x_n| |y_n - y| + |x_n - x| |y|$$

By the boundedness theorem,

there is $M_1 > 0$ such that

$$|x_n| \leq M_1, \quad \text{all } n.$$

Now set $M = \max\{M_1, |y|\}$.

We conclude that

$$|x_n y_n - xy| \leq M |y_n - y| + M |x_n - x|$$

Now let $\epsilon > 0$ be given.

Then there exists K_1

such that

$$|x_n - x| < \frac{\epsilon}{2M} \quad \text{if } n \geq K_1.$$

Similarly, there exists K_2

such that

$$|y_n - y| < \frac{\epsilon}{2M} \quad \text{if } n \geq K_2.$$

Now set $K = \max\{K_1, K_2\}$

If $n \geq K$, then

$$|x_n y_n - xy|$$

$$\leq M|y_n - y| + M|x_n - x|$$

$$< M \cdot \frac{\epsilon}{2M} + M \cdot \frac{\epsilon}{2M} = \epsilon.$$

This proves

$$\lim (x_n y_n) = xy.$$

In order to study limits of sequences of quotients, we need the following result:

Proposition. Suppose that $\lim y_n = c$, where $c \neq 0$. Then there is a $K \in \mathbb{N}$ such that $|c_n| > \frac{|c|}{2}$, for all $n \geq K$.

Set $\epsilon = \frac{|c|}{2}$. Then there is a $K \in \mathbb{N}$ such that if $n \geq K$,

then $|y_n - c| < \frac{|c|}{2}$

20

By the Backwards Triangle

Property

$$|y_n| = |(y_n - c) + c|$$

$$\geq |c| - |y_n - c|$$

$$\geq |c| - \frac{|c|}{2} = \frac{|c|}{2}.$$

Now we can prove that generally, quotients of sequences have limits.

Division

Suppose (y_n) converges 21

to y , where $y \neq 0$. By

the above Property,

$$|y_n| > \frac{|y|}{2}, \text{ if } n \geq K.$$

Hence, if we set $c = y$,

$$\left| \frac{1}{y_n} - \frac{1}{y} \right| = \frac{|y - y_n|}{|y_n||y|}, \text{ so}$$

$$\begin{aligned} \left| \frac{1}{y_n} - \frac{1}{y} \right| &= \frac{|y_n - y|}{|y_n||y|} \\ &\leq \frac{2|y_n - y|}{|y|^2}, \end{aligned}$$

by the Property . Since

$y \neq 0$ and $\lim y_n = y \neq 0$,

there is a $K_2 \in N$. such that

$$|y_n - y| < \frac{|y|^2}{2} \epsilon, \text{ if } n > K_2.$$

If we set $K = \max\{K_1, K_2\}$

and if $n \geq K$, then

$$\left| \frac{1}{y_n} - \frac{1}{y} \right| < \frac{2|y_n - y|}{|y|^2} \cdot \frac{|y|^2}{2} \epsilon$$

$$= \epsilon.$$

Thus, if $\lim y_n = y$ and

$y \neq 0$, then $\lim \frac{1}{y_n} = \frac{1}{y}$.

For the general quotient rule, the Product Rule, implies that

$$\begin{aligned}\lim \frac{x_n}{y_n} &= \lim x_n \cdot \lim \frac{1}{y_n} \\ &= x \cdot \frac{1}{y} = \frac{x}{y},\end{aligned}$$

provided that $y \neq 0$.