

MA 440 (Honors)
Practice Problems For Final

NAME _____ ID number _____

The practice problems for final include the problems from homeworks, quizzes, midterms and the following. The majority problems on the final will be similar to the problems in these 4 sets of problems.

1. If $\xi \in \mathbb{R}$ is irrational and $r \in \mathbb{R}$ and $r \neq 0$, show $r + \xi$ is irrational.
2. If $a > -1, a \in \mathbb{R}$, show that $(1 + a)^n \geq 1 + na$ for all $n \in \mathbb{N}$ by using mathematical induction.
3. If $a > -1, a \in \mathbb{R}$, show that $(1 + a)^r \geq 1 + ra$ for all $r \geq 1$.
4. State the *Supremum Property*
5. Prove the *Archimedean Property*, namely, show for every $x \in \mathbb{R}$, there exists $n \in \mathbb{N}$, such that $x < n$.
6. State the *Nested Cells Property*
7. (Schwarz inequality) Let V be an inner product space. Define

$$\|x\| = \sqrt{x \cdot x} \text{ for } x \in V$$

show $x \cdot y \leq \|x\|\|y\|$.

8. Let S be a set in \mathbb{R}^p . State the definition that a point x is a boundary point of S . State the definition that a point x is a cluster point of S . What are the differences?
9. Give an example such that x is a cluster point, but not a boundary point. Also give an example that x is a boundary point, but not a cluster point.
10. Show that if F is closed, then any cluster point of F is in F .
11. Show that if F is closed, then any boundary point of F is in F .
12. Prove that the set of all cluster points of A which is a subset of \mathbb{R}^p is closed.
13. Show that if $S \subset \mathbb{R}$ is open, then it is the union of a countable collection of open intervals.
14. State the definition for a set K to be compact. Show directly from definition that $K = \{(x, y) : |x| + |y| < 1\}$ is not compact.

15. Show that if a set K in \mathbb{R}^p is compact, then it is bounded.
16. Show that if a set K in \mathbb{R}^p is compact, then it is closed.
17. Show that if a set K in \mathbb{R}^p is compact, then for a sequence (a_n) in K , if (a_n) converges to a , then a is in K .
18. Let D be a subset in \mathbb{R}^p , give the definition for D to be disconnected.
19. Using the fact that \mathbb{R}^p is connected, show that the only subsets of \mathbb{R}^p which are both open and closed are empty set ϕ and \mathbb{R}^p .
20. Let S be a subset in \mathbb{R}^n and denote ∂S be the set of all boundary points of S , Show that ∂S is closed.
21. Give an example that A and B are connected subsets in \mathbb{R}^p , but $A \cap B$ is disconnected.
22. Let K be a compact subset of \mathbb{R}^p and let x be any point in \mathbb{R}^p such that x is not in K . Prove that there exist open sets U and V , where U and V are disjoint, U contains K and V contains x .
23. Let K_1 and K_2 be compact subsets of \mathbb{R}^p . Then there exist $x_1 \in K_1$ and $x_2 \in K_2$ such that for all $z_1 \in K_1$ and $z_2 \in K_2$, $\|z_1 - z_2\| \geq \|x_1 - x_2\|$.
24. Show that if a monotone sequence (x_n) in \mathbb{R} is bounded, then it is convergent. Also $\lim_{n \rightarrow \infty} x_n = \sup x_n$.
25. Show *Bolzano-Weierstrass Theorem*. Namely, let (x_n) be a bounded sequence in \mathbb{R}^p contained infinite distinct values. Then it has a convergent subsequence.
26. State the definition for (x_n) to be a Cauchy sequence. Show that (s_n) where $s_n = \sum_{n=1}^{\infty} \frac{1}{n}$ is not a Cauchy sequence.
27. Show that if a bounded divergent sequence (x_n) must has two convergent subsequences which converge to different values.
28. Let $s_n = (-2)^{(-2)^n}$. Find $\limsup s_n$ and $\liminf s_n$ and justify your answer.
29. Let (x_n) be a positive sequence and $\lim_{n \rightarrow \infty} x_n^{1/n} < 1$, show that there exists a r with $0 < r < 1$, $0 \leq x_n < r^n$ for sufficiently large $n \in \mathbb{N}$.
30. Give the definition for $u \in \mathbb{R}$ to be an infimum of a non-empty subset S of \mathbb{R} .
31. Let $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ be given and satisfy

$$x_n \leq y_n \leq z_n, \quad \lim x_n = \lim z_n = L$$

Prove by definition $\lim y_n = L$.

32. Show that if $\sum a_n$ converges and $a_n \geq 0$, then $\sum \frac{\sqrt{a_n}}{n}$ converges.

33. Show that if $\sum a_n$ diverges and $a_n \geq 0$, then $\sum \frac{1+a_n}{a_n}$ diverges.
34. Let $f(x)$ be continuous, $K \subset D(f)$ and K is compact. Show $f(K)$ is bounded.
35. Let $f(x)$ be continuous, $K \subset D(f)$ and K is compact. Show $f(K)$ is closed.
36. Show that if $f(x)$ is a contraction from R^p to R^p , then $f(x)$ has a fixed point.
37. Show that if $(f_n(x))$ converges uniformly to $f(x)$ and $(f_n(x))$ are continuous on D , then $f(x)$ is continuous on D . (Where the “uniformly” is used?)
38. Show that if $f'_n(x)$ converges uniformly to $g(x)$ in $J = [a, b]$ and $f_n(x)$ converges at x_0 , then $f_n(x)$ converges to $f(x)$ where $f'(x) = g(x)$.

39. Let

$$g(x) = \begin{cases} x^2 & \text{for } 0 \leq x < 2 \\ x^3 & \text{for } 2 \leq x < 3 \end{cases}$$

Evaluate the Riemann-Stieltjes integral

$$\int_0^3 x dg(x)$$

and briefly justifying your computation.

40. Let

$$g_n(x) = \begin{cases} nx & \text{for } 0 \leq x \leq 1/n \\ \frac{n}{n-1}(1-x) & \text{for } 1/n < x \leq 1 \end{cases}$$

Show that (g_n) converges pointwise on $[0, 1]$ and find the limit function. Does it converge uniformly?

41. Let (x_n) be a sequence of real numbers such that $|x_n| \leq \frac{1}{2^n}$, and set $y_n = x_1 + x_2 + \cdots + x_n$. Show the sequence (y_n) converge.
42. If a sequence $(f_n(x))$ converges uniformly to a function $f(x)$ on $[a, b]$, and each $f_n(x)$ is continuous and bounded. Show that $f(x)$ is continuous and bounded.
43. If a sequence $(f_n(x))$ converges uniformly to a function $f(x)$ on $[a, b]$, and each $f_n(x)$ is continuous and bounded. Show directly by definition that $f(x)$ is uniform continuous.
44. Show that if f is continuous and bounded on $[a, b]$, then f is Riemann integrable.
45. Show that if f is a bounded function on $[0, 1]$ and if for every $a > 0$, f is Riemann integrable on $[a, 1]$, then f is integrable on $[0, 1]$.
46. State Taylor's Theorem. Give Taylor's Formula using 3 terms (including the remainder) with $f(x) = \sqrt{x}$ and $x_0 = 1$. In the remainder term, find the point at which the second derivative is evaluated.

47. Prove that if f has a continuous third derivative and satisfies $f(0) = f'(0) = f''(0) = 0$ and $f'''(x) \leq 1$ for $x \geq 0$, then $f(x) \leq x^3/3$ for $x \geq 0$.

48. Let

$$f_n = \frac{(-1)^n}{2^n} \cos(2\pi n x^2), x \in [0, 1], n \in \mathbb{N}$$

show that $\sum_{n=1}^{\infty} f_n(x)$ converges.

49. Prove or disprove the series $\sum_{n=1}^{\infty} \sin(n^{-2}) \cos(n^{-1})$ converges.

50. Let f, f', f'' be bounded and continuous in \mathbb{R} and $f(0) = f'(0) = 0$. Show that $\sum_{n=1}^{\infty} f(\frac{x}{n})$ converges.

51. Let $f(x) = \sum_{k=1}^{\infty} \frac{x^k}{k^2}$. Compute $f'(\frac{1}{3})$ and justify each steps which leads to the result.

52. Show there does not exist a continuous function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, such that

(a) $f\{(x, y) : |x| \leq 1, |y| \leq 2\} = \mathbb{Q} \cap [0, 1]$

(b) $f\{(x, y) : |x| \leq 1, |y| \leq 2\} = [0, \infty)$

(c) $f^{-1}\{x : |x| < 1\} = \{|x| \leq 1, |y| \leq 2\}$

(d) $f^{-1}\{x : |x| \leq 1\} = \{|x| < 1, |y| < 2\}$

53. Let A be a non-compact subset of the real line. Show that there exists a continuous function on A that is unbounded on A .

54. Prove that $2\pi \sin(x) = 1 + x^2$ has at least two real roots and locate disjoint intervals $(a, b), (c, d)$ which contain them.

55. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous and satisfy $\lim_{|x| \rightarrow \infty} f(x) = 0$. Show that $f(x)$ is uniformly continuous.

56. $f(x)$ is continuous on $[0, 1]$ Show that

$$h(x) = \sum \frac{f(x)^n}{(1 + |f(x)|)^n}$$

is also continuous on $[0, 1]$.

57. $f_n(x) = \frac{xn}{n+1}$ and let $f(x)$ be the limit function of $f_n(x)$. Find $f(x)$ and show that $f_n(x)$ does not converge to $f(x)$ uniformly.

58. Let (x_n) be a sequence in R^p with the property that there exists a real number $0 < r < 1$, and an integer N_0 such that

$$\|x_{n+1} - x_n\| \leq r \|x_n - x_{n-1}\| \text{ for } n \geq N_0$$

Then prove (x_n) converges.

59. Let (x_n) be a sequence in \mathbb{R}^p with the property that there exists an integer N_0 such that

$$\|x_{n+1} - x_n\| < \|x_n - x_{n-1}\| \text{ for } n \geq N_0$$

Can you show (x_n) converges? Justify your answer.

60. Let (x_n) be a sequence in a compact set $K \subset \mathbb{R}^p$ that is not convergent. Show there are two subsequences of this sequence that are convergent to different limit points.

61. Let (x_n) be an unbounded monotone increasing sequence, show that $\lim x_n = +\infty$.

62. True or False. Justify your answer.

- (a) Every sequence has a nondecreasing subsequence.
- (b) Every sequence has a bounded subsequence.
- (c) Every bounded sequence has a monotonic subsequence.
- (d) Every subsequence of a bounded monotonic sequence converges.
- (e) Every bounded sequence has a convergent subsequence.