

Disconjugacy and the Secant Conjecture

Alexandre Eremenko*

November 26, 2014

Abstract

We discuss the so-called secant conjecture in real algebraic geometry, and show that it follows from another interesting conjecture, about disconjugacy of vector spaces of real polynomials in one variable.

AMS Class 2010: 14P99, 34C10, 26C10. Keywords: disconjugacy, Wronskian, Schubert calculus.

Let V be a real vector space of dimension n whose elements are real functions on an interval $[a, b]$. The space V is called *disconjugate* if one of the following equivalent conditions is satisfied:

- a) Every $f \in V \setminus \{0\}$ has at most $n - 1$ zeros, or
- b) For every n distinct points z_1, \dots, z_n on $[a, b]$ and every basis f_1, \dots, f_n of V we have $\det(f_i(x_j)) \neq 0$.

One can replace “every basis” by “some basis” in b) and obtain an equivalent condition.

If V is disconjugate then the determinant in b) has constant sign which depends only on the ordering of x_j and on the choice of the basis.

A space of real functions on an open interval is called disconjugate if it is disconjugate on every closed subinterval.

We are only interested here in spaces V consisting of polynomials.

Suppose that a positive integer d is given, and V consists of polynomials of degree at most d . Then every basis f_1, \dots, f_n of V defines a real rational curve $\mathbf{RP}^1 \rightarrow \mathbf{RP}^{n-1}$ of degree d . Indeed, we can replace every $f_j(x)$ by a

*Supported by NSF grant DMS-1361836.

homogeneous polynomial $f_j^*(x_0, x_1)$ of two variables of degree d , such that $f_j(x) = f_j^*(1, x)$, and then f_1^*, \dots, f_n^* define a map $f : \mathbf{RP}^1 \rightarrow \mathbf{RP}^{n-1}$ (if polynomials have a common root, divide it out).

Then the geometric interpretation of disconjugacy is:

c) The curve f constructed from a basis in V is *convex*, that is intersects every hyperplane at most $n - 1$ times.

For every basis f_1, \dots, f_n in V we can consider its Wronski determinant $W = W(f_1, \dots, f_n)$. Changing the basis results in multiplication of W by a non-zero constant, so the roots of W only depend on V .

Conjecture 1. *Suppose that all roots of W are real. Then V is disconjugate on every interval that does not contain these roots.*

This is known for $n = 2$ with arbitrary d (see below), and for $n = 3, d \leq 5$ by direct verification with a computer.

This conjecture arises in real enumerative geometry (Schubert calculus), and we explain the connection. The problem of enumerative geometry we are interested in is the following:

Let $m \geq 2$ and $p \geq 2$ be given integers. Suppose that mp linear subspaces of dimension p in general position in \mathbf{C}^{m+p} are given. How many linear subspaces of dimension m intersect all of them?

The answer was obtained by Schubert in 1886 and it is

$$d(m, p) = \frac{1!2! \dots (p-1)!(mp)!}{m!(m+1)! \dots (m+p-1)!}.$$

Now suppose that all those given subspaces are real. Does it follow that all p -subspaces that intersect all of them are real? The answer is negative, and we are interested in finding a geometric condition on the given p -spaces that ensure that all $d(m, p)$ m -subspaces that intersect all of them are real.

One such condition was proposed by B. and M. Shapiro. Let $F(x) = (1, x, \dots, x^d), d = m + p - 1$ be a rational normal curve, a. k. a. moment curve. Suppose that the given p -spaces are osculating F at some real points $F(x)$. This means that subspaces X_j are spanned by the (row)-vectors $F(x_j), F'(x_j), \dots, F^{(p-1)}(x_j)$ for some real $x_j, 1 \leq j \leq mp$. Then all m -subspaces that intersect all X_j are real.

This was conjectured by B. and M. Shapiro and proved by E. Mukhin,

V. Tarasov and A. Varchenko (MTV) [6]. Earlier it was known for $n = 2$ [1], and in [2] a simplified elementary proof for the case $n = 2$ was given.

We are interested in the following generalization of this result.

Secant Conjecture. *Suppose that each of the mp subspaces X_j , $1 \leq j \leq mp$ is spanned by p distinct real points $F(x_{j,k})$, $0 \leq k \leq p - 1$, and that these sets of points are separated, that is $x_{j,k} \in I_j$, where $I_j \subset \mathbf{RP}^1$ are disjoint intervals. Then all m -subspaces which intersect all X_j are real.*

This is known when $p = 2$, [3] and has been tested on a computer for $p = 3$ and small m , [5, 4]. The special case when the groups $\{x_{j,k}\}_{k=0}^{p-1}$ form arithmetic progressions, $x_{j,k} = x_{j,0} + kh$ has been established [7].

Next we show how the Secant Conjecture follows from Conjecture 1 and the results of MTV.

Let us represent an m -subspace Y that intersects all subspaces X_j as the zero set of p linear forms, and use the coefficients of these forms as coefficients of p polynomials f_0, \dots, f_{p-1} . Then the condition that Y intersects some X_j is equivalent to linear dependence of the p vectors $f_i(x_{j,m})_{m=0}^{p-1}$, $i = 0, \dots, p - 1$. That is

$$\det (f_i(x_{j,m}))_{i,m=0}^{p-1} = 0.$$

These equations for $j = 1, \dots, mp$ define the subspaces Y , and we have to prove that all solutions are real.

Let I_j be the intervals with disjoint closures which contain the $x_{j,k}$. We may assume without loss of generality that $\infty \notin I_j$. We place on each I_j a point y_j , and consider the $d(m, p)$ real rational curves $\mathbf{R} \rightarrow \mathbf{RP}^p$ with inflection points at y_j . These curves exist by the MTV theorem, and they depend continuously on the y_j .

Let $f = (f_0 \dots, f_{p-1})$ be one of these curves. Fix $k \in \{1, \dots, mp\}$. For all $j \neq k$, fix all $y_j \in I_j$. When y_k moves on I_k from the left end to the right end, the determinant $\det(f_i(x_{k,m}))_{i,m=0}^{p-1}$ must change sign. So this determinant is 0 for some position of y_k on I_k .

Then it follows by a well-known topological argument that one can choose all $y_j \in I_j$ in such a way that $\det(f_i(x_{j,m})) = 0$ for all j .

Thus we have constructed $d(m, p)$ real solutions of the secant problem. As the total number of solutions is also $d(m, p)$, for generic data, we obtain the result.

Proof of Conjecture 1 for $n = 2$. We have two real polynomials f_1 and f_2 , such that $f_1'f_2 - f_1f_2'$ has only real zeros. This means that the rational

function $F = f_1/f_2$ is real and all its critical points are real. Let $I \subset \mathbf{R}$ be a closed interval without critical points. Then f is a local homeomorphism on I , so $F(I + i\epsilon)$ belongs to one of the half-planes $\mathbf{C} \setminus \mathbf{R}$, for all sufficiently small $\epsilon > 0$. Suppose without loss of generality that it belongs to the upper half-plane H . Let D be the component of $F^{-1}(H)$ that contains $F(I + i\epsilon)$. Then D is a region in H with piecewise analytic boundary, and $I \subset \partial D$. The map $F : D \rightarrow H$ is a covering because it is proper and has no critical points. As H is simply connected, D must be simply connected and $F : D \rightarrow H$ must be a conformal homeomorphism. Then $F^{-1} : H \rightarrow D$ is a conformal homeomorphism. As ∂D is locally connected, this homeomorphism extends to $F^{-1} : \overline{H} \rightarrow \overline{D}$. This last map must be injective because this is a left inverse of a function. Thus $F^{-1} : \overline{H} \rightarrow \overline{D}$ is a homeomorphism. Then $F : \overline{D} \rightarrow \overline{H}$ must be also a homeomorphism, in particular F is injective on I .

This implies that the linear span of f_1, f_2 is disconjugate.

References

- [1] A. Eremenko and A. Gabrielov, Rational functions with real critical points and the B. and M. Shapiro conjecture in real enumerative geometry, *Ann of Math.*, 155 (2002), 105–129.
- [2] A. Eremenko and A. Gabrielov, Elementary proof of the B. and M. Shapiro conjecture for rational functions, in the book: *Notions of positivity and the geometry of polynomials*, trends in mathematics, Springer, Basel, 2011, p. 167–178.
- [3] A. Eremenko, A. Gabrielov, M. Shapiro and A. Vainshtein, Rational functions and real Schubert calculus, *Proc. AMS*, 134 (2006), no. 4, 949–957.
- [4] L. Garcia-Puente, N. Hein, Ch. Hillar, A. del Campo, J. Ruffo, F. Sottile, Z. Teitler, The Secant Conjecture in the real Schubert calculus, *Exp. Math.* 21 (2012), no. 3, 252–265.
- [5] C. Hillar, L. Garca-Puente, A. Martn del Campo, J. Ruffo, Z. Teitler, S. Johnson, F. Sottile, Experimentation at the frontiers of reality in Schubert calculus. *Gems in experimental mathematics*, 365–380, *Contemp. Math.*, 517, Amer. Math. Soc., Providence, RI, 2010.

- [6] E. Mukhin, V. Tarasov and A. Varchenko, The B and M Shapiro conjecture in real algebraic geometry and the Bethe ansatz, *Ann of Math.*, 170 (2009) 2, 863–881.
- [7] E. Mukhin, V. Tarasov and A. Varchenko, On reality property of Wronski maps, *Confluentes Math.*, 1 (2009) 225–247.

*Department of Mathematics,
Purdue University,
West Lafayette, IN 47907
eremenko@math.purdue.edu*