

# On the “pits effect” of Littlewood and Offord

Alexandre Eremenko\* and Iossif Ostrovskii

October 13, 2007

## Abstract

Asymptotic behavior of the entire functions

$$f(z) = \sum_{n=0}^{\infty} e^{2\pi i \alpha_n} z^n / n!, \quad \text{with real } \alpha_n$$

is studied. It turns out that the Phragmén–Lindelöf indicator of such function is always non-negative, unless  $f(z) = e^{az}$ . For special choice  $\alpha_n = \alpha n^2$  with irrational  $\alpha$ , the indicator is constant and  $f$  has completely regular growth in the sense of Levin–Pfluger. Similar functions of arbitrary order are also considered.

MSC Primary: 30D10, 30D15, 30B10.

In [21] Nassif studied (on Littlewood’s suggestion) the asymptotic behavior and the distribution of zeros of the entire function

$$\sum_{n=0}^{\infty} e^{2\pi i n^2 \alpha} z^n / n!, \tag{1}$$

with  $\alpha = \sqrt{2}$ . This was continued by Littlewood [17, 18], who considered generalizations to Taylor series whose coefficients have smoothly varying moduli and arguments of the form  $\exp(2\pi i \alpha n^2)$ , where  $\alpha$  is a quadratic irrationality.

Such functions behave similarly to random entire functions previously studied by Levy [16] and Littlewood and Offord [19], in particular they display the “pits effect” which Littlewood described as follows:

---

\*Supported by the NSF grants DMS-0555279 and DMS-0244547.

“If we erect an ordinate  $|f(z)|$  at the point  $z$  of the  $z$ -plane, then the resulting surface is an exponentially rapidly rising bowl, approximately of revolution, with exponentially small pits going down to the bottom. The zeros of  $f$ , more generally the  $w$ -points where  $f = w$ , all lie in the pits for  $|z| > R(w)$ . Finally the pits are very uniformly distributed in direction, and as uniformly distributed in distance as is compatible with the order  $\rho$ ”.

The earliest study of functions (1) known to the authors is the thesis of Ålander [1] who considered the case of rational  $\alpha$ . Levy [16] used the results of Hardy and Littlewood on Diophantine approximation to prove the following. Let

$$M(r, f) = \max_{|z|=r} |f(z)| \quad \text{and} \quad m_2^2(r, f) = \frac{1}{2\pi} \int_{-\pi}^{\pi} |f(re^{i\theta})|^2 d\theta.$$

Then

$$M(r, f)/m_2(r, f) \quad \text{is bounded} \tag{2}$$

for  $f$  of the form (1), and  $\alpha$  satisfying a Diophantine condition. This is even stronger regularity than random arguments of coefficients yield [16, 19]. Some other works where the function (1) with various  $\alpha$  was studied or used are [7, 8, 20, 28].

Function (1) is the unique analytic solution of the functional equation

$$f'(z) = qf(q^2z), \quad \text{where} \quad q = e^{2\pi i\alpha}, \quad \text{and} \quad f(0) = 1, \tag{3}$$

which is a special case of the so-called “pantograph equation”. There is a large literature on this equation with real  $q$ , see, for example, [14, 13] and references there.

Recently there was a renewed interest to the functions of the type (1) because they arise as the limits as  $q \rightarrow e^{2\pi i\alpha}$  of the function of two variables

$$\sum_{n=0}^{\infty} q^{n^2} z^n / n!$$

which plays an important role in graph theory [27] and statistical mechanics [25]. This function is the unique solution of (3), for all  $q$  in the closed unit disc.

In the present paper, we first study arbitrary entire functions of the form

$$f(z) = \sum_{n=0}^{\infty} a_n z^n / n!, \quad \text{where} \quad |a_n| = 1. \tag{4}$$

Our Theorem 1 says that such functions cannot decrease exponentially on any ray, unless  $f$  is an exponential. This can be compared with a result of Rubel and Stolarski [23] that there exist exactly five series of the form (4) with  $a_0 = 0$ ,  $a_n = \pm 1$  which are bounded on the negative ray. Our second result, Theorem 2 shows that one cannot replace the condition of exponential decrease in Theorem 1 by boundedness on a ray: there are infinitely many functions of the form (4) which tend to zero as  $z \rightarrow \infty$  in the closed right half-plane.

In the second part of the paper, we consider the case  $\arg a_n = 2\pi in^2\alpha$  with *any irrational*  $\alpha$ . Theorem 3 shows that the qualitative picture of  $|f(z)|$  is the same as described by Littlewood, except that our estimate of the size of the pits is worse than exponential. In particular, we show that

$$\log |f(z)| = |z| + o(|z|),$$

outside some exceptional set of  $z$ . According to the Levin–Pfluger theory [15], this behavior of  $|f|$  has the following consequences about the zeros  $z_k$  of  $f$ :

The number  $n(r, \theta_1, \theta_2)$  of zeros (counting multiplicity) in the sector

$$\{z : \theta_1 < \arg z < \theta_2, |z| < r\}$$

satisfies

$$n(r, \theta_1, \theta_2) = \frac{\theta_2 - \theta_1}{2\pi}(r + o(r)) \quad \text{as } r \rightarrow \infty. \quad (5)$$

Moreover, the limit

$$\lim_{R \rightarrow \infty} \sum_{|z_k| \leq R} \frac{1}{z_k} \quad \text{exists,} \quad (6)$$

where  $z_k$  is the sequence of zeros of  $f$ . It is easy to see from the Taylor series of  $f$  that this limit equals  $-q$ .

Thus the Diophantine conditions used in [16, 28, 21] are unnecessary for the qualitative picture of behavior of  $|f|$ , but with arbitrary irrational  $\alpha$  the results are less precise than those where  $\alpha$  satisfies a Diophantine condition. Theorem 4 shows that Levy’s result (2) cannot be extended to arbitrary irrational  $\alpha$ . Finally we prove a result similar to Theorem 3 where the condition  $|a_n| = 1$  is replaced by a more flexible condition on the moduli of the coefficients allowing the function to have any order of growth.

We denote by

$$F(z) = \sum_{n=1}^{\infty} a_{n-1} z^{-n},$$

the Borel transform of  $f$  in (4) (terminology of [15]). Then  $F$  has an analytic continuation from a neighborhood of infinity to the region  $\overline{\mathbf{C}} \setminus K$ , where  $K$  is a convex compact set in the plane, which is called the conjugate indicator diagram. The indicator

$$h_f(\theta) := \limsup_{r \rightarrow \infty} r^{-1} \log |f(re^{i\theta})|, \quad |\theta| \leq \pi,$$

is the support function of the convex set symmetric to  $K$  with respect to the real axis.

We also consider the function

$$G(z) = \sum_{n=1}^{\infty} a_{n-1} z^n$$

analytic in the unit disc. Transition from  $F$  to  $G$  is by the change of the variable  $1/z$ .

**Pólya's theorem** ([15, Appendix I, §5]). *Suppose that  $G$  has an analytic continuation from the unit disc to infinity through some angle  $|\arg z - \pi| < \delta$ . Then the coefficients  $a_n$  can be interpolated by an entire function  $g$  of exponential type such that the indicator diagram of  $g$  is contained in the horizontal strip  $|\Im z| \leq \pi - \delta$ . That is  $g(n) = a_n$  for  $n \geq 1$ , and  $h_g(\pm\pi/2) \leq \pi - \delta$ .*

**Carlson's theorem** ([15, Ch. IV, Intro.]). *Suppose that the indicator diagram of an entire function  $g$  has width less than  $2\pi$  in the direction of the imaginary axis, that is  $h_g(\pi/2) + h_g(-\pi/2) < 2\pi$ . Then  $g$  cannot vanish on the positive integers, unless  $g = 0$ .*

**Theorem 1.** *Every entire function  $f$  of the form (4) has non-negative indicator, unless  $a_n = \text{const} \cdot a^n$  for some  $a$  on the unit circle, in which case  $f(z) = e^{az}$ .*

By Borel's transform, this is equivalent to

**Theorem 1'.** *Let  $G$  be as above. Then  $G$  cannot have an analytic continuation to infinity through any half-plane containing 0, unless  $a_n = \text{const} \cdot a^n$  for some  $a$ .*

These two theorems give characterizations of the exponential function and the geometric series, respectively, showing that their behavior is quite exceptional. Another somewhat similar characterization follows from the result in [23] mentioned above. As a corollary from Theorem 1 we obtain the following result of Carlson [4]: If  $z_n$  is the sequence of zeros of  $f$  as in (4), then

$$\sum_n 1/|z_n| = \infty, \quad (7)$$

unless  $f$  is an exponential.

*Proof of Theorem 1'.* Suppose that  $G$  has such an analytic continuation. Replacing  $z$  by  $az$  with  $|a| = 1$  we achieve that  $G$  has an analytic continuation to infinity through some left half-plane of the form  $\Re z < \epsilon$ , where  $\epsilon > 0$ .

Pólya's theorem then implies that  $a_n = g(n)$  for some entire function whose indicator diagram is contained in the strip  $|\Im z| < \pi/2 - \delta$ , for some  $\delta > 0$ . Consider the functions

$$g_R(z) = (g(z) + \overline{g(\bar{z})})/2 \quad \text{and} \quad g_I(z) = (g(z) - \overline{g(\bar{z})})/(2i).$$

On the real axis we have  $g_R(x) = \Re g(x)$  and  $g_I(x) = \Im g(x)$ . Consider the entire function

$$H = g_I^2 + g_R^2.$$

Then at positive integers we have

$$H(n) = g_I^2(n) + g_R^2(n) = (\Re g(n))^2 + (\Im g(n))^2 = |a_n|^2 = 1.$$

So the function  $H - 1$  has zeros at all positive integers. Its indicator diagram is contained in the strip

$$|\Im z| \leq \pi - 2\delta,$$

(Squaring stretches the indicator diagram by a factor of 2, and the indicator diagram of the sum of two functions is contained in the convex hull of the union of their diagrams). Now, by Carlson's theorem,  $H \equiv 1$ , so

$$g_I^2 + g_R^2 = 1. \quad (8)$$

The general solution of this functional equation in the class of entire functions is  $g_I = \cos \circ \phi$ ,  $g_R = \sin \circ \phi$ , where  $\phi$  is an entire function. It is well-known and easy to see that for  $g_I$  and  $g_R$  to be of exponential type, it is necessary and sufficient that  $\phi(z) = cz + b$ . As  $g_I$  and  $g_R$  are real on the real line,

we conclude that  $c$  and  $b$  are real. Thus  $a_n = \cos(cn + b) + i \sin(cn + b) = \text{const} \cdot e^{icn} = \text{const} \cdot a^n$ , as advertised.

An alternative way to derive the conclusion from (8) suggested by Katsnelson is to notice that (8) implies

$$|g(x)| \equiv 1 \quad \text{for real } x. \quad (9)$$

The Symmetry Principle then implies that  $g$  has no zeros (if  $z_0$  is a zero then  $\bar{z}_0$  would be a pole). So  $g$  is a function of exponential type without zeros, so  $g = \exp(icz)$ , where  $c$  should be real by (9).  $\square$

As we already noticed, Theorem 1 implies (7). However it does not imply that the sequence of zeros has positive density: there exist functions of exponential type, even with constant indicator, whose zeros have zero density.<sup>1</sup> To construct such examples, take zeros of the form

$$z_k = \left( e^{i \log \log(k+1)} - e^{i \log \log k} \right)^{-1},$$

and construct the canonical product  $W$  of genus one with such zeros. It is not hard to show that the asymptotic behavior of this product will be

$$\log |W(re^{i\theta})| = (cr + o(r)) \cos(\theta - \log \log r), \quad r \rightarrow \infty$$

outside of some small exceptional set, so the indicator  $h_W$  is constant, while the density of zeros is zero.

There exist entire functions of the form (4), other than the exponential, which are bounded in the left half-plane. The simplest example is Hardy's generalization of  $e^z$  defined by the power series

$$E_{s,a} = \sum_{n=1}^{\infty} \frac{(n+a)^s z^n}{n!}, \quad s \in \mathbf{C}, \quad a > 0.$$

For pure imaginary  $s$ , this series is of the form (4). Hardy [10] proved the asymptotic formula

$$E_{s,a}(z) = z^s e^z (1 + o(1)) + \frac{\Gamma(a)}{\Gamma(-s)(-z)^a \log(-z)} (1 + o(1)),$$

---

<sup>1</sup>Valiron [28, p. 415] erroneously asserted the contrary: that for functions with constant indicator, zero cannot be a Borel exceptional value.

as  $z \rightarrow \infty$ ,  $|\arg z \pm \pi/2| < \epsilon$ , for every  $\epsilon \in (0, \pi/2)$ . This formula implies that the functions  $E_{s,a}$  with pure imaginary  $s$  are bounded in the closed left half-plane. For further results on Hardy's function, see [22].

**Theorem 2.** *Let  $\psi$  be a real entire function with the property*

$$\psi(\zeta) = o(|\zeta|), \quad \zeta \rightarrow \infty$$

*in every half-plane  $\Re \zeta > c$ ,  $c \in \mathbf{R}$ . Then the function*

$$f(z) = \sum_{n=0}^{\infty} \frac{e^{i\psi(n)}}{n!} z^n$$

*is of the form (4) and for every  $A > 0$  and every  $\epsilon > 0$  we have*

$$|f(re^{i\phi})| = O(r^{-A}), \quad r \rightarrow \infty, \quad (10)$$

*uniformly for  $|\phi - \pi| \leq \pi/2 - \epsilon$ .*

*Proof.* We have the following integral representation:

$$f(-z) = -\frac{1}{2\pi i} \int_{-A-i\infty}^{-A+i\infty} \frac{\pi e^{i\psi(\zeta)} z^\zeta}{\Gamma(\zeta + 1) \sin \pi \zeta} d\zeta, = \frac{1}{2\pi i} \int_{-A-i\infty}^{-A+i\infty} e^{i\psi(\zeta)} z^\zeta \Gamma(-\zeta) d\zeta, \quad (11)$$

where  $A > 0$  is any positive number. To obtain this representation, we notice that that by Stirling's formula, the modulus of the integrand does not exceed

$$|z|^{-\Re \zeta} \exp((-\pi/2 + \phi + o(1))|\Im \zeta|),$$

as  $|\zeta| \rightarrow \infty$  in every half-plane of the form  $\Re \zeta \geq -A$ . Here  $o(1)$  is independent of  $z$ . Applying the residue formula to the rectangle

$$\{\zeta : -c < \Re \zeta < N + 1/2, |\Im \zeta| < N + 1/2\},$$

and letting  $N$  tend to infinity, we obtain (11). Now the same estimate of the integrand shows that (10) holds.  $\square$

Theorem 1 implies that the indicator diagram of a function of the form (4), other than an exponential, contains zero. Theorem 2 shows that the indicator diagram of such a function can be contained in a closed half-plane. It seems interesting to describe all possible indicator diagrams that can occur for functions of the form (4). We have the following partial result.

**Proposition.** *For arbitrary finite set  $Z$  on the unit circle, there exists an entire function of the form (4) whose indicator diagram coincides with the convex hull of  $Z \cup -Z$ .*

*Proof.* Let  $E$  be the set of all entire functions of the form (4). We consider the following operators on  $E$ :

$$R_\theta[f](z) := f(ze^{-i\theta}),$$

$$C[f](z) := \frac{1}{2}(f(z) + f(-z)),$$

and

$$S[f](z) := \frac{1}{2}(f(z) - f(-z)).$$

Now we define an operator  $E \times E \rightarrow E$  by the formula

$$Q_{\theta_1, \theta_2}[f_1, f_2] = (C \circ R_{\theta_1})[f_1] + (S \circ R_{\theta_2})[f_2].$$

It can be easily shown that if  $f \in E$  is a function with indicator diagram  $[0, 1]$ , then  $(C \circ R_\theta)[f]$  and  $(S \circ R_\theta)[f]$  have indicator diagram  $[-e^{i\theta}, e^{i\theta}]$ . Hence the indicator diagram of  $f_1 = Q_{\theta_1, \theta_2}[f, f]$  is the convex hull of

$$\{e^{i\theta_1}, -e^{i\theta_1}, e^{i\theta_2}, -e^{i\theta_2}\}.$$

This proves the Proposition for the sets  $Z$  of two points. Then we consider  $f_2 = Q_{0, \theta_3}[f_1, f]$  and so on.  $\square$

Now we consider functions of the form (4) with  $\arg a_n = 2\pi n^2\alpha$ ,  $\alpha \in \mathbf{R}$ .

**Theorem 3.** *Let  $f$  be of the form (4) with  $a_n = \exp(2\pi i n^2\alpha)$ , where  $\alpha$  is irrational. Then  $f$  has completely regular growth in the sense of Levin–Pfluger, and  $h_f \equiv 1$ .*

We recall the main facts of the Levin–Pfluger theory in the modern language [2]. Fix a positive number  $\rho$ . Let  $u$  be a subharmonic function in the plane satisfying

$$u(z) \leq O(r^\rho), \quad r \rightarrow \infty.$$

Then the family of subharmonic functions

$$A_t u(z) = t^{-\rho} u(tz), \quad t > 1,$$



is bounded from above on every compact subset of the plane. Such families of subharmonic functions are pre-compact in the topology  $D'$  of Schwartz's distributions [11, Theorem 4.1.9], so from every sequence  $A_{t_k}u$ ,  $t_k \rightarrow \infty$  one can select a convergent subsequence. An entire function  $f$  of order  $\rho$ , normal type is called of *completely regular growth* if the limit

$$u = \lim_{t \rightarrow \infty} A_t \log |f| \tag{12}$$

exists. It is easy to see that this limit is a fixed point for all operators  $A_t$ , so it has the form

$$u(re^{i\theta}) = r^\rho h(\theta),$$

and  $h$  is the indicator of  $f$ . Operators  $A_t$  also act on measures in the plane by the formula

$$A_t \mu(E) = t^{-\rho} \mu(tE) \quad \text{for } E \subset \mathbf{C}.$$

The Riesz measure  $\mu_f$  of  $\log |f|$  is the counting measure of zeros of  $f$ , and one of the results of Levin–Pfluger can be stated as follows: The existence of the limit (12) implies the existence of the limit

$$\mu = \lim_{t \rightarrow \infty} A_t \mu_f.$$

This limit  $\mu$  is also fixed by all operators  $A_t$ , so

$$d\mu = r^{\rho-1} dr d\nu(\theta),$$

where  $\nu$  is a measure on the unit circle which is called the *angular density* of zeros. This measure  $\nu$  is related to the indicator by the formula

$$d\nu = (h'' + \rho^2 h) d\theta,$$

in the sense of distributions.

Thus, as a corollary from Theorem 3, we obtain that the angular density of zeros of  $f$  is a constant multiple of the Lebesgue measure.

Completely regular growth with indicator 1 and order  $\rho = 1$  implies that

$$\log |f(re^{i\theta})| = r + o(r) \quad \text{as } r \rightarrow \infty, \tag{13}$$

uniformly with respect to  $\theta$ , when  $re^{i\theta}$  does not belong to an exceptional set. According to Azarin, [2], for every  $\eta > 0$ , this exceptional set can be covered by discs centered at  $w_k$  and of radii  $r_k$  such that

$$\sum_{|w_k| \leq r} r_k^\eta = o(r^\eta), \quad r \rightarrow \infty. \tag{14}$$

This improves the original condition with  $\eta = 1$  given in [15]. The properties (5) and (6) of zeros of  $f$ , stated in the beginning of the paper, follow from (13) by theorems II.2 and III.4 in [15], see also [24].

The exceptional set (14) is larger than the exceptional set in the work of Nassif. The exceptional set in Theorem 3 could be improved to a set of exponentially small circles if one knew that the zeros of  $f$  are well separated. This seems to be an interesting unsolved problem about the function (1). In particular, *can  $f$  of the form (1) have a multiple zero?* For  $\alpha = \sqrt{2}$ , Nassif proved that all but finitely many zeros are simple and well separated.

That the indicator of  $f$  in Theorem 3 is constant was proved by Valiron [28, p. 412]<sup>2</sup>. This also follows from the result of Cooper [6], who proved that the corresponding function  $G$  has the unit circle as its natural boundary, see also [5, p. 76, Footnote] where a short proof of Cooper's theorem is given. However, as we noticed above, constancy of the indicator by itself only implies (7); it is the statement about completely regular growth that permits to conclude that the zeros have positive density.

*Proof of Theorem 3.* By differentiating the power series it is easy to obtain

$$f'(z) = e^{2\pi i\alpha} f(ze^{i\beta}), \quad \text{where } \beta = 4\pi\alpha. \quad (15)$$

(This is the ‘‘pantograph equation’’ (3) with  $q = e^{2\pi i\alpha}$ .) The assumption that  $|a_n| = 1$  implies the following behavior of  $M(r, f)$

$$\log M(r, f) = r + o(r). \quad (16)$$

This is proved by the standard argument relating the growth of  $M(r, f)$  with the moduli of the coefficients, see, for example [15, Ch. I, §2]. The bounds  $0 \leq r - \log M(r, f) \leq (1/4 + o(1)) \log r$  can be obtained as follows. The upper bound  $M(r, f) \leq e^r$  is trivial, and for the lower bound, use Cauchy's inequality  $M(r, f) \geq r^n/n!$ , and maximize the right hand side with respect to  $n$ . In particular, the order  $\rho = 1$ .

It follows from (16) that the family of subharmonic functions

$$\{u_t = A_t \log |f| : 0 < t < \infty\}$$

---

<sup>2</sup>Valiron obtained the equation which is equivalent to our (20) below, [28, Eq. (11)] but he did not fully explore its consequences. Later in the same paper, on p. 421, Valiron proves that  $f$  is of completely regular growth only under an additional Diophantine condition on  $\alpha$ .

is uniformly bounded from above on compact subsets of  $\mathbf{C}$ . Moreover,  $u_t(0) = 0$ . So every sequence  $\sigma = (t_k) \rightarrow \infty$  contains a subsequence  $\sigma'$  such that the limit

$$u = \lim_{t \in \sigma', t \rightarrow \infty} u_t \quad (17)$$

exists in  $D'$ , the space of Schwartz's distributions in the plane. The set of all possible limits  $u$  for all sequences  $\sigma$  is called the limit set of  $f$  and denoted by  $\text{Fr}[f]$ . It consists of subharmonic functions in the plane satisfying  $u(0) = 0$ . Equation, (16) implies that

$$\max_{|z| \leq r} u(z) = r, \quad 0 \leq r < \infty. \quad (18)$$

If  $u = \lim t_k^{-1} \log |f(t_k z)|$ , and  $v = \lim t_k^{-1} \log |f'(t_k z)|$  with the same sequence  $t_k \rightarrow \infty$ , then

$$v \leq u. \quad (19)$$

Indeed, by Cauchy's inequality, for every  $\epsilon > 0$  and  $|z| > 1/\epsilon$ , we have

$$\log |f'(z)| \leq \max_{|\zeta| \leq \epsilon|z|} \log |f(z + \zeta)|.$$

This implies that for every  $\epsilon > 0$ ,

$$v(z) \leq \max_{|\zeta| \leq \epsilon} u(z + \zeta).$$

Now the upper semi-continuity of subharmonic functions shows that the right hand side of the last equation tends to  $u(z)$  as  $\epsilon \rightarrow 0$ , which proves (19).

The functional equation (15) and (19) imply that  $u(ze^{i\beta}) \leq u(z)$ , and this gives

$$u(ze^{i\beta}) \equiv u(z). \quad (20)$$

As  $\beta$  is irrational,  $u(z)$  is independent of  $\arg z$ , and taking (18) into account we conclude that the limit set  $\text{Fr}[f]$  consists of the single function  $u(z) = |z|$ . This means that  $f$  is of completely regular growth with constant indicator.  $\square$

Now we show that there exist irrational  $\alpha$  such that the corresponding functions  $f_\alpha$  in (1) do not have property (2).

**Theorem 4.** *There is a residual set  $E$  on the unit circle, such that for a function  $f_\alpha$  as in (1) with  $\alpha \in E$ , we have*

$$\limsup_{r \rightarrow \infty} M(r, f)/m_2(r, f) = \infty. \quad (21)$$

We recall that a set is called residual if it is an intersection of countably many dense open sets. By Baire's Category Theorem, residual sets on  $[0, 1]$  have the power of a continuum and thus contain irrational points.

*Proof of Theorem 4.* Consider the sets

$$E_{m,n} = \{\alpha : M(r, f_\alpha)/m_2(r, f_\alpha) \leq m \text{ for } r \geq n\},$$

where  $m$  and  $n$  are positive integers. Evidently, all these sets are closed. Let  $E = [0, 1] \setminus \cup_{m,n} E_{m,n}$ . Then for  $\alpha \in E$  we have (21), and  $E$  is a countable intersection of open sets. It remains to show that  $E$  is dense. We will show that  $E$  contains all rational numbers. Indeed, for rational  $\alpha$ ,  $f_\alpha$  is a finite trigonometric sum:

$$f_\alpha = \sum c_k e^{b_k z}, \quad (22)$$

where  $b_k$  are roots of unity. This representation immediately follows from the functional equation (15): iterating this functional equation finitely many times, we obtain a linear differential equation whose solutions are trigonometric sums (22). It is clear that any finite trigonometric sum  $g$  satisfies

$$M(r, g)/m_2(r, g) \rightarrow \infty \text{ as } r \rightarrow \infty.$$

This proves that  $E$  is dense and thus residual.  $\square$

Now we extend Theorem 3 to the case that  $|a_n|$  is non-constant. For this we need

**Hadamard's Multiplication Theorem** [3]. *Let  $f = \sum_{n=0}^{\infty} c_n z^n$  be an entire function, and  $H = \sum_{n=0}^{\infty} b_n z^n$  a function analytic in  $\overline{\mathbf{C}} \setminus \{1\}$ . Then the function*

$$(f \star H)(z) = \sum_{n=0}^{\infty} a_n c_n z^n$$

*has the integral representation*

$$(f \star H)(z) = \frac{1}{2\pi i} \int_C f(\zeta) H\left(\frac{z}{\zeta}\right) \frac{d\zeta}{\zeta},$$

*where  $C$  is any closed contour going once in positive direction around the point 1.*

This operation  $f \star H$  is called the Hadamard composition of power series. From the integral representation we immediately obtain

$$|(f \star H)(z)| \leq K \max_{|\zeta-z| \leq r|z|} |f(\zeta)|, \quad \text{where } K = \max_{|\zeta-1|=r/(1-r)} |H(\zeta)|. \quad (23)$$

**Theorem 5.** *Let  $h$  be an entire function of minimal exponential type. Let*

$$c_n = h(0)h(1) \dots h(n), \quad n \geq 0, \quad (24)$$

*and assume in addition that*

$$-\log |c_n| = \frac{1}{\rho} n \log n - cn + o(n), \quad n \rightarrow \infty, \quad (25)$$

*with some real constant  $c$ . Then the entire function*

$$f(z) = \sum_{n=0}^{\infty} c_n e^{2\pi i n^2 \alpha} z^n,$$

*with irrational  $\alpha$ , has order  $\rho$ , normal type and completely regular growth with constant indicator.*

The condition that  $c_n/c_{n-1}$  is interpolated by an entire function of minimal exponential type is not so rigid as it may seem. In this connection, we recall a theorem of Keldysh (see, for example, [9]) that every function  $h_1$  analytic in the sector  $|\arg z| < \pi - \epsilon$  and satisfying  $\log |h_1(z)| = O(|z|^\lambda)$ ,  $z \rightarrow \infty$  there, with  $\lambda = \pi/(\pi + \epsilon) < 1$ , can be approximated by an entire function  $h$  of normal type, order  $\lambda$  so that

$$|h(z) - h_1(z)| = O(e^{-|z|^\lambda}), \quad z \rightarrow \infty, \quad |\arg z| < \pi - 2\epsilon.$$

For example, one can take

$$h_1(z) = z^{-1/\rho}, \quad \rho > 0,$$

and apply Keldysh's theorem, to obtain a function  $f$  of normal type, order  $\rho$ , satisfying all conditions of Theorem 5.

*Proof of Theorem 5.* We write:

$$f(z) = \sum_{n=0}^{\infty} c_n e^{2\pi i n^2 \alpha} z^n$$

$$\begin{aligned}
&= 1 + \sum_{n=0}^{\infty} c_{n+1} e^{2\pi i(n+1)^2 \alpha} z^{n+1} \\
&= 1 + z e^{2\pi i \alpha} \sum_{n=0}^{\infty} c_{n+1} e^{2\pi i n^2 \alpha} (z e^{4\pi i n \alpha})^n \\
&= 1 + z e^{2\pi i \alpha} (f \star H)(z e^{i\beta}), \quad \text{where } \beta = 4\pi \alpha,
\end{aligned}$$

and

$$H(z) = \sum_{n=0}^{\infty} \frac{c_{n+1}}{c_n} z^n.$$

By the assumption (24) of the theorem, and Pólya's theorem above,  $H$  is holomorphic in  $\overline{\mathbf{C}} \setminus \{1\}$ , so the estimate (23) holds. Assumption (25) implies that

$$M(r, f) = \sigma r^\rho + o(r^\rho), \quad r \rightarrow \infty,$$

where  $\sigma = e^{c\rho}/(e\rho)$ , see [15, Ch. I, §2]. So from every sequence one can select a subsequence such that the limits

$$u(z) = \lim_{k \rightarrow \infty} t_k^{-\rho} \log |f(t_k z)| \quad \text{and} \quad v(z) = \lim_{k \rightarrow \infty} t_k^{-\rho} \log |(f \star H)(t_k z)|$$

exist, and (23) implies that  $v \leq u$ , by a similar argument as (19) was derived. Now the equation

$$f(z) = 1 + z e^{2\pi i \alpha} (f \star H)(z e^{i\beta}) \tag{26}$$

implies

$$u(z) \leq \max\{0, v(z e^{i\beta})\} \leq \max\{0, u(z e^{i\beta})\},$$

and as  $\beta$  is irrational, we conclude that  $u$  does not depend on  $\arg z$ . This completes the proof.  $\square$

The authors thank Alan Sokal whose questions stimulated this research, and Victor Katsnelson and Vitaly Bergelson for their help with literature.

## References

- [1] M. Ålander, Sur le déplacement des zéros des fonctions entières par leur dérivation, Thèse pour le doctorat, Upsal, 1914.
- [2] V. S. Azarin, Asymptotic behavior of subharmonic functions of finite order. Mat. Sbornik 108 (1979) no. 2, 147–167.

- [3] L. Bieberbach, *Analytische Fortsetzung*, Springer, Berlin, 1955.
- [4] F. Carlson, Sur une fonction entière, *Arkiv för Mat., Astron. och Fys.*, 10, (1915) Nr. 16, 1–5.
- [5] F. W. Carroll, On some classes of noncontinuable analytic functions, *Trans. AMS* 94 (1960) 74–85.
- [6] R. Cooper, The behaviour of certain series associated with the limiting cases of elliptic theta functions, *Proc. London Math. Soc.* 27 (1928) 410–426.
- [7] M. A. Evgrafov, *The Abel–Goncharov interpolation problem*, GITTL, Moscow 1954 (Russian).
- [8] M. A. Evgrafov, On completeness of systems of analytic functions close to  $\{z^n P(z)\}$ ,  $\{[\varphi(z)]^n\}$ , and on some interpolation problems. (Russian) *Izvestiya Akad. Nauk SSSR. Ser. Mat.* 17, (1953) 421–460.
- [9] D. Gaier, *Vorlesungen über Approximation im Komplexen*, Birkhauser, Basel, 1980.
- [10] G. H. Hardy, On the zeros of certain class of integral Taylor series II, *Proc. LMS* 2 (1905) 401–431.
- [11] L. Hörmander, *Analysis of partial differential operators*, vol. 1, Springer 1990.
- [12] L. Hörmander, *Notions of convexity*, Birkhäuser, Boston, MA, 1994.
- [13] A. Iserles, On the generalized pantograph functional-differential equation, *Euro. J. Appl. Math.*, 4 (1993) 1–38.
- [14] J. Langley, A certain functional-differential equation, *J. Math. Anal. Appl.*, 244 (2000) 564–567.
- [15] B. Ya. Levin, *Distribution of zeros of entire functions*, AMS, Providence RI, 1980.
- [16] P. Levy, Sur la croissance des fonctions entières, *Bull. Soc. Math. France*, 58 (1930) 29–59 et 127–149.

- [17] J. E. Littlewood, A “pits effect” for all smooth enough integral functions with a coefficient factor  $\exp(n^2\alpha\pi i)$ ,  $\alpha = \frac{1}{2}(\sqrt{5} - 1)$ , J. London Math. Soc. 43 (1968) 79–92.
- [18] J. E. Littlewood, The “pits effect” for the integral function  $f(z) = \sum \exp\{-\rho^{-1}(n \log n - n) + \pi i \alpha n^2\} z^n$ ,  $\alpha = \frac{1}{2}(\sqrt{5} - 1)$ , Number Theory and Analysis (Papers in Honor of Edmund Landau), 193–215, Plenum, NY 1969.
- [19] J. E. Littlewood and A. C. Offord, On the distribution of zeros and  $a$ -values of a random integral function. II, Ann. of Math. 49, (1948) 885–952; errata 50, (1949) 990–991.
- [20] S. Macintyre, An upper bound for the Whittaker constant, Duke Math. J., 15 (1948) 953–954.
- [21] M. Nassif, On the behaviour of the function  $f(z) = \sum_{n=0}^{\infty} e^{\sqrt{2}\pi i n^2} (z^{2n}/n!)$ , Proc. London Math. Soc. 54 (1952) 201–214.
- [22] I. Ostrovskii, Hardy’s Generalization of  $e^z$  and Related Analogs of Cosine and Sine Computational Methods and Function Theory, 6 (2006) 1–14.
- [23] L. Rubel and K. Stolarsky, Subseries of the power series for  $e^x$ , Amer. Math. Monthly, 87 (1980) 371–376.
- [24] M. Sodin, A remark on the limit sets of subharmonic functions of integer order in the plane, (Russian) Teor. Funktsii Funktsional. Anal. i Prilozh. No 40 (1983) 125–129.
- [25] A. Sokal, Private communication.
- [26] S. Tims, Note on a paper by M. Nassif, Proc. London Math. Soc. (2) 54 (1952) 215–218.
- [27] W. T. Tutte, On dichromatic polynomials, J. Combin. Theory, 2 (1967) 301–320.
- [28] G. Valiron, Sur une équation fonctionnelle et certaines suites de facteurs, J. de Math. pures appl., 12 (1945) 405–423.



*Purdue University, West Lafayette IN 47907-2067*  
*eremenko@math.purdue.edu*

*Bilkent University, Ankara, Turkey*