

Two-parametric PT-symmetric quartic family

Alexandre Eremenko and Andrei Gabrielov*

May 6, 2011

Abstract

We describe a parametrization of the real spectral locus of the two-parametric family of PT-symmetric quartic oscillators. For this family, we find a parameter region where all eigenvalues are real, extending the results of Dorey, Dunning, Tateo and Shin.

MSC: 81Q05, 34M60, 34A05

Key words: one-dimensional Schrödinger operators, PT-symmetry, quasi-exact solvability, Darboux transform.

1 A family of quartic oscillators

We consider the eigenvalue problem in the complex plane

$$w'' + (\zeta^4 + 2b\zeta^2 + 2iJ\zeta + \lambda)w = 0, \quad w(te^{-\pi i/2 \pm \pi i/3}) \rightarrow 0, \quad t \rightarrow +\infty. \quad (1)$$

Here J and b are parameters. This two-parametric family is interesting for several reasons.

When $2J$ is an integer, $2J < 1$, and $b \geq 0$, the problem has the same spectrum as a spherically symmetric quartic oscillator in \mathbf{R}^d . In this case $2J = 2 - 2l - d$, where l is the angular momentum quantum number [5, 4].

When J is a positive integer, problem (1) is quasi-exactly solvable (QES) [2]. This means that there are J eigenfunctions of the form

$$w(\zeta) = p(\zeta) \exp(-i\zeta^3/3 - ib\zeta),$$

*Both authors are supported by NSF grant DMS-1067886.

where p is a polynomial of degree $J - 1$ in ζ whose coefficients are algebraic functions in b .

When J and b are real, the problem is PT-symmetric. The eigenvalues of a PT-symmetric problem can be either real or come in complex conjugate pairs. Both possibilities can be present for $J > 1$. A very interesting feature is level crossing in the real domain: for some real b and J the graphs of the eigenvalues $\lambda_k(b)$ can be real and cross each other. This phenomenon was discovered by Bender and Boettcher [2] numerically, then it was studied in [12], where the presence of infinitely many such real level crossing points was proved for odd positive J .

When $J \rightarrow +\infty$, the QES part of the spectral locus approximates the whole spectral locus of the PT-symmetric cubic family

$$-w'' + (iz^3 + iaz)w = \lambda w, \quad w(\pm\infty) = 0, \quad (2)$$

which was subject of intensive research, see, for example, [3, 7, 8, 11, 16, 11, 20, 23].

By the change of the independent variable $z = i\zeta$ problem (1) is equivalent to

$$L_{b,J}(y) = -y'' + (z^4 - 2bz^2 + 2Jz)y = \lambda y, \quad y(te^{\pm\pi i/3}) \rightarrow 0, \quad t \rightarrow +\infty. \quad (3)$$

Shin's theorem [20] applies to these eigenvalue problems when $J \leq 0$, and implies that for $J \leq 0$ all eigenvalues are real. The proof of Shin's theorem is based on the remarkable ODE-IM correspondence of Dorey, Dunning and Tateo [8]. Here we extend this result.

Theorem 1. *All eigenvalues of (1) or (3) are real for $J \leq 1$.*

The condition $J \leq 1$ is exact, because it is known that for every $J > 1$ there are non-real eigenvalues [12]. Our proof of Theorem 1 is based on purely topological arguments. Using the formulation (3), we establish certain property of eigenfunctions for $J = 0$, and then show that this property persists for $J < 1$ and prevents level crossing.

The *real spectral locus* $Z(\mathbf{R}) \subset \mathbf{R}^3$ is defined as the set of all real triples (b, J, λ) for which there exists $y \neq 0$ satisfying (3). This is an analytic surface in \mathbf{R}^3 . The main result of this paper is a parametrization of $Z(\mathbf{R})$ in terms of Nevanlinna parameters (sections ?-?). In [11] we obtained similar parametrization of the real spectral locus of (2) and another two-parametric family of quartics.

The family considered in this paper is much more complicated because of the presence of QES part and of the real level crossings. In [13] we parametrized the real *quasi-exactly solvable locus* $Z^{QES}(\mathbf{R})$ of (3) which consists of all triples $(b, J, \lambda) \in \mathbf{R}^3$ for which there exists a function $y(z) = p(z) \exp(z^3/3 - bz)$ satisfying (3) with a polynomial p .

The plan of the paper is the following. In the next section we introduce Nevanlinna parameters and state our principal result, Theorem 2, about the correspondence between parameters $(b, J, \lambda) \in Z(\mathbf{R})$ and Nevanlinna parameters. In section 3 we prove the “easy”, algebraic part of this correspondence. In section 4 we study the case $J = 0$ and in section 6 we prove Theorem 1. Then in section 7 we describe the parametrization of the part of the real spectral locus $Z(\mathbf{R})$ where J is an integer in terms of Nevanlinna parameters and complete the proof of Theorem 2. In the last two sections we parametrize the whole real spectral locus.

2 Nevanlinna parameters

The references for this section are [18, 21, 9, 10, 11].

Suppose that b and J are real. Let y be an eigenfunction of (3). Then $y^*(z) = \overline{y(\bar{z})}$ satisfies the same differential equation and the same boundary conditions (3), so $y^* = cy$. We can choose y so that $y(x_0) \in \mathbf{R} \setminus \{0\}$ for some real x_0 . Substituting this x_0 to $y^* = cy$ we conclude that $c = 1$. So our eigenfunction y is real. It is defined up to multiplication by a real constant.

Let y_1 be a solution of the differential equation in (3), which is real and linearly independent of y . Consider the meromorphic function $f = y/y_1$. It is a *Nevanlinna function*, which means that f has no critical points in \mathbf{C} and the only singularities of f^{-1} are finitely many logarithmic branch points.

In the case that y_1 is normalized by $y_1(x) \rightarrow 0$, $x \rightarrow +\infty$, $x \in \mathbf{R}$, we call f a *normalized Nevanlinna function* of (3). The normalized Nevanlinna function is defined up to a real multiple. Existence of y_1 with such normalization is guaranteed by a theorem of Sibuya [21].

Different Nevanlinna functions associated with the same point $(b, J, \lambda) \in Z(\mathbf{R})$ are related by a real fractional-linear transformation of the form $f \mapsto \alpha f / (f - \beta)$ where $\alpha \neq 0$ and β are real.

Nevanlinna functions f of (3) have no critical points, because $f' = (y'y_1 - yy_1')/y_1^2$ and $y'y_1 - yy_1' = \text{const}$. They have 6 asymptotic values in the sectors

$$S_j = \{te^{i\theta} : t > 0, |\theta - \pi j/3| < \pi/6\}, \quad j = 0, \dots, 5.$$

In what follows j is understood as a residue modulo 6. These sectors are in one-to-one correspondence with logarithmic branch points of f^{-1} . The asymptotic values of the normalized Nevanlinna function are 0 in S_1 and S_{-1} , because of the boundary condition, and ∞ in S_0 because of the normalization of y_1 . We denote by c and a the asymptotic values of the normalized Nevanlinna function in S_2 and S_3 , respectively. As f is real, the asymptotic value in S_{-2} is \bar{c} , and a is real.

These asymptotic values a, c are called the *Nevanlinna parameters*. They are related to the Stokes multipliers by simple formulas [21, 17]. Normalized function f and the Nevanlinna parameters are defined modulo multiplication by a real non-zero number, so we can further normalize them. We will use different normalizations, depending of the situation.

Nevanlinna parameters $(c, a) \in \overline{\mathbf{C}} \times (\mathbf{R} \cup \{\infty\})$ are subject to the following conditions:

$$c \neq 0, \quad c \neq a, \tag{4}$$

and the set $\{0, \infty, a, c, \bar{c}\}$ contains at least 3 distinct points, in other words, the combination $c = \infty, a = 0$ is prohibited. [18, 21].

Nevanlinna parameters (modulo multiplication by real constants) serve as local coordinates on the real spectral locus $Z(\mathbf{R})$, [1, 17]. Notice that the set of pairs satisfying (4) modulo proportionality is a non-Hausdorff manifold. But this will cause no difficulties as we always work in local charts.

Relation between (b, J, λ) and (a, c) is very complicated: Nevanlinna's construction of the map $(a, c) \rightarrow (b, J, \lambda)$ involves the uniformization theorem. So it is interesting and challenging to establish any explicit correspondences between sets in the space of Nevanlinna parameters and sets in the (b, J, λ) space.

Our main result in this direction is

Theorem 2. *J is an integer if and only if*

$$a = 0 \quad \text{or} \quad c = \bar{c}.$$

In the next section we prove the “only if” part.

3 QES locus and Darboux transform

In this section we prove the “easy part” of Theorem 2: is J is an integer, then either $a = 0$ or $c = \bar{c}$.

Suppose that for some $(b, J, \lambda) \in Z(\mathbf{R})$ we have $a = 0$. This means that the eigenfunction y tends to zero in S_3 (and also in S_1, S_{-1}), so y is an elementary function of the form pe^q with polynomials p and q . Substitution to (3) gives

$$y(z) = p(z) \exp(z^3/3 - bz),$$

where p is a polynomial. It is known that such eigenfunctions exist if and only if J is a positive integer [2, 12]. Such points (b, J, λ) form the QES spectral locus $Z^{QES}(\mathbf{R})$ which consists of smooth algebraic curves

$$Q_J(b, \lambda) = 0, \quad J = n + 1, \quad n \geq 0,$$

where Q_J are real polynomials of degree J in λ . QES spectral locus was studied in [12, 13]; in the second paper it was parametrized in terms of Nevanlinna parameters. The converse is evident: if $(b, J, \lambda) \in Z^{QES}(\mathbf{R})$, then $a = 0$.

In [12] we obtained the following results:

(i) *If J is a positive integer, and b is real, then all non-QES eigenvalues are real.*

Let $Z_J(\mathbf{R}) = \{(b, \lambda) \in \mathbf{R}^2 : (b, J, \lambda) \in Z(\mathbf{R})\}$ and let $Z_J^{QES}(\mathbf{R})$ be similarly defined. Let Z_J^* be the closure of $Z_J(\mathbf{R}) \setminus Z_J^{QES}(\mathbf{R})$ in \mathbf{R}^2 .

(ii) *When J is even, $Z_J^*(\mathbf{R}) \cap Z_J^{QES}(\mathbf{R}) = \emptyset$. When J is odd, then $c = \bar{c}$ holds at all points (b, λ) of this intersection.*

(iii) *If J is a positive integer, then $Z_J^* = Z_{-J}$.*

To deal with the condition $c = \bar{c}$, we use the Darboux transform [6, 14, 12], which we recall. Let ψ_0, \dots, ψ_n be some eigenfunctions of a differential operator $L = -d^2/dz^2 + V(z)$ with eigenvalues $\lambda_0, \dots, \lambda_n$. Then the differential operator

$$-\frac{d^2}{dz^2} + V - 2\frac{d^2}{dz^2} \log W(\psi_0, \dots, \psi_n), \quad (5)$$

where W is the Wronski determinant, has the same eigenvalues as L , *except* $\lambda_0, \dots, \lambda_n$.

Let $J \geq 0$ be an integer, $(b, \lambda) \in Z_J^*(\mathbf{R})$, and let y be the eigenfunction corresponding to (b, J, λ) . We apply the Darboux transform to our operator $L_{b,J}$ in (3), taking all QES eigenfunctions as ψ_0, \dots, ψ_n . If $J = 0$ the Darboux transform does not change anything. It is easy to see, [12] that the

transformed operator (5) is $L_{b,-J}$ in this case. But $L_{b,J}(y) = \lambda y$, and if we define $y^*(z) = y(-z)$, then $L_{b,-J}(y^*) = \lambda y^*$. However, this y^* is not an eigenfunction of $L_{b,-J}$, because it does not satisfy the normalization condition in (3). Instead it tends to zero in S_2 and in S_{-2} . This means that y^* is linearly independent of the eigenfunction y_0 of $L_{b,-J}$, and $g = y_0/y^*$ has asymptotic values ∞ in S_2 and in S_{-2} .

Since g has equal asymptotic values in S_2 and S_{-2} any other Nevanlinna function for the same (b, J, λ) has the same property. So $c = \bar{c}$ at the point $(b, -J, \lambda)$.

The case of negative J is treated similarly, applying the inverse Darboux transform. Thus $c = \bar{c}$ on $Z_J^*(\mathbf{R})$ when J is an integer. This proves the “only if” part of Theorem 2.

As a byproduct we obtain the following

(iv) If $J > 0$ is odd, then the condition $(b, \lambda) \in Z_J^{QES}(\mathbf{R}) \cap Z_J^(\mathbf{R})$ is equivalent to the condition $c = \infty$ at the point $(b, \lambda) \in Z_{-J}(\mathbf{R})$.*

Indeed, the eigenfunction y is elementary if and only if the asymptotic values of f in S_3, S_1, S_{-1} are zero. This happens if and only if the function g defined above has asymptotic values ∞ in S_0, S_2, S_{-2} .

To prove the second part of Theorem 2, we need to find which part of the spectral locus corresponds to real c .

4 Line complexes

Assigning asymptotic values in sectors S_j is not enough to define a Nevanlinna function f , one needs additional information about the topology of the covering

$$f : \mathbf{C} \setminus f^{-1}(\text{asymptotic values}) \rightarrow \overline{\mathbf{C}} \setminus \{\text{asymptotic values}\}.$$

Such information is encoded in the following way [18, 9, 10]. One chooses a cell decomposition Φ of the Riemann sphere $\overline{\mathbf{C}}$ such that each 2-cell contains one asymptotic value, and takes the f -preimage of this cell decomposition. This preimage is a cell decomposition Ψ of the plane which locally looks like Φ . There are many ways to choose Φ , and here we describe cell decompositions, used by Nevanlinna, see also [15, Ch. 7] for a comprehensive treatment.

First we fix two points in $\overline{\mathbf{C}}$ which are distinct from the asymptotic values. We call these points \times and \circ . They are the vertices of Φ . Suppose that we have q asymptotic values. We connect \times and \circ with q edges which do not intersect except at the ends. These edges and vertices form the 1-skeleton of Φ ; we require that faces of Φ contain one asymptotic value each. This choice of Φ is fixed in this section.

The preimage $\Psi = f^{-1}(\Phi)$ is called the *line complex*. It has the following property: *the 1-skeleton of Ψ is a bipartite connected graph embedded in \mathbf{C} whose all vertices have degree q* . This property completely characterizes all possible line complexes. We label the faces of Φ and Ψ with the asymptotic values, so that a face of Ψ has the same label as its image. Two cell decompositions Ψ' and Ψ'' are considered equivalent if one can be mapped onto another by a homeomorphism of the plane preserving orientation and labels of faces and vertices. If two Nevanlinna functions f and g have equivalent cell decompositions, then $f(z) = g(\alpha z + \beta)$, $\alpha \neq 0$.

The following properties are evident. The cyclic order of face labels around a vertex of Ψ is the same as for the image vertex in Φ . Two faces of Ψ with the same labels have disjoint closures. Each face is bounded either by two edges or by infinitely many edges.

By erasing the multiple edges of the 1-skeleton of Ψ and discarding the labels of bounded faces, we obtain a new cell decomposition with labeled faces and vertices, whose 1-skeleton is a tree. The line complex can be uniquely recovered from its associated tree.

If f is real, so asymptotic values are symmetric with respect to complex conjugation, sometimes it is possible to choose a symmetric cell decomposition Φ . Then Ψ is also symmetric. Then f can be made real by a precomposition with an affine map of \mathbf{C} .

Line complexes are convenient for study of the limits of families of Nevanlinna functions when two asymptotic values collide. We have the following compactness theorem [22]. Fix a cell decomposition Φ . Let f_n be a sequence of Nevanlinna functions with line complexes $\Psi_n = f_n^{-1}(\Phi)$. Suppose that $v_n = 0$ is a vertex of Ψ_n , and that f_n are normalized by conditions $|f'_n(0)| = 1$. Then one can choose a subsequence from f_n that tends to a limit, and this limit is a Nevanlinna function.

We need only the special case when the cell decompositions Ψ_n corresponding to f_n are all equivalent. If distinct asymptotic values of f_n tend to distinct limits, then f has the same cell decomposition Ψ . If two asymptotic values of f_n which are labeling adjacent faces of Ψ , collide in the limit, then

one has to erase from the 1-skeleton of Φ all edges on the common boundaries of faces with these collided asymptotic values. The component of the remaining graph containing the vertex v is the cell decomposition of the limit function.

Now we return to our Nevanlinna functions corresponding to problem (3). One technical problem we are facing is that it is not always possible to choose a symmetric Φ . However this is possible when c is real, and in the next sections we consider this case. We begin with the simplest case when $J = 0$. In this case the Nevanlinna function has additional symmetry.

The cell decompositions considered in [13] are *different* from line complexes, because in the situation considered in that paper, it is impossible to define a line complex with the required symmetry properties.

5 Subfamily $J = 0$

In this section we begin to prove Theorem 1. For $J = 0$, we know from the results of [4] and [20] that all eigenvalues are real.

Let y be an eigenfunction of $L_{b,0}$. Function $y_1(z) = y(-z)$ satisfies the differential equation in (3) with $J = 0$, but does not satisfy the boundary conditions. So y_1 is linearly independent of y , and we consider the Nevanlinna function

$$f = y/y_1. \tag{6}$$

It is *not normalized* in the sense of section 2. This function f has the following symmetry properties:

$$f^* = f, \quad f(-z) = 1/f(z).$$

The asymptotic values are 0 in S_1 and S_{-1} , ∞ in S_2 and S_{-2} , $A \in \mathbf{R}$ in S_0 and $1/A$ in S_3 . This A is the Nevanlinna parameter. As f is defined up to multiplication by a real non-zero number, we can normalize so that $A > 0$.

Proposition 1. $A \in (0, 1)$. *Each eigenfunction has at most one zero on the real line.*

Proof. Function f has the property that

$$f : \mathbf{C} \setminus f^{-1}(0, A, 1/A, \infty) \rightarrow \overline{\mathbf{C}} \setminus \{0, A, 1/A, \infty\}$$

is a covering. To construct the line complex, we choose the cell decomposition Φ of the target sphere shown in Fig. 1.




Fig. 1.

It has two vertices, four edges and four faces labeled by the asymptotic values. We denote by A' the asymptotic value which is on $(0, 1)$, so that A' is either A or A^{-1} , and our first goal is to find out which of these possibilities holds.

The line complex $\Psi = f^{-1}(\Phi)$ is a labeled cell decomposition of \mathbf{C} . It has 6 unbounded faces. Moreover, Ψ is symmetric with respect to the real line and with respect to the imaginary line. The symmetry with respect to the real line does not change the labels, while the symmetry with respect to the imaginary line interchanges \times with \circ , 0 with ∞ and A' with $1/A'$. Unbounded faces of Ψ are asymptotic to the sectors S_j .

For any two pairs of vertices of Ψ connected by several edges, we replace these several edges with one edge. The result is a simpler cell decomposition T whose 1-skeleton is a tree. The faces of T are labeled with asymptotic values, and T has all symmetry properties described above. The label of a face asymptotic to S_j is the asymptotic value in S_j .

It is easy to classify all possible labeled trees satisfying the above conditions. They all have two vertices of order 4 and the number k of edges between these two vertices is odd. These trees depend on one non-negative integer parameter m , such that $k = 2m + 1$, and we denote them by T_m . The tree T_1 and the corresponding line complex are shown in Fig. 2.




Fig. 2. The tree T_1 (left) and the corresponding line complex (right).

Comparing the cyclic order of labels of faces adjacent to a vertex of order 4 in Fig. 1 and Fig. 2 we conclude that $A = A'$ so $A \in (0, 1)$. This proves the first part of the proposition. The second part is immediately clear from the classification of the trees T_m : the cell decomposition Ψ has at most one face labeled with 0 which intersects the real line. (A similar argument was used in [9].) Notice that

$$\text{the number of real zeros of } y \text{ is } 0 \text{ if } m \text{ is even and } 1 \text{ if } m \text{ is odd.} \quad (7)$$

Function f depends continuously on the parameter b . It follows that A also depends continuously on b . As all eigenvalues are real, for every $b \in \mathbf{R}$, they cannot collide as b changes, and we can label them in increasing order, $\lambda_0(b) < \lambda_1(b) < \dots$. To each eigenvalue λ_k corresponds one cell decomposition Ψ , and one tree $T = T_m$. So we have some correspondence $m(k)$, which is defined so that $T_{m(k)}$ corresponds to λ_k .

Proposition 2. $m(k) = k$. So the eigenfunction y_k corresponding to λ_k has no real zeros when k is even and one real zero when k is odd.

Proof. We use the asymptotic result in [12, 11] to degenerate (3) as $b \rightarrow +\infty$ to the harmonic oscillator

$$Y''(u) + 4u^2Y(u) = \mu Y(u), \quad Y(it) \rightarrow 0, \quad t \rightarrow \pm\infty.$$

The boundary condition here comes from the boundary condition in (3). Eigenfunctions Y_k corresponding to eigenvalues μ_k , labeled in the increasing order, have all zeros on the imaginary axis. There is one zero on the real axis (namely at the origin) when k is odd and none if k is even.

To analyse the behavior of zeros of eigenfunctions of (3) as $b \rightarrow +\infty$, we consider the function $g = f/(f - A)$. This corresponds to a different choice of y_1 in the basis (y, y_1) of solutions of the differential equation in (3). Asymptotic values of g are ∞ in S_0 , 0 in $S_{\pm 1}$, 1 in $S_{\pm 2}$, and $a := 1/(1 - A^2)$ in S_3 .

According to the result of [11], to every eigenfunction Y_k with eigenvalue μ_k , and to every positive b large enough corresponds a unique eigenfunction y of (3) with eigenvalue $\lambda(b) \rightarrow \mu_k$. Let $m = m(k)$ and let T_m be the tree corresponding to y . Then T_m must collapse to the tree T'_k corresponding to Y_k (see Fig. 3).




Fig. 3. Tree T'_1 .

By collapse of a tree we mean the following: two face labels become equal, and all edges on the common boundary of these faces are erased. Such collapse can happen only if $1/(1 - A^2) \rightarrow 1$ or $1/(1 - A^2) \rightarrow \infty$. In the second case we must multiply g by $1 - A^2$, so the asymptotic value in S_3 becomes 1 and in $S_{\pm 2}$ it becomes $1 - A^2 \rightarrow 0$. Then as the tree $T_m(k)$ collapses to the tree T'_k , we must have $m(k) = k$. This proves the proposition.

Proposition 2 gives a parametrization of the spectral locus for $J = 0$ by Nevanlinna parameter A . For each m , the correspondence $b \rightarrow A$ is a real analytic homeomorphism of the real line onto $(0, 1)$. Later we will prove that this correspondence is increasing.

6 Region $J < 1$

Now we can prove Theorem 1. When b, J and λ are real, the eigenvalue problem is PT-symmetric, and we can define a real eigenfunction y . We choose the second linearly independent solution y_1 of the differential equation in (3) from the conditions that $y_1 \rightarrow 0$ in S_0 , and y_1 is real. Then $f = y/y_1$ is real. The asymptotic values of f in S_0, \dots, S_5 are $\infty, 0, c, a, \bar{c}, 0$ in this order, where $a \in \mathbf{R}$. Thus for every (b, J, λ) on the real spectral locus, Nevanlinna parameters $a \in \overline{\mathbf{R}}$ and $c \in \overline{\mathbf{C}}$ are defined.

As long as $a \neq 0$, the number of real zeros cannot change, because y and f never have multiple zeros.

Equality $a = 0$ can happen on the spectral locus if and only if J is a positive integer and the eigenfunction y is elementary [2, 12].

Lemma 1. *Let $\gamma(t), t \in [0, 1]$ be a curve in the real (b, J) plane. Suppose that $\gamma(0) = (0, 0)$, and that $J \leq 1$ on γ . Then for $(b, J) = \gamma(1)$, all eigenvalues are real.*

This implies Theorem 1.

Proof of Lemma 1. It is sufficient to prove the lemma for curves in the open half-plane $J < 1$. The general case follows by continuity.

For $(b, J) = (0, 0)$ all eigenvalues are real and distinct, so we can order them as $\lambda_0 < \lambda_1 < \dots$. Then each λ_k can be analytically continued along γ for $t \in [0, t_k]$. We denote these analytic continuations by $\lambda_k(t)$. According to a theorem of Shin [19] all but finitely many of the λ_k remain real for $t \in [0, 1]$, and by [10, Thm. 1], all but finitely many λ_k can be analytically continued for $t \in [0, 1]$. Let G be a bounded simply connected neighborhood of γ in the half-plane $\{(b, J) \in \mathbf{R}^2 : J < 1\}$. Then all but finitely many branches of $\lambda(b, J)$ are holomorphic, distinct and real in G . The remaining branches satisfy a minimal algebraic equation of the form

$$\lambda^m + a_{m-1}(b, J)\lambda^{m-1} + \dots + a_m(b, J) = 0, \quad (8)$$

with a_j analytic in G . The zeros of discriminants of these equations form a closed subset K of G .

We claim that all solutions of equations (8) are distinct for all $(b, J) \in \gamma$. Indeed, if any two eigenvalues collide as t increases, then some adjacent eigenvalues $\lambda_j(t)$ and $\lambda_{j+1}(t)$ must collide at some point t_0 . The corresponding eigenfunctions will tend to the same limit as $t \nearrow t_0$. But this is impossible because one of them has no real zeros and another has one.

Thus K does not intersect γ , and there is an analytic continuation of eigenfunctions to $\gamma(1)$.

7 Classification of line complexes

In this section we consider the general case of real asymptotic values a and c . In this case the normalized Nevanlinna function f has asymptotic values $(\infty, 0, c, a, c, 0)$ in S_0, \dots, S_5 . We choose the cell decomposition Φ similar to that in Fig. 1, see Figs. 5, 7, 9.

There are three generic cases:

Case \mathcal{L} . $c < 0 < a$, Case \mathcal{R} . $0 < a < c$, Case \mathcal{E} . $0 < c < a$.

Letters \mathcal{L}, \mathcal{R} and \mathcal{E} stand for “left”, “right” and “even”, the meaning of this notation will be clear later (Figs. 11-15). There are also non-generic cases $a = 0$ and $c = \infty$. Assuming that $a \neq 0$, we classify all possible line complexes.

In all cases \mathcal{L}, \mathcal{R} and \mathcal{E} , we first classify all possible bipartite trees symmetric with respect to the real line, with 6 faces labeled by $\infty, 0, c, a, c, 0$ in this cyclic order, the face labeled ∞ bisected by the positive ray, and satisfying the condition that faces with the same label have disjoint closures. There are three types of such trees shown in Fig. 4. They depend on two integer parameters, $k > 0$ and l , where k is the number of edges between two ramified vertices as shown in Fig. 4. Parameter l takes all integer values, but $|l|$ is the number of edges between ramified vertices as indicated in Fig. 4. We say that a tree has type 0 if $l = 0$, type 1 if $l > 0$ and type 2 if $l < 0$.

For the trees that occurred in section 5 we have $T_n = X_{2n+1,0}$.




Fig. 4. Three types of trees.

Now we consider all cases separately, and argue by the following scheme. First, for a given case of ordering $a, c, 0$ on the real line, and for each tree, we decide whether this tree can come from a line complex, and if it can, we recover the line complex.

To do this, we begin with a ramified vertex v of the tree. Comparing the cyclic order of the labels around this vertex v with the cyclic order around the vertices of the cell decomposition Φ of the sphere we determine whether this vertex of the tree is a \times or \circ , and add the missing edges, step by step. When we come to another ramified vertex, either the cyclic order is correct or not. If it is not correct, the tree does not correspond to a line complex in the considered case. Otherwise, we recover the line complex uniquely.

Each line complex defines a map from an interval to the real spectral locus $Z(\mathbf{R})$ as follows: let $t = c/a$. For each t , we construct a normalized Nevanlinna function f with this given line complex and asymptotic values $a = 1$ and $c = t$. Then b, J, λ can be found from the formula

$$\frac{f'''}{f'} - \frac{3}{2} \left(\frac{f''}{f'} \right)^2 = -2(z^4 - 2bz^2 + 2Jz - \lambda).$$

We write $(b, J, \lambda) = F(\Psi, t)$, where Ψ is the line complex, and call F the Nevanlinna map.

Then we consider possible limits as $a \rightarrow 0$ or $c \rightarrow \infty$ on the spectral locus. This gives degenerations to the intersections of the QES spectral locus with the non-QES spectral locus, and description of these intersections in [13] permits to recover the value of J from the tree in cases 1 and 2. Then we consider the degeneration as $c \rightarrow \infty$ or $c \rightarrow a$ whenever possible, as it was done in [11, 12, 13]. We will conclude that real values of c correspond to integer J and we obtain the parametrization of the whole spectral locus for integer J .

Case \mathcal{L} . $c < 0 < a$, see Figs. 5–6.




Fig. 5. Cell decomposition Φ in case \mathcal{L} .




Fig. 6. $\mathcal{L}_{m,l}$ complexes.

Trees of type 0 (with $l = 0$) are impossible in this case.

Trees $X_{k,l}$ of type 1 (with $l > 0$) are possible in this case if and only if $k = 2m + 1$, $m = 0, 1, \dots$ and $l \geq 1$ is odd. Such line complexes will be called $\mathcal{L}_{m,l}$. When $a \searrow 0$, $t = c/a \rightarrow -\infty$ the corresponding Nevanlinna function has a limit on the QES locus. This limit function has $l - 1$ zeros,

none of them real. It follows that $J = l$ for the limit function. By the first part of Theorem 2 which we proved in section 5, c is real for integer J on the non-QES part of the spectral locus, so we conclude that the whole image of the Nevanlinna map $t \mapsto F(\mathcal{L}_{m,l}, t)$ belongs to $Z_J(\mathbf{R})$ with $J = l$.

In the limit when $c \nearrow 0$, that is $t = c/a \nearrow 0$, we obtain a Nevanlinna function for the harmonic oscillator with m zeros.

Trees of type 2 (with $l < 0$) are possible in this case if and only if $k = 2m + 1$ and $l \leq -1$ is odd. Such line complexes will be called $\mathcal{L}_{m,l}$. The only possible limit on the spectral locus is $c \rightarrow -\infty$. This corresponds to an elementary *second* solution of the differential equation in (3) (solution which is linearly independent of the eigenfunction). These points are marked in Fig. 15. Thus the chart $\mathcal{L}_{m,l}$, $l \leq -1$ corresponds to the chart $\mathcal{L}_{m,-l}$ via Darboux transform. We have $J = l < 0$ in this case.

Case \mathcal{R} . $0 < a < c$, Figs. 7–8.




Fig. 7. Cell decomposition Φ in case \mathcal{R} .




Fig. 8. $\mathcal{R}_{m,l}$ complexes.

Trees of type 0 (with $l = 0$) are impossible in this case.

Trees $X_{k,l}$ of the type 1 (with $l > 0$) are possible with $k = 2m + 1$ and l positive odd. We call the complex $\mathcal{R}_{m,l}$. Degeneration $a \rightarrow 0$, $t = c/a \rightarrow +\infty$ is possible, and the limit belongs to the QES spectral locus. The limit Nevanlinna function has $l - 1$ zeros, so $J = l$ for this limit function. Again, using the first part of Theorem 2, we conclude that $J = l$ on the whole image of the Nevanlinna map $F(\mathcal{R}_{m,l}, \cdot)$. Degeneration $c \rightarrow +\infty$ (that is $t = c/a \rightarrow 0+$) gives a Nevanlinna function for the harmonic oscillator

with $J - 1 + m$ zeros. As $\mathcal{L}_{m,l}$ and $\mathcal{R}_{m,l}$ have common limit of the QES spectral locus, their Nevanlinna images form a single curve. The results in [13] about QES spectral locus together with counting of zeros of degeneration to harmonic oscillator show that $\mathcal{L}_{m,l}$ lies on the left and $\mathcal{R}_{m,l}$ lies on the right from the intersection point with the QES spectral locus. See Figs. 11, 13, where Nevanlinna images of $\mathcal{L}_{m,l}$ and $\mathcal{R}_{m,l}$ are shown with the solid lines, and the QES locus with the dotted line.

Trees $X_{k,l}$ of type 2 (with $l < 0$) give line complexes when $k = 2m + 1$ and l odd negative. We call these complexes $\mathcal{R}_{m,l}$. Degeneration $c \rightarrow \infty$ is possible on the spectral locus. These charts correspond to the charts $\mathcal{R}_{m,-l}$ by Darboux transform.

Thus cases \mathcal{L} and \mathcal{R} and trees of types 1 and 2 cover all cases when J is odd. We conclude from this: a) J is constant on the Nevanlinna images of $\mathcal{L}_{m,l}$ and $\mathcal{R}_{m,l}$, namely $J = l$. b) Even values of J must be covered by the remaining trees from our classification.

Case \mathcal{E} . $0 < c < a$, see Figs. 9–10.




Fig. 9. Cell decomposition Φ in case \mathcal{E} .




Fig. 10. $\mathcal{E}_{m,l}$ complexes.

Trees $X_{k,0}$ of type 0 have parameter $k = 2m + 1$, $m \geq 0$ and we denote the corresponding line complex by $\mathcal{E}_{m,0}$. This line complex represents Nevanlinna functions from section 5 which correspond to $J = 0$.

Trees $X_{k,l}$ of type 2 have $k = 2m + 1$ and l negative, even. We call the corresponding line complex $\mathcal{E}_{m,l}$. No degeneration on the spectral locus is possible. We know that for even J the non-QES spectral locus consists

of graphs of functions. Degeneration as $c \rightarrow 0$ and $c \rightarrow a$ gives a Nevanlinna function for the harmonic oscillator with m zeros. Thus these trees parametrize the whole spectrum for negative even J .

Trees $X_{k,l}$ of type 1 have $k = 2m + 1$ and l positive, even. We call the corresponding complex $\mathcal{E}_{m,l}$. It corresponds to $\mathcal{E}_{m,-l}$ by the Darboux transform. Degeneration as $c \rightarrow 0$ and $c \rightarrow a$ gives a Nevanlinna function for the harmonic oscillator with m and $m + l$ zeros, respectively.

These arguments show that real c and $a \neq 0$ correspond to integer J and that we obtain a parametrization of the whole non-QES spectral locus in this way. This completes the proof of Theorem 2.

The parametrization of the spectral locus for integer J is represented in Figs. 11-15. The symbols of line complexes are written below the corresponding curves. The QES spectral locus is shown with dotted lines. It was parametrized with different cell decompositions (not with line complexes!) in [13]. Symbols $X_{k,l}$ in the figures refer to the charts on the QES locus described in [13].




Fig. 11. $Z_1(\mathbf{R})$.




Fig. 12. $Z_2(\mathbf{R})$.




Fig. 13. $Z_3(\mathbf{R})$.




Fig. 14. $Z_0(\mathbf{R})$.




Fig. 15. $Z_{-1}(R)$ superimposed with $Z_1^{QES}(\mathbf{R})$ (thin dotted line).

References

- [1] I. Bakken, A multiparameter eigenvalue problem in the complex plane, Amer. J. Math. 99 (1977), no. 5, 1015–1044.

- [2] C. Bender and S. Boettcher, Quasi-exactly solvable quartic potential, *J. Phys. A* 31 (1998), no. 14, L273–L277, arXiv:physics/9801007.
- [3] C. Bender and S. Boettcher, Real spectra in non-Hermitian Hamiltonians having PT symmetry, *Phys. Rev. Lett.*, 80 (1998) 52435246.
- [4] C. Bender, D. Brody, J-H. Chen, H. Jones, K. Milton and C. Ogilvie, Equivalence of a complex PT-symmetric quartic Hamiltonian and a Hermitian quartic Hamiltonian with an anomaly, arXiv:hep-th/0605066v2.
- [5] V. Buslaev and V. Grecchi, Equivalence of unstable anharmonic oscillators and double wells, *J. Phys. A*, 26 (1993) 5541–5549.
- [6] M. Crum, Associated Sturm–Liouville systems, *Quart. J. Math.*, 6 (1955) 121–127.
- [7] E. Delabaere, D. T. Trinh, Spectral analysis of the complex cubic oscillator, *J. Phys. A* 33 (2000), no. 48, 87718796.
- [8] P. Dorey, C. Dunning and R. Tateo, The ODE/IM correspondence. *J. Phys. A* 40 (2007), no. 32, R205R283.
- [9] A. Eremenko A. Gabrielov and B. Shapiro, Zeros of eigenfunctions of some anharmonic oscillators, *Ann. Inst. Fourier, Grenoble*, 58, 2 (2008) 603-624.
- [10] A. Eremenko and A. Gabrielov, Analytic continuation of eigenvalues of a quartic oscillator, *Comm. Math. Phys.*, v. 287, No. 2 (2009) 431-457.
- [11] A. Eremenko and A. Gabrielov, Singular perturbation of polynomial potentials in the complex domain with applications to PT-symmetric families, to appear in *Moscow Math. J.*, arXiv:1005.1696.
- [12] A. Eremenko and A. Gabrielov, Quasi-exactly solvable quartic: elementary integrals and asymptotic, arXiv:1104.2305.
- [13] A. Eremenko and A. Gabrielov, Quasi-exactly solvable quartic: real algebraic spectral locus, arXiv:1104.4980.
- [14] J. Gibbons and A. P. Veselov, On the rational monodromy-free potentials with sextic growth, *J. Math. Phys.* 50 (2009), no. 1, 013513, 25 pp.

- [15] A. A. Goldberg and I. V. Ostrovskii, *Distribution of values of meromorphic functions*, AMS, Providence RI, 2008.
- [16] V. Grecchi, M. Maioli and A. Martinez, Padé summability of the cubic oscillator, *J. Math. Phys. A: Math. Theor.* 42 (2009) 425208, 17pp.
- [17] D. Masoero, Y-System and Deformed Thermodynamic Bethe Ansatz, arXiv:1005.1046.
- [18] R. Nevanlinna, Über Riemannsche Flächen mit endlich vielen Windungspunkten, *Acta Math.*, 58 (1932) 295373.
- [19] K. Shin, Eigenvalues of PT-symmetric oscillators with polynomial potentials, *J. Phys. A* 38 (2005), no. 27, 6147–6166.
- [20] K. Shin, On the reality of the eigenvalues for a class of PT-symmetric oscillators, *Comm. Math. Phys.* 229 (2002), no. 3, 543–564.
- [21] Y. Sibuya, *Global theory of a second order linear ordinary differential equation with a polynomial coefficient*, North-Holland, Amsterdam; American Elsevier, NY, 1975.
- [22] L. Volkovyski, Converging sequences of Riemann surfaces, *Mat. Sbornik*, 23 (65) N3 (1948) 361–382.
- [23] J. Zinn-Justin and U. Jentschura, Imaginary cubic perturbation: numerical and analytic study, *J. Phys. A*, 43 (2010) 425301.